

STERNE, KESSLER, GOLDSTEIN & FOX P.L.L.C.

ATTORNEYS AT LAW

1100 NEW YORK AVENUE, N.W., SUITE 600  
WASHINGTON, D.C. 20005-3934

(202) 371-2600

FACSIMILE (202) 371-2540, (202) 371-6566

ROBERT GREENE STERNE  
EDWARD J. KESSLER  
JORGE A. GOLDSTEIN  
SAMUEL L. FOX  
DAVID K.S. CORNWELL  
ROBERT W. ESMOND  
TRACY-GENE G. DURKIN  
MICHAEL A. CIMBALA  
MICHAEL B. RAY  
JAMES E. SOKOHL

ERIC K. STEFFE  
MICHAEL O. LEE  
STEVEN R. LUDWIG  
JOHN M. COVERT\*  
LINDA E. ALCORN  
RAZ E. FLESHNER  
ROBERT C. MILLONIG  
MICHAEL V. MESSINGER  
JUDITH U. KIM  
TIMOTHY J. SHEA, JR.

DONALD R. MCPHAIL  
PATRICK E. GARRETT  
STEPHEN G. WHITESIDE  
NOEL B. WHITLEY  
JEFFREY T. HELVEY\*  
HEIDI L. KRAUS  
JEFFREY R. KURIN  
RAYMOND MILLIEN  
PATRICK D. O'BRIEN  
BRIAN S. ROSENBLUM

LAWRENCE B. BUGAISKY  
CRYSTAL D. SAYLES\*  
EDWARD W. YEE  
ALBERT L. FERRO\*  
DONALD R. BANOWIT\*  
PETER A. JACKMAN\*  
MOLLY A. MCCALL\*  
TERESA U. MEDLER\*  
JEFFREY S. WEAVER\*

DONALD J. FEATHERSTON  
KAREN R. MARKOWICZ\*\*  
GRANT E. REED\*\*  
SUZANNE E. ZISKA\*\*  
BRIAN J. DEL BUONO\*\*  
CAMERON H. TOUSI\*\*  
VINCENT L. CAPUANO\*\*

ROBERT H. BENSON\*  
OF COUNSEL

\*BAR OTHER THAN D.C.  
\*\*REGISTERED PATENT AGENTS

June 29, 1999

WRITER'S DIRECT NUMBER:  
(202) 789-5509

**Box Patent Application**

Assistant Commissioner for Patents  
Washington, D.C. 20231

Re: U.S. Non-Provisional Utility Patent Application under 37 C.F.R. § 1.53(b)  
Appl. No. To be assigned; Filed: Herewith  
(Divisional of Appl. No. 08/741,095; Filed: October 30, 1996)  
For: **Human Tumor Necrosis Factor Receptor-Like 2**  
Inventors: Ni *et al.*  
Our Ref: 1488.0770007/EKS/SGW

Sir:

The following documents are forwarded herewith for appropriate action by the  
U.S. Patent and Trademark Office:

1. USPTO Utility Patent Application Transmittal Form PTO/SB/05;
2. Fee Transmittal (Form PTO/SB/17) (*in duplicate*);
3. 37 C.F.R. § 1.136(a)(3) Authorization to Treat a Reply As Incorporating An  
Extension of Time (*in duplicate*);
4. U.S. Utility Patent Application entitled:

**Human Tumor Necrosis Factor Receptor-Like 2**

06/29/99  
1595 U.S. PTO  
09/340690

06/29/99

06/29/99

1595 U.S. PTO  
09340690-062999

Assistant Commissioner for Patents

June 29, 1999

Page 2

and naming as inventors:

Jian Ni

Craig A. Rosen

Reiner L. Gentz

Sally Doreen Patricia Lyn

Mark Robert Hurle

the application comprising:

a. A specification containing:

(i) 84 pages of description prior to the claims;

(ii) 20 pages of sequence listing;

(iii) 8 pages of claims (Claims 1-26);

(iv) a one (1) page abstract;

b. 27 sheets of drawings: (Figures 1A, 1B, 2, 3, 4A, 4B, 5, 6, 7A, 7B, 8-12, 13A-13C, 14A-14C, 15A-15E, and 16);

5. A copy of the executed Declaration, as filed in U.S. Appl. No. 08/741,095, by Jian Ni, Craig A. Rosen, and Reiner L. Gentz;
6. A copy of the executed Declaration, as filed in U.S. Appl. No. 08/741,095, by Sally Doreen Patricia Lyn and Mark Robert Hurle;
7. First Preliminary Amendment;
8. Second Preliminary Amendment and Submission of Sequence Listing with Exhibits A and B;
9. ATCC Deposit Receipt for Deposit Nos. 97057, 97058, and 97059;
10. Computer Readable Copy of Sequence Listing;
11. Paper Copy of Sequence Listing (Pages 85-106);

Assistant Commissioner for Patents

June 29, 1999

Page 3

12. Our Check No. 24788 in the amount of \$4,906.00 for:


- (i) \$ 760.00 Basic Utility Filing Fee (37 C.F.R. § 1.16(a));
- (ii) \$1,482.00 Independent Claims in Excess of Three (37 C.F.R. § 1.16(b));
- (iii) \$2,664.00 Claims in excess of twenty (37 C.F.R. § 1.16(c)); and

13. Two (2) return postcards.

It is respectfully requested that, of the two attached postcards, one be stamped with the filing date of these documents and returned to our courier, and the other, prepaid postcard, be stamped with the filing date and unofficial application number and returned as soon as possible.

Respectfully submitted,

STERNE, KESSLER, GOLDSTEIN & FOX P.L.L.C.



Stephen G. Whiteside  
Attorney for Applicants  
Registration No. 42,224

Enclosures

Please type a sign(+) inside this box



PTO/SB/05 (2/98)

Approved for use through 09/30/2000 OMB 0651-0032

Patent and Trademark Office U S DEPARTMENT OF COMMERCE

Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it displays a valid OMB control number

<b>UTILITY PATENT APPLICATION TRANSMITTAL</b> (Only for new nonprovisional applications under 37 CFR § 1.53(b))	Attorney Docket No.	1488 0770007/EKS/SGW
	First Inventor or Application Identifier	Jian Ni
	Title	Human Tumor Necrosis Factor Receptor-Like 2
	Express Mail Label No.	

<b>APPLICATION ELEMENTS</b> See MPEP chapter 600 concerning utility patent application contents.	<b>ADDRESS TO</b> Assistant Commissioner for Patents Box Patent Application Washington, DC 20231
<p>1. <input checked="" type="checkbox"/> * Fee Transmittal Form (e.g., PTO/SB/17) (Submit an original, and a duplicate for fee processing)</p> <p>2. <input checked="" type="checkbox"/> Specification [Total Pages <u>113</u> ] (preferred arrangement set forth below) - Descriptive title of the invention - Cross References to Related Applications - Statement Regarding Fed sponsored R &amp; D - Reference to Microfiche Appendix - Background of the invention - Brief Summary of the invention - Brief Description of the Drawings (if filed) - Detailed Description - Claim(s) - Abstract of the Disclosure</p> <p>3. <input checked="" type="checkbox"/> Drawing(s) (35 U.S.C. 113) [Total Sheets <u>27</u> ]</p> <p>4. <input checked="" type="checkbox"/> Oath or Declaration [Total Pages <u>6</u> ] a. <input type="checkbox"/> Newly executed (original or copy) b. <input checked="" type="checkbox"/> Copy from a prior application (37 CFR 1.63(d)) (for continuation/divisional with Box 17 completed) [Note Box 5 below] i. <input type="checkbox"/> <b>DELETION OF INVENTOR(S)</b> Signed statement attached deleting inventor(s) named in the prior application, see 37 CFR §§ 1.63(d)(2) and 1.33(b).</p> <p>5. <input checked="" type="checkbox"/> Incorporation By Reference (useable if Box 4b is checked) The entire disclosure of the prior application, from which a copy of the oath or declaration is supplied under Box 4b, is considered as being part of the disclosure of the accompanying application and is hereby incorporated by reference therein</p> <p>6. <input type="checkbox"/> Microfiche Computer Program (Appendix)</p> <p>7. Nucleotide and/or Amino Acid Sequence Submission (if applicable, all necessary) a. <input checked="" type="checkbox"/> Computer Readable Copy b. <input checked="" type="checkbox"/> Paper Copy (identical to computer copy) c. <input checked="" type="checkbox"/> Statement verifying identity of above copies</p>	
<b>ACCOMPANYING APPLICATION PARTS</b>	
<p>8. <input type="checkbox"/> Assignment Papers (cover sheet &amp; document(s))</p> <p>9. <input type="checkbox"/> 37 CFR 3 73(b) Statement <input type="checkbox"/> Power of Attorney (when there is an assignee)</p> <p>10. <input type="checkbox"/> English Translation Document (if applicable)</p> <p>11. <input type="checkbox"/> Information Disclosure Statement (IDS)/PTO-1449 <input type="checkbox"/> Copies of IDS Citations</p> <p>12. <input checked="" type="checkbox"/> First Preliminary Amendment; Second Preliminary Amendment and Submission of Sequence Listing</p> <p>13. <input checked="" type="checkbox"/> Return Receipt Postcard (MPEP 503) (Should be specifically itemized)</p> <p>14. <input type="checkbox"/> *Small Entity Statement(s) <input type="checkbox"/> Statement filed in prior application, Status still proper and desired (PTO/SB/09-12)</p> <p>15. <input type="checkbox"/> Certified Copy of Priority Document(s) (if foreign priority is claimed)</p> <p>16. <input checked="" type="checkbox"/> Other 37 C.F.R. § 1.136(a)(3) Authorization <input type="checkbox"/> Other.</p> <p><small>*NOTE FOR ITEMS 1 &amp; 14 IN ORDER TO BE ENTITLED TO PAY SMALL ENTITY FEES, A SMALL ENTITY STATEMENT IS REQUIRED (37 C.F.R. § 1.27), EXCEPT IF ONE FILED IN A PRIOR APPLICATION IS RELIED UPON (37 C.F.R. § 1.28)</small></p>	
<p>17. If a CONTINUING APPLICATION, check appropriate box, and supply the requisite information below and in a preliminary amendment</p> <p><input type="checkbox"/> Continuation <input checked="" type="checkbox"/> Divisional <input type="checkbox"/> Continuation-in-Part (CIP) of prior application No: <u>08/741,095</u></p> <p>Prior application information: Examiner <u>Draper, G.</u> Group/Art Unit: <u>1646</u></p>	

<b>18. CORRESPONDENCE ADDRESS</b>					
<input type="checkbox"/> Customer Number or Bar Code Label <input type="checkbox"/> Correspondence address below					
(Insert Customer No. or Attach bar code label here)					
NAME	STERNE, KESSLER, GOLDSTEIN & FOX P.L.L.C.				
	Attorneys at Law				
ADDRESS	Suite 600, 1100 New York Avenue, N.W.				
CITY	Washington	STATE	DC	ZIP CODE	20005-3934
COUNTRY	USA	TELEPHONE	(202) 371-2600	FAX	(202) 371-2540

NAME (Print/Type)	Stephen G. Whiteside	Registration No. (Attorney/Agent)	42,224
SIGNATURE		Date	6/29/99

Burden Hour Statement: this form is estimated to take 0.2 hours to complete. Time will vary depending upon the needs of the individual case. Any comments on the amount of time you are required to complete this form should be sent to the Chief Information Officer, Patent and Trademark Office, Washington, DC 20231. DO NOT SEND FEES OR COMPLETED FORMS TO THIS ADDRESS. SEND TO: Assistant Commissioner for Patents, Washington, DC 20231.



IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re application of:

Ni *et al.*

Appl. No. To be assigned  
(Divisional of Appl. No. 08/741,095,  
Filed: October 30, 1996)

Filed: Herewith

For: **Human Tumor Necrosis Factor  
Receptor-Like 2**

Art Unit: To be assigned

Examiner: To be assigned

Atty. Dkt.: 1488.0770007/EKS/SGW

**First Preliminary Amendment**

Assistant Commissioner for Patents  
Washington, D.C. 20231

Sir:

In advance of prosecution, please amend the captioned application as follows:

***In the Claims:***

Please cancel claims 1-20 and 22-26 without prejudice of or disclaimer to the subject matter therein.

***Remarks***

***I. Status of the Claims***


Claims 1-20 and 22-26 have been canceled. Accordingly, claim 21 is active in the present application. No new matter has been added by this amendment.

**II. Second Preliminary Amendment**

A Second Preliminary Amendment is being filed herewith. In the Second Preliminary Amendment, claim 21 is canceled and new claims 27-174 are added.

Respectfully submitted,

STERNE, KESSLER, GOLDSTEIN & FOX P.L.L.C.



Stephen G. Whiteside  
Attorney for Applicants  
Registration No. 42,224

Date: 6/29/99

1100 New York Avenue, N.W.  
Suite 600  
Washington, D.C. 20005  
(202) 371-2600

666290"05907E60

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re application of:

Ni *et al.*

Appl. No. To be assigned  
(Divisional of Appl. No. 08/741,095,  
Filed: October 30, 1996)

Filed: Herewith

For: **Human Tumor Necrosis Factor  
Receptor-Like 2**

Art Unit: To be assigned

Examiner: To be assigned

Atty. Dkt.: 1488.0770007/EKS/SGW

**Second Preliminary Amendment  
and Submission of Sequence Listing**

Assistant Commissioner for Patents  
Washington, D.C. 20231

Sir:

It is respectfully requested that the following amendments to the specification and claims be entered in advance of substantive examination.

It is not believed that extensions of time or fees for net addition of claims are required beyond those that may otherwise be provided for in documents accompanying this paper. However, if additional extensions of time are necessary to prevent abandonment of this application, then such extensions of time are hereby petitioned under 37 C.F.R. § 1.136(a), and any fees required therefor (including fees for net addition of claims) are hereby authorized to be charged to our Deposit Account No. 19-0036.

***In the Specification:***

Please amend the specification as follows:

On page 1, line 2, please delete the section heading "***Background of the Invention***" and insert therefor as a section heading: --***Cross-Reference to Related Applications***--.

On page 1, lines 3-9, please delete the text of the Cross-Reference to Related Applications section and replace it with the following:

--The present application is a divisional of U.S. Application No. 08/741,095, filed October 30, 1996, which is incorporated herein by reference; said U.S. Application No. 08/741,095 is a continuation-in-part of U.S. Application No. 08/464,595, U.S. Application No. 08/462,962, and U.S. Application No. 08/462,315, each of which was filed June 5, 1995 and are incorporated herein by reference; said U.S. Application Nos. 08/464,595, 08/462,962 and 08/462,315 are each continuations-in-part of PCT/US95/05058, filed April 27, 1995, which is incorporated herein by reference.--

On page 1, line 10, immediately before "Field of the Invention" please insert as a section heading --***Background of the Invention***--.

On page 4, line 6, please delete "Pfefferk" and insert therefor --Pfeffer, K--.

On page 4, line 26, please delete "in bacterial hosts".

On page 9, line 4, please delete "12301 Park Lawn Drive, Rockville, Maryland 20852" and insert therefor --10801 University Blvd., Manassas, VA 20110-2209, USA--.

On page 10, line 4, please delete "The" and insert therefor --All of the--.

On page 17, line 16, after second occurrence of "acids", please insert ---.

On page 21, line 18, please delete "(150 mM NaCl, 15 mM trisodium citrate)" and insert --(750 mM NaCl, 75 mM trisodium citrate)--.

On page 21, line 19, please delete "20 g/ml" and insert therefor --20 µg/ml--.

On page 42, line 15, please delete "Rockville, Maryland" and insert therefor --Manassas, VA--.

On page 75, line 13, please delete "(Rockville, MD)" and insert therefor --(Manassas, VA)--.

065290" 06904E60

After Example 6, on page 84, after line 3, please insert the following example:

**--Example 7**

***Expression Pattern of TNF Receptor Expression in Human Tissue***

Northern blot analysis is carried out to examine the levels of expression of TNF receptor in human tissues. Total cellular RNA samples are isolated with RNazol™ B system (Biotecx Laboratories, Inc., Houston, Tex.). About 10 µg of total RNA isolated from each human tissue specified is separated on 1% agarose gel and blotted onto a nylon filter (Sambrook, Fritsch, and Maniatis, Molecular Cloning, Cold Spring Harbor Press, (1989)). The labeling reaction is done according to the Stratagene Prime-It kit with 50 ng DNA fragment. The labeled DNA is purified with a Select-G-50 column (5 Prime - 3 Prime, Inc. Boulder, CO). The filter is then hybridized with radioactive labeled full length TNF receptor gene at 1,000,000 cpm/ml in 0.5 M NaPO<sub>4</sub>, pH 7.4 and 7% SDS overnight at 65°C. After washing twice at room temperature and twice at 60°C with 0.5 X SSC, 0.1% SDS, the filter is then exposed at -70°C overnight with an intensifying screen.--

Please remove pages 85-104 and insert therefor pages 85-106 included herewith, which contain the amended Sequence Listing for the subject invention. Please renumber the pages which follow accordingly.

***In the Claims:***

Please cancel claim 21 without prejudice or disclaimer of the subject matter therein.

Please add the following new claims:

- 27. An isolated protein comprising amino acids 1 to 245 of SEQ ID NO:26.
28. The isolated protein of claim 27, which comprises amino acids -38 to 245 of SEQ ID NO:26.
29. The isolated protein of claim 27, which is produced by a recombinant host cell.
30. The isolated protein of claim 27, which comprises a heterologous polypeptide.
31. The isolated protein of claim 30, wherein said a heterologous polypeptide comprises the Fc portion of an antibody molecule.
32. A composition comprising the isolated protein of claim 27 and a pharmaceutically acceptable carrier.
33. An isolated protein comprising the mature amino acid sequence encoded by the cDNA clone contained in ATCC Deposit No. 97059.
34. The isolated protein of claim 33, which comprises the complete amino acid sequence encoded by the cDNA clone contained in ATCC Deposit No. 97059.
35. The isolated protein of claim 33, which is produced by a recombinant host cell.

36. The isolated protein of claim 33, which comprises a heterologous polypeptide.
37. The isolated protein of claim 36, wherein said a heterologous polypeptide comprises the Fc portion of an antibody molecule.
38. A composition comprising the isolated protein of claim 33 and a pharmaceutically acceptable carrier.
39. An isolated protein comprising 30 contiguous amino acids of the complete amino acid sequence encoded by the cDNA clone contained in ATCC Deposit No. 97059.
40. The isolated protein of claim 39, which comprises 50 contiguous amino acids of the complete amino acid sequence encoded by the cDNA clone contained in ATCC Deposit No. 97059.
41. The isolated protein of claim 39, which is produced by a recombinant host cell.
42. The isolated protein of claim 39, which comprises a heterologous polypeptide.
43. The isolated protein of claim 42, wherein said a heterologous polypeptide comprises the Fc portion of an antibody molecule.
44. A composition comprising the isolated protein of claim 39 and a pharmaceutically acceptable carrier.

666230-06304650

45. An isolated protein comprising the mature amino acid sequence encoded by the cDNA clone contained in ATCC Deposit No. 97058.
46. The isolated protein of claim 45, which comprises the complete amino acid sequence encoded by the cDNA clone contained in ATCC Deposit No. 97058.
47. The isolated protein of claim 45, which is produced by a recombinant host cell.
48. The isolated protein of claim 45, which comprises a heterologous polypeptide.
49. The isolated protein of claim 48, wherein said a heterologous polypeptide comprises the Fc portion of an antibody molecule.
50. A composition comprising the isolated protein of claim 45 and a pharmaceutically acceptable carrier.
51. An isolated protein comprising 30 contiguous amino acids of the complete amino acid sequence encoded by the cDNA clone contained in ATCC Deposit No. 97058.
52. The isolated protein of claim 51, which comprises 50 contiguous amino acids of the complete amino acid sequence encoded by the cDNA clone contained in ATCC Deposit No. 97058.
53. The isolated protein of claim 51, which is produced by a recombinant host cell.

06904660



54. The isolated protein of claim 51, which comprises a heterologous polypeptide.
55. The isolated protein of claim 54, wherein said a heterologous polypeptide comprises the Fc portion of an antibody molecule.
56. A composition comprising the isolated protein of claim 51 and a pharmaceutically acceptable carrier.
57. An isolated protein comprising the mature amino acid sequence encoded by the cDNA clone contained in ATCC Deposit No. 97057.
58. The isolated protein of claim 57, which comprises the complete amino acid sequence encoded by the cDNA clone contained in ATCC Deposit No. 97057.
59. The isolated protein of claim 57, which is produced by a recombinant host cell.
60. The isolated protein of claim 57, which comprises a heterologous polypeptide.
61. The isolated protein of claim 60, wherein said a heterologous polypeptide comprises the Fc portion of an antibody molecule.
62. A composition comprising the isolated protein of claim 57 and a pharmaceutically acceptable carrier.

63. An isolated protein comprising 30 contiguous amino acids of the complete amino acid sequence encoded by the cDNA clone contained in ATCC Deposit No. 97057.
64. The isolated protein of claim 63, which comprises 50 contiguous amino acids of the complete amino acid sequence encoded by the cDNA clone contained in ATCC Deposit No. 97057.
65. The isolated protein of claim 63, which is produced by a recombinant host cell.
66. The isolated protein of claim 63, which comprises a heterologous polypeptide.
67. The isolated protein of claim 66, wherein said a heterologous polypeptide comprises the Fc portion of an antibody molecule.
68. A composition comprising the isolated protein of claim 63 and a pharmaceutically acceptable carrier.
69. An isolated protein comprising a fragment of the amino acid sequence of SEQ ID NO:26; wherein said protein binds to an antibody having specificity for a polypeptide consisting of the complete amino acid sequence of SEQ ID NO:26.
70. The isolated protein of claim 69, which is produced by a recombinant host cell.
71. The isolated protein of claim 69, which comprises a heterologous polypeptide.

72. The isolated protein of claim 71, wherein said a heterologous polypeptide comprises the Fc portion of an antibody molecule.

73. A composition comprising the isolated protein of claim 69 and a pharmaceutically acceptable carrier.

74. An isolated protein comprising a fragment of an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:5, wherein said protein binds to an antibody having specificity for a polypeptide consisting of the complete amino acid sequence of SEQ ID NO:5; and

(b) the amino acid sequence of SEQ ID NO:8, wherein said protein binds to an antibody having specificity for a polypeptide consisting of the complete amino acid sequence of SEQ ID NO:8.

75. The isolated protein of claim 74, which comprises a fragment of SEQ ID NO:5.

76. The isolated protein of claim 74, which comprises a fragment of SEQ ID NO:8.

77. The isolated protein of claim 74, which is produced by a recombinant host cell.

78. The isolated protein of claim 74, which comprises a heterologous polypeptide.

05904650

79. The isolated protein of claim 78, wherein said a heterologous polypeptide comprises the Fc portion of an antibody molecule.
80. A composition comprising the isolated protein of claim 74 and a pharmaceutically acceptable carrier.
81. An isolated protein comprising amino acids 1 to 162 of SEQ ID NO:26.
82. The isolated protein of claim 81, which comprises amino acids -38 to 162 of SEQ ID NO:26.
83. The isolated protein of claim 81, which is produced by a recombinant host cell.
84. The isolated protein of claim 81, which comprises a heterologous polypeptide.
85. The isolated protein of claim 84, wherein said a heterologous polypeptide comprises the Fc portion of an antibody molecule.
86. A composition comprising the isolated protein of claim 81 and a pharmaceutically acceptable carrier.
87. An isolated protein comprising 30 contiguous amino acids of SEQ ID NO:26.
88. The isolated protein of claim 87, which comprises 50 contiguous amino acids of SEQ ID NO:26.

666230" 06904660

89. The isolated protein of claim 87, which is produced by a recombinant host cell.
90. The isolated protein of claim 87, which comprises a heterologous polypeptide.
91. The isolated protein of claim 90, wherein said a heterologous polypeptide comprises the Fc portion of an antibody molecule.
92. A composition comprising the isolated protein of claim 87 and a pharmaceutically acceptable carrier.
93. An isolated protein comprising an amino acid sequence at least 95% identical to amino acids 1 to 164 of SEQ ID NO:2;  
wherein % identity is determined using the Bestfit program with parameters that calculate % identity over the full length of amino acids 1 to 164 of SEQ ID NO:2 and that allow gaps of up to 5% of the total number of residues in amino acids 1 to 164 of SEQ ID NO:2.
94. The isolated protein of claim 93, comprising amino acids 1 to 164 of SEQ ID NO:2.
95. The isolated protein of claim 93, wherein said amino acid sequence is at least 95% identical to amino acids 1 to 247 of SEQ ID NO:2;  
wherein % identity is determined using the Bestfit program with parameters that calculate % identity over the full length of amino acids 1 to 247 of SEQ ID NO:2 and that

666290 06904660

allow gaps of up to 5% of the total number of residues in amino acids 1 to 247 of SEQ ID NO:2.

96. The isolated protein of claim 95, comprising amino acids 1 to 247 of SEQ ID NO:2.

97. The isolated protein of claim 93, wherein said amino acid sequence is at least 95% identical to amino acids -35 to 247 of SEQ ID NO:2;

wherein % identity is determined using the Bestfit program with parameters that calculate % identity over the full length of amino acids -35 to 247 of SEQ ID NO:2 and that allow gaps of up to 5% of the total number of residues in amino acids -35 to 247 of SEQ ID NO:2.

98. The isolated protein of claim 97, comprising amino acids -35 to 247 of SEQ ID NO:2.

99. The isolated protein of claim 93, wherein said amino acid sequence is at least 95% identical to amino acids -36 to 247 of SEQ ID NO:2;

wherein % identity is determined using the Bestfit program with parameters that calculate % identity over the full length of amino acids -36 to 247 of SEQ ID NO:2 and that allow gaps of up to 5% of the total number of residues in amino acids -36 to 247 of SEQ ID NO:2.

100. The isolated protein of claim 99, comprising amino acids -36 to 247 of SEQ ID NO:2.

101. The isolated protein of claim 93, which is produced by a recombinant host cell.
102. The isolated protein of claim 93, which comprises a heterologous polypeptide.
103. The isolated protein of claim 102, wherein said a heterologous polypeptide comprises the Fc portion of an antibody molecule.
104. A composition comprising the isolated protein of claim 93 and a pharmaceutically acceptable carrier.
105. An isolated protein comprising an amino acid sequence at least 95% identical to the mature amino acid sequence encoded by the cDNA clone in ATCC Deposit No. 97059;  
wherein % identity is determined using the Bestfit program with parameters that calculate % identity over the full length of the mature amino acid sequence encoded by the cDNA clone in ATCC Deposit No. 97059 and that allow gaps of up to 5% of the total number of residues of the mature amino acid sequence encoded by the cDNA clone in ATCC Deposit No. 97059.
106. The isolated protein of claim 105, wherein said amino acid sequence is at least 95% identical to the complete amino acid sequence encoded by the cDNA clone contained in ATCC Deposit No. 97059;  
wherein % identity is determined using the Bestfit program with parameters that calculate % identity over the full length of the complete amino acid sequence encoded by the cDNA clone in ATCC Deposit No. 97059 and that allow gaps of up to 5% of the total number

666230 06904660

of residues of the complete amino acid sequence encoded by the cDNA clone in ATCC  
Deposit No. 97059.

107. The isolated protein of claim 105, which is produced by a recombinant host cell.

108. The isolated protein of claim 105, which comprises a heterologous polypeptide.

109. The isolated protein of claim 108, wherein said a heterologous polypeptide comprises the Fc portion of an antibody molecule.

110. A composition comprising the isolated protein of claim 105 and a pharmaceutically acceptable carrier.

111. An isolated protein comprising an amino acid sequence at least 95% identical to amino acids 1 to 149 of SEQ ID NO:5;

wherein % identity is determined using the Bestfit program with parameters that calculate % identity over the full length of amino acids 1 to 149 of SEQ ID NO:5 and that allow gaps of up to 5% of the total number of residues in amino acids 1 to 149 of SEQ ID NO:5.

112. The isolated protein of claim 111, comprising amino acids 1 to 149 of SEQ ID NO:5.



113. The isolated protein of claim 111, wherein said amino acid sequence is at least 95% identical to amino acids -35 to 149 of SEQ ID NO:5;

wherein % identity is determined using the Bestfit program with parameters that calculate % identity over the full length of amino acids -35 to 149 of SEQ ID NO:5 and that allow gaps of up to 5% of the total number of residues in amino acids -35 to 149 of SEQ ID NO:5.

114. The isolated protein of claim 113, comprising amino acids -35 to 149 of SEQ ID NO:5.

115. The isolated protein of claim 111, wherein said amino acid sequence is at least 95% identical to amino acids -36 to 149 of SEQ ID NO:5;

wherein % identity is determined using the Bestfit program with parameters that calculate % identity over the full length of amino acids -36 to 149 of SEQ ID NO:5 and that allow gaps of up to 5% of the total number of residues in amino acids -36 to 149 of SEQ ID NO:5.

116. The isolated protein of claim 115, comprising amino acids -36 to 149 of SEQ ID NO:5.

117. The isolated protein of claim 111, which is produced by a recombinant host cell.

118. The isolated protein of claim 111, which comprises a heterologous polypeptide.

069301650

119. The isolated protein of claim 118, wherein said a heterologous polypeptide comprises the Fc portion of an antibody molecule.

120. A composition comprising the isolated protein of claim 111 and a pharmaceutically acceptable carrier.

121. An isolated protein comprising an amino acid sequence at least 95% identical to the mature amino acid sequence encoded by the cDNA clone in ATCC Deposit No. 97058; wherein % identity is determined using the Bestfit program with parameters that calculate % identity over the full length of the mature amino acid sequence encoded by the cDNA clone in ATCC Deposit No. 97058 and that allow gaps of up to 5% of the total number of residues of the mature amino acid sequence encoded by the cDNA clone in ATCC Deposit No. 97058.

122. The isolated protein of claim 121, wherein said amino acid sequence is at least 95% identical to the complete amino acid sequence encoded by the cDNA clone contained in ATCC Deposit No. 97058;

wherein % identity is determined using the Bestfit program with parameters that calculate % identity over the full length of the complete amino acid sequence encoded by the cDNA clone in ATCC Deposit No. 97058 and that allow gaps of up to 5% of the total number of residues of the complete amino acid sequence encoded by the cDNA clone in ATCC Deposit No. 97058.

123. The isolated protein of claim 121, which is produced by a recombinant host cell.

124. The isolated protein of claim 121, which comprises a heterologous polypeptide.

125. The isolated protein of claim 124, wherein said a heterologous polypeptide comprises the Fc portion of an antibody molecule.

126. A composition comprising the isolated protein of claim 121 and a pharmaceutically acceptable carrier.

127. An isolated protein comprising an amino acid sequence at least 95% identical to amino acids 2 to 136 of SEQ ID NO:8;

wherein % identity is determined using the Bestfit program with parameters that calculate % identity over the full length of amino acids 2 to 136 of SEQ ID NO:8 and that allow gaps of up to 5% of the total number of residues in amino acids 2 to 136 of SEQ ID NO:8.

128. The isolated protein of claim 127, comprising amino acids 2 to 136 of SEQ ID NO:8.

129. The isolated protein of claim 127, wherein said amino acid sequence is at least 95% identical to amino acids 1 to 136 of SEQ ID NO:8;

555230 06904660

wherein % identity is determined using the Bestfit program with parameters that calculate % identity over the full length of amino acids 1 to 136 of SEQ ID NO:8 and that allow gaps of up to 5% of the total number of residues in amino acids 1 to 136 of SEQ ID NO:8.

130. The isolated protein of claim 129, comprising amino acids 1 to 136 of SEQ ID NO:8.

131. The isolated protein of claim 127, which is produced by a recombinant host cell.

132. The isolated protein of claim 127, which comprises a heterologous polypeptide.

133. The isolated protein of claim 132, wherein said a heterologous polypeptide comprises the Fc portion of an antibody molecule.

134. A composition comprising the isolated protein of claim 127 and a pharmaceutically acceptable carrier.

135. An isolated protein comprising an amino acid sequence at least 95% identical to the mature amino acid sequence encoded by the cDNA clone in ATCC Deposit No. 97057;  
wherein % identity is determined using the Bestfit program with parameters that calculate % identity over the full length of the mature amino acid sequence encoded by the cDNA clone in ATCC Deposit No. 97057 and that allow gaps of up to 5% of the total number

of residues of the mature amino acid sequence encoded by the cDNA clone in ATCC Deposit No. 97057.

136. The isolated protein of claim 135, wherein said amino acid sequence is at least 95% identical to the complete amino acid sequence encoded by the cDNA clone contained in ATCC Deposit No. 97057;

wherein % identity is determined using the Bestfit program with parameters that calculate % identity over the full length of the complete amino acid sequence encoded by the cDNA clone in ATCC Deposit No. 97057 and that allow gaps of up to 5% of the total number of residues of the complete amino acid sequence encoded by the cDNA clone in ATCC Deposit No. 97057.

137. The isolated protein of claim 135, which is produced by a recombinant host cell.

138. The isolated protein of claim 135, which comprises a heterologous polypeptide.

139. The isolated protein of claim 138, wherein said a heterologous polypeptide comprises the Fc portion of an antibody molecule.

140. A composition comprising the isolated protein of claim 135 and a pharmaceutically acceptable carrier.

065290" 06904E60

141. An isolated protein comprising an amino acid sequence selected from the group consisting of:
- (a) amino acids 3 to 34 in SEQ ID NO:2;
  - (b) amino acids 70 to 84 in SEQ ID NO:2;
  - (c) amino acids 106 to 153 in SEQ ID NO:2; and
  - (d) amino acids 240 to 247 in SEQ ID NO:2.
142. The isolated protein of claim 141, wherein said amino acid sequence is (a).
143. The isolated protein of claim 141, wherein said amino acid sequence is (b).
144. The isolated protein of claim 141, wherein said amino acid sequence is (c).
145. The isolated protein of claim 141, wherein said amino acid sequence is (d).
146. The isolated protein of claim 141, which is produced by a recombinant host cell.
147. The isolated protein of claim 141, which comprises a heterologous polypeptide.
148. The isolated protein of claim 147, wherein said a heterologous polypeptide comprises the Fc portion of an antibody molecule.

666250"06904660

149. A composition comprising the isolated protein of claim 141 and a pharmaceutically acceptable carrier.
150. An isolated protein comprising 15 contiguous amino acids of SEQ ID NO:2.
151. The isolated protein of claim 150, which is produced by a recombinant host cell.
152. The isolated protein of claim 150, which comprises a heterologous polypeptide.
153. The isolated protein of claim 152, wherein said a heterologous polypeptide comprises the Fc portion of an antibody molecule.
154. A composition comprising the isolated protein of claim 150 and a pharmaceutically acceptable carrier.
155. An isolated protein comprising an amino acid sequence selected from the group consisting of:
- (a) amino acid 3 to 34 in SEQ ID NO:5;
  - (b) amino acid 63 to 100 in SEQ ID NO:5; and
  - (c) amino acid 135 to 149 in SEQ ID NO:5;
156. The isolated protein of claim 155, wherein said amino acid sequence is (a).

157. The isolated protein of claim 155, wherein said amino acid sequence is (b).
158. The isolated protein of claim 155, wherein said amino acid sequence is (c).
159. The isolated protein of claim 155, which is produced by a recombinant host cell.
160. The isolated protein of claim 155, which comprises a heterologous polypeptide.
161. The isolated protein of claim 160, wherein said a heterologous polypeptide comprises the Fc portion of an antibody molecule.
162. A composition comprising the isolated protein of claim 155 and a pharmaceutically acceptable carrier.
163. An isolated protein comprising an amino acid sequence selected from the group consisting of:
- (a) amino acid 56 to 68 in SEQ ID NO:8; and
  - (b) amino acid 93 to 136 in SEQ ID NO:8.
164. The isolated protein of claim 163, wherein said amino acid sequence is (a).
165. The isolated protein of claim 163, wherein said amino acid sequence is (b).



166. The isolated protein of claim 163, which is produced by a recombinant host cell.

167. The isolated protein of claim 163, which comprises a heterologous polypeptide.

168. The isolated protein of claim 167, wherein said a heterologous polypeptide comprises the Fc portion of an antibody molecule.

169. A composition comprising the isolated protein of claim 163 and a pharmaceutically acceptable carrier.

170. An isolated protein comprising an amino acid sequence encoded by a first polynucleotide which hybridizes to a second polynucleotide consisting of the coding region of SEQ ID NO:25, or the complement thereof, under conditions comprising:

(a) incubating overnight at 65°C in a solution consisting of 0.5M NaPO<sub>4</sub>, pH 7.4 and 7% SDS; and

(b) washing at 60°C in a solution consisting of 0.5 X SSC with 0.1% SDS.

171. The isolated protein of claim 170, which is produced by a recombinant host cell.

172. The isolated protein of claim 170, which comprises a heterologous polypeptide.

09240690-06999

173. The isolated protein of claim 172, wherein said a heterologous polypeptide comprises the Fc portion of an antibody molecule.

174. A composition comprising the isolated protein of claim 170 and a pharmaceutically acceptable carrier.--

### ***Remarks***

After entry of the foregoing amendments, claims 27-174 will be pending in the captioned application with claims 27, 33, 39, 45, 51, 57, 63, 69, 74, 81, 87, 93, 105, 111, 121, 127, 135, 141, 150, 155, 163 and 170 being the independent claims.

#### ***I. The Priority Applications***

The captioned application claims priority benefit of U.S. Application No. 08/741,095, filed October 30, 1996; U.S. Application Nos. 08/464,595, 08/462,962, and 08/462,315, filed June 5, 1995; and PCT/US95/05058 (the '058 application), filed April 27, 1995. Each of these applications have been incorporated by reference into the captioned application. (Specification, page 1, lines 3-7.)

#### ***II. The Amendments and the New Claims are Supported by the Specification***

The specification of the captioned application has been amended to introduce subject matter from the '058 priority application. In particular, the Sequence Listing has been amended by adding SEQ ID NOs:25 and 26. SEQ ID NOs:25 and 26 are identical to SEQ ID NOs:1 and

2 of the '058 application, which was incorporated by reference into the present application. (Specification, page 1, lines 3-9.) Further support for added SEQ ID NOs:25 and 26 appears in the present specification at page 9, lines 20-28, where it is stated that the TR2 receptors of the invention include allelic variants. Allelic variants of TR2 are then described which contain either an adenine or a guanine at nucleotide position 314 of SEQ ID NO:1, resulting in the codon at positions 313-315 of SEQ ID NO:1 encoding either a lysine or arginine residue. The nucleotide sequence shown in SEQ ID NO:25 contains a guanine residue at position 314 whereas SEQ ID NO:1 of the captioned application contains an adenine residue at this location. Further, SEQ ID NO:26 contains an arginine at the -22 position whereas SEQ ID NO:2 of the captioned application contains a lysine residue at the analogous location (-20 position).

In accordance with 37 C.F.R. § 1.821(f), the paper copy of the Sequence Listing and the computer readable copy of the Sequence Listing submitted herewith in the above application are the same.

In accordance with 37 C.F.R. § 1.821(g), this submission includes no new matter.

Example 7 has also been added to the specification. This example is essentially the same as Example 4 of the '058 application which, as noted above, has been incorporated by reference into the captioned application. Thus, the addition of Example 7 into the specification does not introduce new matter.

The amendment of the "Cross-Reference to Related Applications" section of the captioned application is supported by page 1, lines 3-9 of the originally filed specification.

The specification has also been amended to correct an error in the form of the biological materials deposited as ATCC Deposit Nos. 75059, 75058, and 75057. In particular, page 4, lines 26-27, of the specification states that the cDNA clones encoding the TR2 (ATCC Deposit No. 75059), TR2-SV1 (ATCC Deposit No. 75058), and TR2-SV2 (ATCC Deposit No. 75057) polypeptides were deposited in bacterial hosts. However, as evidenced by the attached ATCC deposit receipts, these cDNAs were deposited as plasmid DNAs.

The specification has been amended to include the new address of the American Type Culture Collection (ATCC). The ATCC has recently moved from 12301 Park Lawn Drive, Rockville, MD 20852 to 10801 University Blvd., Manassas, VA 20110-2209. Applicants were advised on May 19, 1998, in a notification published in the Official Gazette, to amend pending applications to refer to the current address of the ATCC. (1210 OFF. GAZ. PAT. OFFICE 74 (May 19, 1998).) The amendments to page 9, lines 3-4; page 42, line 15; and page 75, line 13 are required to incorporate the new address of the ATCC into the specification. No new matter has been added by these amendments.

Page 21, line 18, of the specification has been amended to correct an obvious typographical error in the amount of ingredients listed for 5x SSC (sodium chloride/sodium citrate). An amendment to correct an obvious error does not constitute new matter where one skilled in the art would not only recognize the existence of the error in the specification, but also the appropriate correction. (M.P.E.P. § 2163.07.) Here, the recognition of the typographical error, along with the correction of the error, in the amount of the ingredients listed for 5x SSC, is obvious to one skilled in the art, and, therefore, the correction does not constitute new matter.

5x SSC is a well known solution used in hybridization solutions. (See, e.g., Exhibit A, CURRENT PROTOCOLS IN MOLECULAR BIOLOGY, John Wiley and Sons, N.Y., Supplement 35, page 2.10.7 (1996).) SSC is normally made as a 20x stock solution, and then diluted accordingly for a particular use. Exhibit A shows that a 20x SSC stock solution contains 3M NaCl and 0.3M trisodium citrate. (Exhibit B, CURRENT PROTOCOLS, page A.2.5.) To make a 5x SSC solution, the 20x SSC solution must be diluted by one-fourth. Therefore, a 5x SSC solution contains 750mM NaCl ( $3\text{M} \div 4 = 750\text{mM}$ ) and 75mM trisodium citrate ( $0.3\text{M} \div 4 = 75\text{mM}$ ).

One skilled in the art would have immediately recognized that the amount of ingredients listed in the specification for a 5x SSC solution was incorrect. Rather than describing a 5x SSC solution, made up of 750mM NaCl and 75mM trisodium citrate, the specification inaccurately

listed the ingredients for a 1x SSC solution. Further, the skilled artisan, in recognizing the typographical error, could easily have adjusted the amount of ingredients described in the specification to properly make a 5x SSC solution. Therefore, the correction of this typographical error does not introduce new matter.

Page 21, line 19, of the specification has also been amended to correct an obvious typographical error in the amount of denatured, sheared salmon sperm DNA in the hybridization solution used in an example of "stringent hybridization conditions". The originally filed specification refers to the inclusion of 20 g/ml denatured, sheared salmon sperm DNA but should recite 20 µg/ml.

The inclusion of agents such as salmon sperm DNA as blocking agents is well known in the art. (*See, e.g.,* Exhibit A, CURRENT PROTOCOLS, page 2.10.7.) One skilled in the art would know that salmon sperm DNA is present in hybridization solutions in µg/ml quantities and thus would immediately recognize the above-described typographical error in the specification. *See id.* Further, the skilled artisan, in recognizing the typographical error, could easily have adjusted the amount of ingredients described in the specification to properly included 20 µg/ml denatured, sheared salmon sperm DNA in the hybridization solution. Therefore, the correction of this typographical error does not introduce new matter.

The specification has been amended to include a section heading and to update the priority application data.

Support for claims 27-174 can be found throughout the specification and original claims. In particular, support for claims 27-28, 81-82 and 87-88 can be found, *inter alia*, in the specification in SEQ ID NO:26; at page 6, lines 11-21; at page 9, lines 20-28; at page 34, lines 7-20; and at page 36, lines 16-22. Support for claims 33-34, 45-46 and 57-58 can be found, *inter alia*, in the specification at page 15, line 8, to page 16, line 2. Support for claims 39-40, 51-52 and 63-64 can be found, *inter alia*, in the specification at page 18, lines 24-27, and at page 34, lines 7-20. Support for claims 69 and 74-76 can be found, *inter alia*, in the specification at page

556290-0624650

20, line 20, to page 21, line 11; at page 30, lines 15-19; and at page 35, line 24, to page 36, line 4. Support for claims 93-100, 105-106, 111-116, 121-122, 127-130, and 135-136 can be found, *inter alia*, in the specification in SEQ ID NO:2; at page 6, lines 11-21; at page 15, line 8, to page 16, line 2; and at page 34, line 7, to page 35, line 20. Support for claims 141-145, 155-158, and 163-165 can be found, *inter alia*, in the specification at page 6, line 25, to page 7, line 3; at page 7, lines 12-18; and at page 7, line 25, to page 8, line 2. Support for claim 150 can be found, *inter alia*, in the specification at page 36, lines 14-22. Support for claim 170 can be found, *inter alia*, in the specification in SEQ ID NO:25 and in Example 7 on page 84. Support for claims 29-32, 35-38, 41-44, 47-50, 53-56, 59-62, 65-68, 70-73, 77-80, 83-86, 89-92, 101-104, 107-110, 117-120, 123-126, 131-134, 137-140, 146-149, 151-154, 159-162, 166-169 and 171-174 can be found, *inter alia*, in the specification at page 27, line 16, to page 30, line 13, and at page 53, line 1, to page 54, line 11.

Portions of the TR2 protein comprising amino acids 1-245 and 1-162 of SEQ ID NO:26 are supported, *inter alia*, by SEQ ID NO:26 of the captioned application. More specifically, as evidenced by the negative numbers assigned to the amino acid residues at the N-terminus of the TR2 amino acid sequence shown in SEQ ID NO:26, the leader sequence for the TR2 protein was predicted in the '058 application to consist of 38 amino acids. Thus, one skilled in the art would recognize that the mature TR2 protein would be made up of amino acid 1-245 in SEQ ID NO:26. Further, one skilled in the art would recognize that the predicted extracellular domain of the TR2 protein would be made up of amino acid 1-162 in SEQ ID NO:26. This second point is so because the predicted C terminus of the extracellular domain has remained consistent in all of the priority applications and in the captioned application. As a result, amino acids 1-162 of SEQ ID NO:26 correspond to amino acids 3-164 in SEQ ID NO:2.

In view of the above, claims 27 and 81 are supported by the captioned application.

**Conclusion**

It is respectfully believed that this application is now in condition for substantive examination. Early notice to this effect is respectfully requested.

If the Examiner believes, for any reason, that personal communication will expedite prosecution of this application, the Examiner is invited to telephone the undersigned at the number provided.

Respectfully submitted,

STERNE, KESSLER, GOLDSTEIN & FOX P.L.L.C.



Stephen G. Whiteside  
Attorney for Applicants  
Registration No. 42,224

Date: 6/29/99

1100 New York Avenue, N.W.  
Suite 600  
Washington, D.C. 20005  
(202) 371-2600

# Human Tumor Necrosis Factor Receptor-Like 2

## *Background of the Invention*

The present application is a continuation-in-part of U.S. Application No. 08/464,595, U.S. Application No. 08/462,962, and U.S. Application No 08/462,315, each of which was filed June 5, 1995, the disclosures of which are herein incorporated by reference. U.S. Application Nos. 08/464,595, 08/462,962 and 08/462,315 each claim priority benefit under 35 U.S.C. § 365(c) to PCT/US95/05058, filed April 27, 1995, which disclosure is also herein incorporated by reference

## *Field of the Invention*

The present invention relates to novel members of the Tumor Necrosis Factor (TNF) receptor family. More specifically, isolated nucleic acid molecules are provided encoding a human TNF receptor-related protein, referred to herein as the TR2 receptor of FIG. 1A-1B, having considerable homology to murine CD40. Two different TR2 splice variants, referred to as TR2-SV1 and TR2-SV2, are also provided. TR2 polypeptides are also provided with homology to human type 2 TNF receptor (TNF-RII). Further provided are vectors, host cells and recombinant methods for producing the same. The invention also relates to both the inhibition and enhancement of the activity of TR2 receptor polypeptides and diagnostic methods for detecting TR2 receptor gene expression.

## *Related Art*

Human tumor necrosis factors  $\alpha$  (TNF- $\alpha$ ) and  $\beta$  (TNF- $\beta$  or lymphotoxin) are related members of a broad class of polypeptide mediators, which includes the interferons, interleukins and growth factors, collectively called cytokines (Beutler, B. and Cerami, A., *Annu. Rev. Immunol.*, 7:625-655 (1989)).

Tumor necrosis factor (TNF- $\alpha$  and TNF- $\beta$ ) was originally discovered as a result of its anti-tumor activity, however, now it is recognized as a pleiotropic cytokine playing important roles in a host of biological processes and pathologies



To date, there are ten known members of the TNF-related cytokine family, TNF- $\alpha$ , TNF- $\beta$  (lymphotoxin- $\alpha$ ), LT- $\beta$ , TRAIL and ligands for the Fas receptor, CD30, CD27, CD40, OX40 and 4-1BB receptors. These proteins have conserved C-terminal sequences and variable N-terminal sequences which are often used as membrane anchors, with the exception of TNF- $\beta$ . Both TNF- $\alpha$  and TNF- $\beta$  function as homotrimers when they bind to TNF receptors.

TNF is produced by a number of cell types, including monocytes, fibroblasts, T-cells, natural killer (NK) cells and predominately by activated macrophages. TNF- $\alpha$  has been reported to have a role in the rapid necrosis of tumors, immunostimulation, autoimmune disease, graft rejection, producing an anti-viral response, septic shock, cerebral malaria, cytotoxicity, protection against deleterious effects of ionizing radiation produced during a course of chemotherapy, such as denaturation of enzymes, lipid peroxidation and DNA damage (Nata *et al.*, *J. Immunol.* 136(7):2483 (1987)), growth regulation, vascular endothelium effects and metabolic effects. TNF- $\alpha$  also triggers endothelial cells to secrete various factors, including PAI-1, IL-1, GM-CSF and IL-6 to promote cell proliferation. In addition, TNF- $\alpha$  up-regulates various cell adhesion molecules such as E-Selectin, ICAM-1 and VCAM-1. TNF- $\alpha$  and the Fas ligand have also been shown to induce programmed cell death.

TNF- $\beta$  has many activities, including induction of an antiviral state and tumor necrosis, activation of polymorphonuclear leukocytes, induction of class I major histocompatibility complex antigens on endothelial cells, induction of adhesion molecules on endothelium and growth hormone stimulation (Ruddle, N. and Homer, R., *Prog. Allergy* 40:162-182 (1988)).

Both TNF- $\alpha$  and TNF- $\beta$  are involved in growth regulation and interact with hemopoietic cells at several stages of differentiation, inhibiting proliferation of various types of precursor cells, and inducing proliferation of immature myelomonocytic cells. Porter, A., *Tibtech* 9:158-162 (1991).

Recent studies with "knockout" mice have shown that mice deficient in TNF- $\beta$  production show abnormal development of the peripheral lymphoid organs and morphological changes in spleen architecture (reviewed in Aggarwal *et al.*, *Eur Cytokine Netw*, 7(2):93-124 (1996)). With respect to the lymphoid organs, the popliteal, inguinal, para-aortic, mesenteric, axillary and cervical lymph nodes failed to develop in TNF- $\beta$   $-/-$  mice. In addition, peripheral blood from TNF- $\beta$   $-/-$  mice contained a three fold reduction in white blood cells as compared to normal mice. Peripheral blood from TNF- $\beta$   $-/-$  mice, however, contained four fold more B cells as compared to their normal counterparts. Further, TNF- $\beta$ , in contrast to TNF- $\alpha$  has been shown to induce proliferation of EBV-infected B cells. These results indicate that TNF- $\beta$  is involved in lymphocyte development.

The first step in the induction of the various cellular responses mediated by TNF- $\alpha$  or TNF- $\beta$  is their binding to specific cell surface or soluble receptors. Two distinct TNF receptors of approximately 55-KDa (TNF-RI) and 75-KDa (TNF-RII) have been identified (Hohman *et al.*, *J. Biol. Chem.*, 264:14927-14934 (1989)), and human and mouse cDNAs corresponding to both receptor types have been isolated and characterized (Loetscher *et al.*, *Cell*, 61:351 (1990)). Both TNF-Rs share the typical structure of cell surface receptors including extracellular, transmembrane and intracellular regions.

These molecules exist not only in cell bound forms, but also in soluble forms, consisting of the cleaved extra-cellular domains of the intact receptors (Nophar *et al.*, *EMBO Journal*, 9 (10):3269-76 (1990)) and otherwise intact receptors wherein the transmembrane domain is lacking. The extracellular domains of TNF-RI and TNF-RII share 28% identity and are characterized by four repeated cysteine-rich motifs with significant intersubunit sequence homology. The majority of cell types and tissues appear to express both TNF receptors and both receptors are active in signal transduction, however, they are able to mediate distinct cellular responses. Further, TNF-RII was shown to exclusively mediate human T-cell proliferation by TNF as shown in PCT WO 94/09137.

5 TNF-RI dependent responses include accumulation of C-FOS, IL-6, and  
manganese superoxide dismutase mRNA, prostaglandin E2 synthesis, IL-2  
receptor and MHC class I and II cell surface antigen expression, growth inhibition,  
and cytotoxicity. TNF-RI also triggers second messenger systems such as  
phospholipase A<sub>2</sub>, protein kinase C, phosphatidylcholine-specific phospholipase  
C and sphingomyelinase (Pfefferk *et al.*, *Cell*, 73:457-467 (1993)).

10 Several interferons and other agents have been shown to regulate the  
expression of TNF receptors. Retinoic acid, for example, has been shown to  
induce the production of TNF receptors in some cells type while down regulating  
production in other cells. In addition, TNF- $\alpha$  has been shown effect the  
localization of both types of receptor. TNF- $\alpha$  induces internalization of TNF-RI  
and secretion of TNF-RII (reviewed in Aggarwal *et al.*, *supra*). Thus, the  
production and localization of both TNF-Rs are regulated by a variety of agents.

15 Both the yeast two hybrid system and co-precipitation and purification  
have been used to identify ligands which associate with both types of the TNF-Rs  
(reviewed in Aggarwal *et al.*, *supra* and Vandenabeele *et al.*, *Trends in Cell Biol.*  
5:392-399 (1995)). Several proteins have been identified which interact with the  
cytoplasmic domain of a murine TNF-R. Two of these proteins appear to be  
related to the baculovirus inhibitor of apoptosis, suggesting a direct role for TNF-  
20 R in the regulation of programmed cell death.

### *Summary of the Invention*

25 The present invention provides isolated nucleic acid molecules comprising  
polynucleotides encoding a TR2 receptor and splice variants thereof having the  
amino acid sequences shown in FIG. 1A-1B (SEQ ID NO:2), FIG. 4A-4B (SEQ  
ID NO:5) and FIG. 7A-7B (SEQ ID NO:8) or the amino acid sequence encoded  
by the cDNA clone encoding the TR2 receptors deposited in bacterial hosts as  
ATCC Deposit Numbers 97059, 97058 and 97057 on February 13, 1995. The  
present invention also relates to recombinant vectors, which include the isolated

nucleic acid molecules of the present invention, and to host cells containing the recombinant vectors, as well as to methods of making such vectors and host cells and for using them for production of TR2 polypeptides or peptides by recombinant techniques.

5           The invention further provides isolated TR2 polypeptides having amino acid sequences encoded by the polynucleotides described herein.

10           The present invention also provides a screening method for identifying compounds capable of enhancing or inhibiting a cellular response induced by TR2 receptors, which involves contacting cells which express TR2 receptors with the candidate compound, assaying a cellular response, and comparing the cellular response to a standard cellular response, the standard being assayed when contact is made in absence of the candidate compound; whereby, an increased cellular response over the standard indicates that the compound is an agonist and a decreased cellular response over the standard indicates that the compound is an antagonist.

15           In another aspect, a screening assay for agonists and antagonists is provided which involves determining the effect a candidate compound has on the binding of cellular ligands to TR2 receptors. In particular, the method involves contacting TR2 receptors with a ligand polypeptide and a candidate compound and determining whether ligand binding to the TR2 receptors is increased or decreased due to the presence of the candidate compound.

20           The invention further provides a diagnostic method useful during diagnosis or prognosis of a disease states resulting from aberrant cell proliferation due to alterations in TR2 receptor expression.

25           An additional aspect of the invention is related to a method for treating an individual in need of an increased level of a TR2 receptor activity in the body comprising administering to such an individual a composition comprising a therapeutically effective amount of isolated TR2 polypeptides of the invention or an agonist thereof.

A still further aspect of the invention is related to a method for treating an individual in need of a decreased level of a TR2 receptor activity in the body comprising, administering to such an individual a composition comprising a therapeutically effective amount of a TR2 receptor antagonist.

5 The invention additionally provides soluble forms of the polypeptides of the present invention. Soluble peptides are defined by amino acid sequences wherein the sequence comprises the polypeptide sequences lacking a transmembrane domain. Such soluble forms of the TR2 receptors are useful as antagonists of the membrane bound forms of the receptors.

### *Brief Description of the Figures*

10 FIG. 1A-1B shows the nucleotide (SEQ ID NO:1) and deduced amino acid (SEQ ID NO:2) sequences of a TR2 receptor. The protein has a predicted leader sequence of about 36 amino acid residues (underlined) (amino acid residues -36 to -1 in SEQ ID NO:2) and a deduced molecular weight of about 30,417 kDa. It is further predicted that amino acid residues from about 37 to about 200 (amino acid residues 1 to 164 in SEQ ID NO:2) constitute the extracellular domain; from about 201 to about 225 (amino acid residues 165 to 189 in SEQ ID NO:2) the transmembrane domain (underlined); and from about 226 to about 283 (amino acid residues 190 to 247 in SEQ ID NO:2) the intracellular domain. Two potential asparagine-linked glycosylation sites are located at amino acid positions 110 and 173 (amino acid residues 74 to 137 in SEQ ID NO:2).

20 FIG. 2 shows the regions of similarity between the amino acid sequences of the TR2 receptor protein of FIG. 1A-1B and a murine CD40 protein (SEQ ID NO:3).

25 FIG. 3 shows an analysis of the TR2 receptor amino acid sequence of FIG. 1A-1B. Alpha, beta, turn and coil regions; hydrophilicity and hydrophobicity; amphipathic regions; flexible regions; antigenic index and surface probability are shown. In the "Antigenic Index - Jameson-Wolf" graph, amino acid residues 39

to 70, 106 to 120, 142 to 189 and 276 to 283 in FIG. 1A-1B (amino acid residues 3 to 34, 70 to 84, 106 to 153 and 240 to 247 in SEQ ID NO:2) correspond to the shown highly antigenic regions of the TR2 receptor protein.

FIG. 4A-4B shows the nucleotide (SEQ ID NO:4) and deduced amino acid (SEQ ID NO:5) sequences of the TR2-SV1 receptor. The protein has a predicted leader sequence of about 36 amino acid residues (underlined) (amino acid residues -36 to -1 in SEQ ID NO:5) and a deduced molecular weight of about 19.5 kDa.

FIG. 5 shows the regions of similarity between the amino acid sequences of the full-length TR2-SV1 receptor protein and a human type 2 TNF receptor (SEQ ID NO:6).

FIG. 6 shows an analysis of the TR2-SV1 receptor amino acid sequence. Alpha, beta, turn and coil regions; hydrophilicity and hydrophobicity; amphipathic regions; flexible regions; antigenic index and surface probability are shown. In the "Antigenic Index - Jameson-Wolf" graph, amino acid residues 39 to 70, 99 to 136 and 171 to 185 in FIG. 4A-4B (amino acid residues 3 to 34, 63 to 100 and 135 to 149 in SEQ ID NO:5) correspond to the shown highly antigenic regions of the TR2-SV1 receptor protein.

FIG. 7A-7B shows the nucleotide (SEQ ID NO:7) and deduced amino acid (SEQ ID NO:8) sequences of the TR2-SV2 receptor. This protein lacks a putative leader sequence and has a deduced molecular weight of about 14 kDa.

FIG. 8 shows the regions of similarity between the amino acid sequences of the TR2-SV2 receptor protein and a human type 2 TNF receptor (SEQ ID NO:9).

FIG. 9 shows an analysis of the TR2-SV2 receptor amino acid sequence. Alpha, beta, turn and coil regions; hydrophilicity and hydrophobicity; amphipathic regions; flexible regions; antigenic index and surface probability are shown. In the "Antigenic Index - Jameson-Wolf" graph, amino acid residues 56 to 68 and 93 to

136 in FIG. 7A-7B (SEQ ID NO:8) correspond to the shown highly antigenic regions of the TR2-SV2 receptor protein.

FIG. 10 shows the regions of similarity between the amino acid sequences of the TR2 receptor protein of FIG. 1A-1B and the TR2-SV1 receptor protein of FIG. 4A-4B.

FIG. 11 shows the regions of similarity between the amino acid sequences of the TR2 receptor protein of FIG. 1A-1B and the TR2-SV2 receptor protein of FIG. 7A-7B.

FIG. 12 shows the regions of similarity between the amino acid sequences of the TR2-SV1 and the TR2-SV2 receptor proteins.

FIG. 13A-13C shows the regions of similarity between the nucleotide sequences encoding the TR2 receptor protein of FIG. 1A-1B and the TR2-SV1 receptor protein of FIG. 4A-4B.

FIG. 14A-14C shows the regions of similarity between the nucleotide sequences encoding the TR2 receptor protein of FIG. 1A-1B and the TR2-SV2 receptor protein of FIG. 7A-7B.

FIG. 15A-15E shows the regions of similarity between the nucleotide sequences encoding the TR2-SV1 and the TR2-SV2 receptor proteins.

FIG. 16 shows an alignment of the amino acid sequence of the TR2 receptor of FIG. 1A-1B (SEQ ID NO:2) with other TNFR family members. The amino acid sequence of TR2 was aligned with those of TNFR-I (SEQ ID NO:10), TNFR-II (SEQ ID NO:11), CD40 (SEQ ID NO:12) and 4-1BB (SEQ ID NO:13) on the basis of sequence homology and conserved cysteine residues.

### ***Detailed Description of the Preferred Embodiments***

The present invention provides isolated nucleic acid molecules comprising polynucleotides encoding a TR2 polypeptide (FIG. 1A-1B (SEQ ID NO:2)) and splice variants thereof, TR2-SV1 (FIG. 4A-4B (SEQ ID NO:5)) and TR2-SV2 (FIG. 7A-7B (SEQ ID NO:8)), the amino acid sequences of which were

determined by sequencing cloned cDNAs. The TR2 protein shown in FIG. 1A-1B shares sequence homology with the murine CD40 receptor (FIG. 2 (SEQ ID NO:3)). On February 13, 1995 a deposit was made at the American Type Culture Collection, 12301 Parklawn Drive, Rockville, Maryland 20852, and given accession number 97059. The nucleotide sequence shown in FIG. 1A-1B (SEQ ID NO:1) was obtained by sequencing a cDNA clone (Clone ID HLHAB49) containing the same amino acid coding sequences as the clone in ATCC Accession No. 97059 with minor deviation. The cDNA sequence shown in FIG. 1A-1B (SEQ ID NO:1) differs from that of the ATCC deposit in the 5' and 3' noncoding nucleotide sequences and three nucleotides.

The clone deposited in ATCC Accession No. 97059 contains 8 nucleotides 5' to the TR2 initiation codon and 21 nucleotides 3' to the TR2 stop codon. In contrast, the TR2 cDNA sequence of HLHAB49, shown in FIG. 1A-1B (SEQ ID NO:1), contains considerably longer non-coding nucleotides sequence on both the 5' and 3' ends of the TR2 coding sequences. Further, the TR2 receptor nucleotide sequence shown in FIG. 1A-1B (SEQ ID NO:1) contains an adenine at nucleotide 314, a cytosine at nucleotide 386, and a cytosine at nucleotide 627. In contrast, the clone of ATCC Accession No. 97059 contains a guanine at nucleotide 314, a thymine at nucleotide 386, and a thymine at nucleotide 627.

The TR2 receptors of the present invention include several allelic variants containing alterations in at least these three nucleotides and two amino acids. Nucleotide sequence variants which have been identified include either guanine or adenine at nucleotide 314, thymine or cytosine at nucleotide 386, and thymine or cytosine at nucleotide 627 shown in FIG. 1A-1B (SEQ ID NO:1). While the identified alteration at nucleotide 627 is silent, the alteration at nucleotide 386 results in the codon at nucleotides 385 to 387 encoding either serine or phenylalanine and the alteration at nucleotide 314 results in the codon at nucleotides 313 to 315 encoding either lysine or arginine.



5 The nucleotide sequences shown in FIG. 4A-4B (SEQ ID NO:4) and FIG. 7A-7B (SEQ ID NO:7) were also obtained by sequencing cDNA clones deposited on February 13, 1995 at the American Type Culture Collection and given accession numbers 97058 (TR2-SV1) and 97057 (TR2-SV2), respectively. The deposited clones are contained in the pBluescript SK(-) plasmid (Stratagene, LaJolla, CA).

10 As used herein the phrase "splice variant" refers to cDNA molecules produced from a RNA molecules initially transcribed from the same genomic DNA sequence which have undergone alternative RNA splicing. Alternative RNA splicing occurs when a primary RNA transcript undergoes splicing, generally for the removal of introns, which results in the production of more than one mRNA molecule each of which may encode different amino acid sequences. The term "splice variant" also refers to the proteins encoded by the above cDNA molecules.

15 As used herein, "TR2 proteins", "TR2 receptors", "TR2 receptor proteins" and "TR2 polypeptides" refer to all proteins resulting from the alternate splicing of the genomic DNA sequences encoding proteins having regions of amino acid sequence identity and receptor activity which correspond to the proteins shown in FIG. 1A-1B (SEQ ID NO:2), FIG. 4A-4B (SEQ ID NO:5) or FIG. 7A-7B (SEQ ID NO:8). The TR2 protein shown in FIG. 1A-1B, the TR2-SV1 protein shown FIG. 4A-4B and the TR2-SV2 protein shown in FIG. 7A-7B are examples of such receptor proteins.

### *Nucleic Acid Molecules*

20 Unless otherwise indicated, all nucleotide sequences determined by sequencing a DNA molecule herein were determined using an automated DNA sequencer (such as the Model 373 from Applied Biosystems, Inc.), and all amino acid sequences of polypeptides encoded by DNA molecules determined herein were predicted by translation of a DNA sequence determined as above. Therefore, as is known in the art for any DNA sequence determined by this

25

automated approach, any nucleotide sequence determined herein may contain some errors. Nucleotide sequences determined by automation are typically at least about 90% identical, more typically at least about 95% to at least about 99.9% identical to the actual nucleotide sequence of the sequenced DNA molecule. The actual sequence can be more precisely determined by other approaches including manual DNA sequencing methods well known in the art. As is also known in the art, a single insertion or deletion in a determined nucleotide sequence compared to the actual sequence will cause a frame shift in translation of the nucleotide sequence such that the predicted amino acid sequence encoded by a determined nucleotide sequence will be completely different from the amino acid sequence actually encoded by the sequenced DNA molecule, beginning at the point of such an insertion or deletion

Using the information provided herein, such as the nucleotide sequence in FIG. 1A-1B, FIG. 4A-4B or FIG. 7A-7B, nucleic acid molecules of the present invention encoding TR2 polypeptides may be obtained using standard cloning and screening procedures, such as those used for cloning cDNAs using mRNA as starting material. Illustrative of the invention, the nucleic acid molecule described in FIG. 1A-1B (SEQ ID NO:1) was discovered in a cDNA library derived from activated human T-lymphocytes. The nucleic acid molecules described in FIG. 4A-4B (SEQ ID NO:4) and FIG. 7A-7B (SEQ ID NO:7) were discovered in cDNAs library derived from human fetal heart and human stimulated monocytes, respectively.

As described in Example 6, TR2 mRNA was detected in numerous tissues including lung, spleen and thymus and may be ubiquitously expressed in human cells. TR2 RNA was also found to be expressed in B lymphocytes (CD19<sup>+</sup>), both CD4<sup>+</sup> (T<sub>H1</sub> and T<sub>H2</sub> clones) and CD8<sup>+</sup> T lymphocytes, monocytes and endothelial cells.

As also noted in Example 6, the production of TR2 mRNA was inducible in MG 63 cells by TNF $\alpha$ . Further, the accumulation of TR2 mRNA was observed

in HL60, U937 and THP1 cells upon PMA or DMSO treatment. PMA and DMSO are agents known to induce differentiation of these three cell types.

The determined nucleotide sequence of the TR2 cDNA of FIG. 1A-1B (SEQ ID NO:1) contains an open reading frame encoding a protein of about 283 amino acid residues, with a predicted leader sequence of about 36 amino acid residues, and a deduced molecular weight of about 30,417 kDa. The amino acid sequence of the predicted mature TR2 receptor is shown in FIG. 1A-1B from amino acid residue about 37 to residue about 283 (amino acid residues 1 to 247 in SEQ ID NO:2). As noted in Example 6, the location of the leader sequence cleavage site was confirmed for a TR2-Fc fusion protein and found to be between amino acids 36 and 37 shown in FIG. 1A-1B (amino acid residues -1 to 1 in SEQ ID NO:2). The TR2 protein shown in FIG. 1A-1B (SEQ ID NO:2) is about 29% identical and about 47% similar to the murine CD40 protein shown in SEQ ID NO:3 (see FIG. 2).

Similarly, the determined cDNA nucleotide sequences of the TR2-SV1 splice variant of TR2 (FIG. 4A-4B (SEQ ID NO:4)) contains an open reading frame encoding a protein of about 185 amino acid residues, with a predicted leader sequence of about 36 amino acid residues, and a deduced molecular weight of about 19.5 kDa. The amino acid sequence of the predicted mature TR2-SV1 receptor is shown in FIG. 4A-4B (SEQ ID NO:5) from amino acid residue about 37 to residue about 185 (amino acid residues 1 to 149 in (SEQ ID NO:5). The TR2-SV1 protein shown in FIG. 4A-4B (SEQ ID NO:5) is about 25% identical and about 48% similar to the human type 2 TNF receptor protein shown in SEQ ID NO:6 (see FIG. 5).

The determined cDNA nucleotide sequences of the TR2-SV2 splice variant of TR2 (FIG. 7A-7B (SEQ ID NO:7)) contains an open reading frame encoding a protein of about 136 amino acid residues, without a predicted leader sequence, and a deduced molecular weight of about 14 kDa. The amino acid sequence of the predicted TR2-SV2 receptor is shown in FIG. 7A-7B (SEQ ID

NO:8) from amino acid residue about 1 to residue about 136. The TR2-SV2 protein shown in FIG. 7A-7B (SEQ ID NO:8) is about 27% identical and about 45% similar to the human type 2 TNF receptor protein shown in SEQ ID NO: 9 (see FIG. 8).

5 A comparison of both the nucleotide and amino acid sequences of the TR2, TR2-SV1 and TR2-SV2 receptor proteins shown in FIG. 1A-1B, FIG. 4A-4B and FIG. 7A-7B shows several regions of near identity. While the amino acid sequence of the TR2 receptor protein, shown in FIG. 1A-1B (SEQ ID NO:2), is about 60% identical and about 73% similar to the amino acid sequence of the TR2-SV1 receptor protein, shown in FIG. 4A-4B (SEQ ID NO:5), in approximately the first one hundred amino acids of their respective sequences the two proteins differ in one location (FIG. 10).

10 Similarly, the amino acid sequence of the TR2 receptor protein of FIG. 1A-1B (SEQ ID NO:2) is about 60% identical and about 71% similar to the amino acid sequence of the TR2-SV2 receptor protein, shown in FIG. 7A-7B (SEQ ID NO:8); however, the two proteins are almost identical over a 60 amino acid stretch in the central portion of the TR2-SV2 protein (FIG. 11).

15 In contrast, the TR2-SV1 and TR2-SV2 proteins are only about 20% identical and about 38% similar at the amino acid level to each other. Unlike the comparisons of either of these proteins to the TR2 protein shown in FIG. 1A-1B (SEQ ID NO:2), these proteins share their homology over the entire 136 amino acid sequence of the TR2-SV2 protein (FIG. 12).

20 With respect to their nucleotide sequences of the cDNAs encoding the disclosed TR2 proteins, a comparison of these sequences indicates that the TR2 cDNAs share large regions of near identity at the nucleic acid level (FIG. 13A-13C, FIG. 14A-13C and FIG. 15A-15E). The cDNA sequences encoding the TR2 and TR2-SV1 proteins, for example, share large regions of near identity in their nucleotide sequences which encode both the N termini of the respective proteins and their 5' and 3' noncoding regions (FIG. 13A-13C). Further, the

25

nucleotide sequences of the cDNAs encoding the TR2-SV1 and TR2-SV2 proteins share considerable homology but this identity is limited to their 3' regions well beyond their respective coding sequences (FIG. 15A-15E).

Such regions of near identity between two different cDNA sequences, when maintained over an extended stretch of sequence, indicates to one skilled in the art that the respective molecules were originally transcribed from the same genomic DNA sequence. One skilled in the art would further recognize that, since more than one codon can encode the same amino acid, identity between two proteins at the amino acid level does not necessarily mean that the DNA sequences encoding those proteins will share similar regions of identity. The above data indicates that the TR2 receptors of the present invention are transcribed from a single genomic DNA sequence and represent multiple splice variants of one initial RNA transcript.

Related proteins which are produced from alternately spliced RNA, referred to as splice variants, are known in the art. The transcript of the *src* gene, for example, undergoes alternate RNA splicing to produce cell type specific products. In most cells the Src protein consists of 533 amino acids while in nerve cells an additional short exon is included in the mRNA resulting in a protein of 539 amino acids. See Alberts, B. *et al.*, MOLECULAR BIOLOGY OF THE CELL (3rd Edition, Garland Publishing, Inc., 1994), 455. Similarly, sex specific mRNA transcripts have been identified in *Drosophila* where alternate mRNA splicing results in a protein named Dsx which is approximately 550 amino acids in length in males and 430 amino acids in length in females. These two splice variant proteins share a common core sequence of about 400 amino acids. See *id.* at 457.

In the present instance, the TR2 receptor protein shown in FIG. 1A-1B (SEQ ID NO:2) is believed to be the full-length polypeptide encoded by the RNA from which the TR2 receptor proteins are translated. The RNA encoding the TR2-SV1 splice variant shown in FIG. 4A-4B (SEQ ID NO:5) is believed to contain an insertion in the region encoding amino acid residue 102 of the amino

acid sequence shown in FIG. 1A-1B and a deletion in the region encoding amino acid residue 184 of the amino acid sequence shown in FIG. 1A-1B. The RNA encoding the TR2-SV2 splice variant shown in FIG. 7A-7B is believed to begin with the nucleotide sequence encoding amino acid residue 102 of the amino acid sequence shown in FIG. 1A-1B and contain insertions in the regions encoding amino acid residues 184 and 243 of the amino acid sequence shown in FIG. 1A-1B.

As indicated, the present invention also provides the mature forms of the TR2 receptors of the present invention. According to the signal hypothesis, proteins secreted by mammalian cells have a signal or secretory leader sequence which is cleaved from the mature protein once export of the growing protein chain across the rough endoplasmic reticulum has been initiated. Most mammalian cells and even insect cells cleave secreted proteins with the same specificity. However, in some cases, cleavage of a secreted protein is not entirely uniform, which results in two or more mature species on the protein. Further, it has long been known that the cleavage specificity of a secreted protein is ultimately determined by the primary structure of the complete protein, that is, it is inherent in the amino acid sequence of the polypeptide. Therefore, the present invention provides nucleotide sequences encoding mature TR2 polypeptides having the amino acid sequences encoded by the cDNA clones contained in the host identified as ATCC Deposit Numbers 97059 and 97058 and as shown in FIG. 1A-1B (SEQ ID NO:2) and FIG. 4A-4B (SEQ ID NO:5). By the mature TR2 polypeptides having the amino acid sequences encoded by the cDNA clones contained in the host identified as ATCC Deposit Numbers 97059 and 97058 is meant the mature form(s) of the TR2 receptors produced by expression in a mammalian cell (e.g., COS cells, as described below) of the complete open reading frame encoded by the human DNA sequence of the clone contained in the vector in the deposited host

The invention also provides nucleic acid sequences encoding the TR2-SV2 receptor protein of FIG. 7A-7B (SEQ ID NO:8), having the amino acid sequence

encoded by the cDNA clone contained in ATCC Deposit Number 97057, which does not contain a secretory leader sequence.

Methods for predicting whether a protein has a secretory leader as well as the cleavage point for that leader sequence are available. For instance, the methods of McGeoch (*Virus Res.* 3:271-286 (1985)) and von Heinje (*Nucleic Acids Res.* 14:4683-4690 (1986)) can be used. The accuracy of predicting the cleavage points of known mammalian secretory proteins for each of these methods is in the range of 75-80%. von Heinje, *supra*. However, the two methods do not always produce the same predicted cleavage point(s) for a given protein.

In the present case, the predicted amino acid sequences of the complete TR2 polypeptides shown in FIG. 1A-1B (SEQ ID NO:2), FIG. 4A-4B (SEQ ID NO:5) and FIG. 7A-7B (SEQ ID NO:8) were analyzed by a computer program ("PSORT") (K. Nakai and M. Kanehisa, *Genomics* 14:897-911 (1992)), which is an expert system for predicting the cellular location of a protein based on the amino acid sequence. As part of this computational prediction of localization, the methods of McGeoch and von Heinje are incorporated. The analysis by the PSORT program predicted the cleavage sites between amino acids -1 and 1 in SEQ ID NO:2 and SEQ ID NO:5. Thereafter, the complete amino acid sequences were further analyzed by visual inspection, applying a simple form of the (-1,-3) rule of von Heine. von Heinje, *supra*. Thus, the leader sequences for the TR2 protein shown in SEQ ID NO.2 and the TR2-SV1 protein are predicted to consist of amino acid residues -36 to -1 in both SEQ ID NO:2 and SEQ ID NO:5, while the predicted mature TR2 proteins consist of amino acid residues 1 to 247 for the TR2 protein shown in SEQ ID NO:2 and residues 1 to 149 for the TR2-SV1 protein shown in SEQ ID NO:5.

As noted in Example 6, the cleavage site of the leader sequence of a TR2-Fc fusion protein was confirmed using amino acid analysis of the expressed fusion protein. This fusion protein was found to begin at amino acid 37, which corresponds to amino acid 1 in SEQ ID NO:2 and SEQ ID NO:5, indicating that

the cleavage site of the leader sequence is between amino acids 36 and 37 in this protein (corresponding to amino acid residues -1 to 1 in SEQ ID NO:2 and SEQ ID NO:5).

As one of ordinary skill would appreciate, however, due to the possibilities of sequencing errors, as well as the variability of cleavage sites for leaders in different known proteins, the TR2 receptor polypeptide encoded by the cDNA of ATCC Deposit Number 97059 comprises about 283 amino acids, but may be anywhere in the range of 250 to 316 amino acids; and the leader sequence of this protein is about 36 amino acids, but may be anywhere in the range of about 30 to about 42 amino acids. Similarly, the TR2-SV1 receptor polypeptide encoded by the cDNA of ATCC Deposit Number 97058 comprises about 185 amino acids, but may be anywhere in the range of 163-207 amino acids; and the leader sequence of this protein is about 36 amino acids, but may be anywhere in the range of about 30 to about 42 amino acids. Further, the TR2-SV2 receptor polypeptide encoded by the cDNA of ATCC Deposit Number 97057 comprises about 136 amino acids, but may be anywhere in the range of 120-152 amino acids

As indicated, nucleic acid molecules of the present invention may be in the form of RNA, such as mRNA, or in the form of DNA, including, for instance, cDNA and genomic DNA obtained by cloning or produced synthetically. The DNA may be double-stranded or single-stranded. Single-stranded DNA or RNA may be the coding strand, also known as the sense strand, or it may be the non-coding strand, also referred to as the anti-sense strand.

By "isolated" nucleic acid molecule(s) is intended a nucleic acid molecule, DNA or RNA, which has been removed from its native environment. For example, recombinant DNA molecules contained in a vector are considered isolated for the purposes of the present invention. Further examples of isolated DNA molecules include recombinant DNA molecules maintained in heterologous host cells or purified (partially or substantially) DNA molecules in solution. Isolated RNA molecules include *in vivo* or *in vitro* RNA transcripts of the DNA molecules of the



present invention. Isolated nucleic acid molecules according to the present invention further include such molecules produced synthetically.

Isolated nucleic acid molecules of the present invention include DNA molecules comprising an open reading frame (ORF) shown in FIG. 1A-1B (SEQ ID NO:1); DNA molecules comprising the coding sequence for the mature TR2 receptor shown in FIG. 1A-1B (SEQ ID NO:2) (last 247 amino acids); and DNA molecules which comprise a sequence substantially different from those described above but which, due to the degeneracy of the genetic code, still encode the TR2 receptor protein shown in FIG. 1A-1B (SEQ ID NO:2). Of course, the genetic code is well known in the art. Thus, it would be routine for one skilled in the art to generate such degenerate variants.

Similarly, isolated nucleic acid molecules of the present invention include DNA molecules comprising an open reading frame (ORF) shown in FIG. 4A-4B (SEQ ID NO:4); DNA molecules comprising the coding sequence for the mature TR2-SV1 receptor shown in FIG. 4A-4B (SEQ ID NO:5) (last 149 amino acids); and DNA molecules which comprise a sequence substantially different from those described above but which, due to the degeneracy of the genetic code, still encode the TR2-SV1 receptor.

Further, isolated nucleic acid molecules of the present invention include DNA molecules comprising an open reading frame (ORF) shown in FIG. 7A-7B (SEQ ID NO:7) and DNA molecules which comprise a sequence substantially different from those described above but which, due to the degeneracy of the genetic code, still encode the TR2-SV2 receptor.

In another aspect, the invention provides isolated nucleic acid molecules encoding the TR2, TR2-SV1 and TR2-SV2 polypeptides having the amino acid sequences encoded by the cDNA clones contained in the plasmid deposited as ATCC Deposit No. 97059, 97058 and 97057, respectively, on February 13, 1995. In a further embodiment, these nucleic acid molecules will encode a mature polypeptide or the full-length polypeptide lacking the N-terminal methionine. The

invention further provides isolated nucleic acid molecules having the nucleotide sequences shown in FIG. 1A-1B (SEQ ID NO:1), FIG. 4A-4B (SEQ ID NO:4), and FIG. 7A-7B (SEQ ID NO:7); the nucleotide sequences of the cDNAs contained in the above-described deposited clones; or nucleic acid molecules having a sequence complementary to one of the above sequences. Such isolated molecules, particularly DNA molecules, are useful as probes for gene mapping, by *in situ* hybridization with chromosomes, and for detecting expression of the TR2 receptor genes of the present invention in human tissue, for instance, by Northern blot analysis.

The present invention is further directed to fragments of the isolated nucleic acid molecules described herein. By a fragment of an isolated nucleic acid molecule having the nucleotide sequence of the deposited cDNAs or the nucleotide sequence shown in FIG. 1A-1B (SEQ ID NO:1), FIG. 4A-4B (SEQ ID NO:4), or FIG. 7A-7B (SEQ ID NO:7) is intended fragments at least about 15 nt, and more preferably at least about 20 nt, still more preferably at least about 30 nt, and even more preferably, at least about 40 nt in length which are useful as diagnostic probes and primers as discussed herein. Of course, larger fragments 50-400 nt in length are also useful according to the present invention as are fragments corresponding to most, if not all, of the nucleotide sequences of the deposited cDNAs or as shown in FIG. 1A-1B (SEQ ID NO:1), FIG. 4A-4B (SEQ ID NO:4), or FIG. 7A-7B (SEQ ID NO:7). By a fragment at least 20 nt in length, for example, is intended fragments which include 20 or more contiguous bases from the nucleotide sequences of the deposited cDNAs or the nucleotide sequences as shown in FIG. 1A-1B (SEQ ID NO:1), FIG. 4A-4B (SEQ ID NO:4), or FIG. 7A-7B (SEQ ID NO:7).

Preferred nucleic acid fragments of the present invention include nucleic acid molecules encoding: a polypeptide comprising the TR2 receptor protein of FIG. 1A-1B (SEQ ID NO:2) extracellular domain (predicted to constitute amino acid residues from about 37 to about 200 in FIG. 1A-1B (amino acid residues 1

to 164 in SEQ ID NO:2)); a polypeptide comprising the TR2 receptor transmembrane domain (amino acid residues 201 to 225 in FIG. 1A-1B (amino acid residues 165 to 189 in SEQ ID NO:2)); a polypeptide comprising the TR2 receptor intracellular domain (predicted to constitute amino acid residues from about 226 to about 283 in FIG. 1A-1B (amino acid residues 190 to 247 in SEQ ID NO:2)); and a polypeptide comprising the TR2 receptor protein of FIG. 1A-1B (SEQ ID NO:2) extracellular and intracellular domains with all or part of the transmembrane domain deleted.

Preferred nucleic acid fragments of the present invention also include nucleic acid molecules encoding polypeptides comprising the mature TR2-SV1 receptor (predicted to constitute amino acid residues from about 37 to about 185 in FIG. 4A-4B (amino acid residues 1 to 149 in SEQ ID NO:5)) and the complete TR2-SV2 receptor (predicted to constitute amino acid residues from about 1 to about 136 in FIG. 7A-7B (SEQ ID NO:8)).

As above with the leader sequence, the amino acid residues constituting the extracellular, transmembrane and intracellular domains have been predicted by computer analysis. Thus, as one of ordinary skill would appreciate, the amino acid residues constituting these domains may vary slightly (e.g., by about 1 to about 15 amino acid residues) depending on the criteria used to define each domain

Preferred nucleic acid fragments of the present invention also include nucleic acid molecules encoding epitope-bearing portions of the TR2 receptor proteins. In particular, such nucleic acid fragments of the present invention include nucleic acid molecules encoding: a polypeptide comprising amino acid residues from about 39 to about 70 in FIG. 1A-1B (amino acid residues 3 to 34 in SEQ ID NO:2); a polypeptide comprising amino acid residues from about 106 to about 120 in FIG. 1 (amino acid residues 70 to 84 in SEQ ID NO:2); a polypeptide comprising amino acid residues from about 142 to about 189 in FIG. 1A-1B (amino acid residues 106 to 153 in SEQ ID NO:2); a polypeptide comprising amino acid residues from about 276 to about 283 in FIG. 1A-1B

(amino acid residues 240 to 247 in SEQ ID NO:2); a polypeptide comprising amino acid residues from about 39 to about 70 in FIG. 4A-4B (amino acid residues 3 to 34 in SEQ ID NO:5); amino acid residues from about 99 to about 136 in FIG. 4A-4B (amino acid residues 63 to 100 in SEQ ID NO:5); amino acid residues from about 171 to about 185 in FIG. 4A-4B (amino acid residues 135 to 149 in SEQ ID NO:5); amino acid residues from about 56 to about 68 in FIG. 7A-7B (SEQ ID NO:8); amino acid residues from about 93 to about 136 in FIG. 7A-7B (SEQ ID NO:8). The inventors have determined that the above polypeptide fragments are antigenic regions of the TR2 receptors. Methods for determining other such epitope-bearing portions of the TR2 proteins are described in detail below.

In another aspect, the invention provides isolated nucleic acid molecules comprising polynucleotides which hybridizes under stringent hybridization conditions to a portion of the polynucleotide of one of the nucleic acid molecules of the invention described above, for instance, the cDNA clones contained in ATCC Deposits 97059, 97058 and 97057. By "stringent hybridization conditions" is intended overnight incubation at 42°C in a solution comprising: 50% formamide, 5x SSC (150 mM NaCl, 15 mM trisodium citrate), 50 mM sodium phosphate (pH 7.6), 5x Denhardt's solution, 10% dextran sulfate, and 20 g/ml denatured, sheared salmon sperm DNA, followed by washing the filters in 0.1x SSC at about 65°C

By a polynucleotide which hybridizes to a "portion" of a polynucleotide is intended a polynucleotide (either DNA or RNA) hybridizing to at least about 15 nucleotides (nt), and more preferably at least about 20 nt, still more preferably at least about 30 nt, and even more preferably about 30-70 nt of the reference polynucleotide. These are useful as diagnostic probes and primers as discussed above and in more detail below.

By a portion of a polynucleotide of "at least 20 nt in length," for example, is intended 20 or more contiguous nucleotides from the nucleotide sequence of the

reference polynucleotide (e.g., the deposited cDNAs or the nucleotide sequences as shown in FIG. 1A-1B (SEQ ID NO:1), FIG. 4A-4B (SEQ ID NO:4), or FIG. 7A-7B (SEQ ID NO:7)).

Of course, a polynucleotide which hybridizes only to a poly A sequence (such as the 3' terminal poly(A) tract of the TR2 receptor cDNA sequences shown in FIG. 1A-1B (SEQ ID NO:1), FIG. 4A-4B (SEQ ID NO:4), or FIG. 7A-7B (SEQ ID NO:7)), or to a complementary stretch of T (or U) residues, would not be included in a polynucleotide of the invention used to hybridize to a portion of a nucleic acid of the invention, since such a polynucleotide would hybridize to any nucleic acid molecule containing a poly (A) stretch or the complement thereof (e.g., practically any double-stranded cDNA clone).

As indicated, nucleic acid molecules of the present invention which encode TR2 polypeptides may include, but are not limited to those encoding the amino acid sequences of the mature polypeptides, by itself; the coding sequence for the mature polypeptides and additional sequences, such as those encoding the about 36 amino acid leader or secretory sequences, such as pre-, or pro- or prepro-protein sequences; the coding sequence of the mature polypeptides, with or without the aforementioned additional coding sequences, together with additional, non-coding sequences, including for example, but not limited to introns and non-coding 5' and 3' sequences, such as the transcribed, non-translated sequences that play a role in transcription, mRNA processing, including splicing and polyadenylation signals, for example - ribosome binding and stability of mRNA; an additional coding sequence which codes for additional amino acids, such as those which provide additional functionalities. Thus, the sequence encoding the polypeptides may be fused to a marker sequence, such as a sequence encoding a peptide which facilitates purification of the fused polypeptide. In certain preferred embodiments of this aspect of the invention, the marker amino acid sequence is a hexa-histidine peptide, such as the tag provided in a pQE vector (Qiagen, Inc.), among others, many of which are commercially available. As described in Gentz

5 *et al., Proc. Natl. Acad. Sci. USA* 86:821-824 (1989), for instance, hexa-histidine provides for convenient purification of the fusion protein. The "HA" tag is another peptide useful for purification which corresponds to an epitope derived from the influenza hemagglutinin protein, which has been described by Wilson *et al., Cell* 37: 767 (1984). As discussed below, other such fusion proteins include the TR2 receptors fused to IgG-Fc at the N- or C-terminus.

10 The present invention further relates to variants of the nucleic acid molecules of the present invention, which encode portions, analogs or derivatives of the TR2 receptors. Variants may occur naturally, such as a natural allelic variant. By an "allelic variant" is intended one of several alternate forms of a gene occupying a given locus on a chromosome of an organism. *Genes II*, Lewin, B , ed., John Wiley & Sons, New York (1985). Non-naturally occurring variants may be produced using art-known mutagenesis techniques.

15 Such variants include those produced by nucleotide substitutions, deletions or additions, which may involve one or more nucleotides. The variants may be altered in coding regions, non-coding regions, or both. Alterations in the coding regions may produce conservative or non-conservative amino acid substitutions, deletions or additions. Especially preferred among these are silent substitutions, additions and deletions, which do not alter the properties and activities of the TR2  
20 receptors or portions thereof. Also especially preferred in this regard are conservative substitutions.

25 Further embodiments of the invention include isolated nucleic acid molecules comprising a polynucleotide having a nucleotide sequence at least 90% identical, and more preferably at least 95%, 96%, 97%, 98% or 99% identical to (a) a nucleotide sequence encoding the TR2 polypeptide having the complete amino acid sequence shown in FIG. 1A-1B (amino acid residues -36 to 247 in SEQ ID NO:2), FIG. 4A-4B (amino acid residues -36 to 149 in SEQ ID NO:5), or FIG. 7A-7B (amino acid residues 1 to 136 in SEQ ID NO:8); (b) a nucleotide encoding the complete amino sequence shown in FIG. 1A-1B (amino acid residues

-35 to 247 in SEQ ID NO:2), FIG. 4A-4B (amino acid residues -35 to 149 in SEQ ID NO:5), or FIG. 7A-7B (amino acid residues 2 to 136 in SEQ ID NO:8) but lacking the N-terminal methionine; (c) a nucleotide sequence encoding the mature TR2 receptors (full-length polypeptide with any attending leader sequence removed) having the amino acid sequence at positions from about 37 to about 283 in FIG. 1A-1B (amino acid residues 1 to 247 in SEQ ID NO:2) or the amino acid sequence at positions from about 37 to about 185 in FIG. 4A-4B (amino acid residues 1 to 149 in SEQ ID NO:5), or the amino acid sequence at positions from about 1 to about 136 in FIG. 7A-7B (SEQ ID NO:8); (d) a nucleotide sequence encoding the TR2, TR2-SV1 or TR2-SV2 polypeptides having the complete amino acid sequence including the leader encoded by the cDNA clones contained in ATCC Deposit Numbers 97059, 97058, and 97057, respectively; (e) a nucleotide sequence encoding the mature TR2 or TR2-SV1 receptors having the amino acid sequences encoded by the cDNA clones contained in ATCC Deposit Numbers 97059 and 97058, respectively; (f) a nucleotide sequence encoding the TR2 or TR2-SV1 receptor extracellular domain; (g) a nucleotide sequence encoding the TR2 receptor transmembrane domain; (h) a nucleotide sequence encoding the TR2 receptor intracellular domain; (i) a nucleotide sequence encoding the TR2 receptor extracellular and intracellular domains with all or part of the transmembrane domain deleted; and (j) a nucleotide sequence complementary to any of the nucleotide sequences in (a), (b), (c), (d), (e), (f), (g), (h), or (i).

By a polynucleotide having a nucleotide sequence at least, for example, 95% "identical" to a reference nucleotide sequence encoding a TR2 polypeptide is intended that the nucleotide sequence of the polynucleotide is identical to the reference sequence except that the polynucleotide sequence may include up to five point mutations per each 100 nucleotides of the reference nucleotide sequence encoding the TR2 receptors. In other words, to obtain a polynucleotide having a nucleotide sequence at least 95% identical to a reference nucleotide sequence,

up to 5% of the nucleotides in the reference sequence may be deleted or substituted with another nucleotide, or a number of nucleotides up to 5% of the total nucleotides in the reference sequence may be inserted into the reference sequence. These mutations of the reference sequence may occur at the 5' or 3' terminal positions of the reference nucleotide sequence or anywhere between those terminal positions, interspersed either individually among nucleotides in the reference sequence or in one or more contiguous groups within the reference sequence.

As a practical matter, whether any particular nucleic acid molecule is at least 90%, 95%, 96%, 97%, 98% or 99% identical to, for instance, the nucleotide sequence shown in FIG. 1A-1B (SEQ ID NO:1) or to the nucleotides sequence of the deposited cDNA clone encoding that protein can be determined conventionally using known computer programs such as the Bestfit program (Wisconsin Sequence Analysis Package, Version 8 for Unix, Genetics Computer Group, University Research Park, 575 Science Drive, Madison, WI 53711. Bestfit uses the local homology algorithm of Smith and Waterman, *Advances in Applied Mathematics* 2: 482-489 (1981), to find the best segment of homology between two sequences. When using Bestfit or any other sequence alignment program to determine whether a particular sequence is, for instance, 95% identical to a reference sequence according to the present invention, the parameters are set, of course, such that the percentage of identity is calculated over the full length of the reference nucleotide sequence and that gaps in homology of up to 5% of the total number of nucleotides in the reference sequence are allowed.

The present application is directed to nucleic acid molecules at least 90%, 95%, 96%, 97%, 98% or 99% identical to the nucleic acid sequences shown in FIG. 1A-1B (SEQ ID NO:1), FIG. 4A-4B (SEQ ID NO:4), or FIG. 7A-7B (SEQ ID NO:7) or to the nucleic acid sequence of the deposited cDNAs, irrespective of whether they encode a polypeptide having TR2 receptor activity. This is because even where a particular nucleic acid molecule does not encode a polypeptide



having TR2 receptor activity, one of skill in the art would still know how to use the nucleic acid molecule, for instance, as a hybridization probe or a polymerase chain reaction (PCR) primer. Uses of the nucleic acid molecules of the present invention that do not encode a polypeptide having TR2 receptor activity include, *inter alia*, (1) isolating a TR2 receptor gene or allelic or splice variants thereof in a cDNA library; (2) *in situ* hybridization (e.g., "FISH") to metaphase chromosomal spreads to provide precise chromosomal location of a TR2 receptor gene, as described in Verma *et al.*, *Human Chromosomes: A Manual of Basic Techniques*, Pergamon Press, New York (1988); and (3) Northern Blot analysis for detecting TR2 receptor mRNA expression in specific tissues.

Preferred, however, are nucleic acid molecules having sequences at least 90%, 95%, 96%, 97%, 98% or 99% identical to the nucleic acid sequence shown in FIG. 1A-1B (SEQ ID NO:1), FIG. 4A-4B (SEQ ID NO:4), or FIG. 7A-7B (SEQ ID NO:7) or to the nucleic acid sequence of the deposited cDNAs which do, in fact, encode a polypeptide having TR2 receptor activity. By "a polypeptide having TR2 receptor activity" is intended polypeptides exhibiting activity similar, but not necessarily identical, to an activity of the TR2 receptors of the present invention (either the full-length protein, the splice variants, or, preferably, the mature protein), as measured in a particular biological assay. For example, TR2 receptor activity can be measured by determining the ability of a polypeptide-Fc fusion protein to inhibit lymphocyte proliferation as described below in Example 6. TR2 receptor activity may also be measured by determining the ability of a polypeptide, such as cognate ligand which is free or expressed on a cell surface, to confer proliferatory activity in intact cells expressing the receptor.

Of course, due to the degeneracy of the genetic code, one of ordinary skill in the art will immediately recognize that a large number of the nucleic acid molecules having a sequence at least 90%, 95%, 96%, 97%, 98%, or 99% identical to the nucleic acid sequence of the deposited cDNAs or the nucleic acid sequences shown in FIG. 1A-1B (SEQ ID NO:1), FIG. 4A-4B (SEQ ID NO:4),

or FIG. 7A-7B (SEQ ID NO:7) will encode polypeptides "having TR2 receptor activity." In fact, since degenerate variants of any of these nucleotide sequences all encode the same polypeptide, this will be clear to the skilled artisan even without performing the above described comparison assay. It will be further recognized in the art that, for such nucleic acid molecules that are not degenerate variants, a reasonable number will also encode a polypeptide having TR2 protein activity. This is because the skilled artisan is fully aware of amino acid substitutions that are either less likely or not likely to significantly effect protein function (e.g., replacing one aliphatic amino acid with a second aliphatic amino acid).

For example, guidance concerning how to make phenotypically silent amino acid substitutions is provided in Bowie, J. U. *et al.*, "Deciphering the Message in Protein Sequences: Tolerance to Amino Acid Substitutions," *Science* 247:1306-1310 (1990), wherein the authors indicate that proteins are surprisingly tolerant of amino acid substitutions.

### *Vectors and Host Cells*

The present invention also relates to vectors which include the isolated DNA molecules of the present invention, host cells which are genetically engineered with the recombinant vectors, and the production of TR2 polypeptides or fragments thereof by recombinant techniques.

The polynucleotides may be joined to a vector containing a selectable marker for propagation in a host. Generally, a plasmid vector is introduced in a precipitate, such as a calcium phosphate precipitate, or in a complex with a charged lipid. If the vector is a virus, it may be packaged *in vitro* using an appropriate packaging cell line and then transduced into host cells.

The DNA insert should be operatively linked to an appropriate promoter, such as the phage lambda PL promoter, the *E. coli lac*, *trp* and *tac* promoters, the SV40 early and late promoters and promoters of retroviral LTRs, to name a few.

Other suitable promoters will be known to the skilled artisan. The expression constructs will further contain sites for transcription initiation, termination and, in the transcribed region, a ribosome binding site for translation. The coding portion of the mature transcripts expressed by the constructs will preferably include a translation initiating at the beginning and a termination codon (UAA, UGA or UAG) appropriately positioned at the end of the polypeptide to be translated.

As indicated, the expression vectors will preferably include at least one selectable marker. Such markers include dihydrofolate reductase or neomycin resistance for eukaryotic cell culture and tetracycline or ampicillin resistance genes for culturing in *E. coli* and other bacteria. Representative examples of appropriate heterologous hosts include, but are not limited to, bacterial cells, such as *E. coli*, *Streptomyces* and *Salmonella typhimurium* cells; fungal cells, such as yeast cells; insect cells such as *Drosophila* S2 and *Spodoptera* Sf9 cells; animal cells such as CHO, COS and Bowes melanoma cells; and plant cells. Appropriate culture mediums and conditions for the above-described host cells are known in the art.

Among vectors preferred for use in bacteria include pQE70, pQE60 and pQE-9, available from Qiagen; pBS vectors, Phagescript vectors, Bluescript vectors, pNH8A, pNH16a, pNH18A, pNH46A, available from Stratagene; and ptrc99a, pKK223-3, pKK233-3, pDR540, pRIT5 available from Pharmacia. Among preferred eukaryotic vectors are pWLNEO, pSV2CAT, pOG44, pXT1 and pSG available from Stratagene; and pSVK3, pBPV, pMSG and pSVL available from Pharmacia. Other suitable vectors will be readily apparent to the skilled artisan.

Introduction of the construct into the host cell can be effected by calcium phosphate transfection, DEAE-dextran mediated transfection, cationic lipid-mediated transfection, electroporation, transduction, infection or other methods. Such methods are described in many standard laboratory manuals, such as Davis *et al.*, *Basic Methods In Molecular Biology* (1986).

5 The polypeptide may be expressed in a modified form, such as a fusion protein, and may include not only secretion signals, but also additional heterologous functional regions. For instance, a region of additional amino acids, particularly charged amino acids, may be added to the N-terminus of the polypeptide to improve stability and persistence in the host cell, during purification, or during subsequent handling and storage. Also, peptide moieties may be added to the polypeptide to facilitate purification. Such regions may be removed prior to final preparation of the polypeptide. The addition of peptide moieties to polypeptides to engender secretion or excretion, to improve stability and to facilitate purification, among others, are familiar and routine techniques in the art. A preferred fusion protein comprises a heterologous region from immunoglobulin that is useful to solubilize proteins. For example, EP-A-O 464 533 (Canadian counterpart 2045869) discloses fusion proteins comprising various portions of constant region of immunoglobulin molecules together with another human protein or part thereof. In many cases, the Fc part in a fusion protein is thoroughly advantageous for use in therapy and diagnosis and thus results, for example, in improved pharmacokinetic properties (EP-A 0232 262). On the other hand, for some uses it would be desirable to be able to delete the Fc part after the fusion protein has been expressed, detected and purified in the advantageous manner described. This is the case when Fc portion proves to be a hindrance to use in therapy and diagnosis, for example when the fusion protein is to be used as antigen for immunizations. In drug discovery, for example, human proteins, such as, human hIL-5 receptor has been fused with Fc portions for the purpose of high-throughput screening assays to identify antagonists of hIL-5. See, D. Bennett *et al.*, *Journal of Molecular Recognition*, Vol. 8:52-58 (1995) and K. Johanson *et al.*, *The Journal of Biological Chemistry*, Vol. 270, No. 16:9459-9471 (1995).

25 TR2 receptors can be recovered and purified from recombinant cell cultures by well-known methods including ammonium sulfate or ethanol

precipitation, acid extraction, anion or cation exchange chromatography, phosphocellulose chromatography, hydrophobic interaction chromatography, affinity chromatography, hydroxylapatite chromatography and lectin chromatography. Most preferably, high performance liquid chromatography ("HPLC") is employed for purification. Polypeptides of the present invention include naturally purified products, products of chemical synthetic procedures, and products produced by recombinant techniques from a prokaryotic or eukaryotic host, including, for example, bacterial, yeast, higher plant, insect and mammalian cells. Depending upon the host employed in a recombinant production procedure, the polypeptides of the present invention may be glycosylated or may be non-glycosylated. In addition, polypeptides of the invention may also include an initial modified methionine residue, in some cases as a result of host-mediated processes.

#### *TR2 Polypeptides and Fragments*

The invention further provides isolated TR2 polypeptides having the amino acid sequence encoded by the deposited cDNAs, or the amino acid sequence in FIG. 1A-1B (SEQ ID NO:2), FIG. 4A-4B (SEQ ID NO:5), or FIG. 7A-7B (SEQ ID NO:8), or a peptide or polypeptide comprising a portion of the above polypeptides.

The polypeptides of this invention may be membrane bound or may be in a soluble circulating form. Soluble peptides are defined by amino acid sequence wherein the sequence comprises the polypeptide sequence lacking the transmembrane domain. One example of such a soluble form of the TR2 receptor is the TR2-SV1 splice variant which has a secretory leader sequence but lacks both the intracellular and transmembrane domains. Thus, the TR2-SV1 receptor protein appears to be secreted in a soluble form from cells which express this protein.

The polypeptides of the present invention may exist as a membrane bound receptor having a transmembrane region and an intra- and extracellular region or they may exist in soluble form wherein the transmembrane domain is lacking. One example of such a form of the TR2 receptor is the TR2 receptor shown in FIG. 1A-1B (SEQ ID NO:2) which contains, in addition to a leader sequence, transmembrane, intracellular and extracellular domains. Thus, this form of the TR2 receptor appears to be localized in the cytoplasmic membrane of cells which express this protein

It will be recognized in the art that some amino acid sequences of the TR2 receptors can be varied without significant effect to the structure or function of the protein. If such differences in sequence are contemplated, it should be remembered that there will be critical areas on the protein which determine activity. Thus, the invention further includes variations of the TR2 receptors which show substantial TR2 receptor activity or which include regions of TR2 proteins such as the protein portions discussed below. Such mutants include deletions, insertions, inversions, repeats, and type substitutions. As indicated above, guidance concerning which amino acid changes are likely to be phenotypically silent can be found in Bowie, J. U., *et al.*, "Deciphering the Message in Protein Sequences: Tolerance to Amino Acid Substitutions," *Science* 247:1306-1310 (1990).

Thus, the fragment, derivative or analog of the polypeptides of FIG. 1A-1B (SEQ ID NO:2), FIG. 4A-4B (SEQ ID NO:5), and FIG. 7A-7B (SEQ ID NO:8), or that encoded by the deposited cDNAs, may be (i) one in which one or more of the amino acid residues are substituted with a conserved or non-conserved amino acid residue (preferably a conserved amino acid residue) and such substituted amino acid residue may or may not be one encoded by the genetic code, or (ii) one in which one or more of the amino acid residues includes a substituent group, or (iii) one in which the mature polypeptide is fused with another compound, such as a compound to increase the half-life of the polypeptide

(for example, polyethylene glycol), or (iv) one in which the additional amino acids are fused to the mature polypeptide, such as an IgG Fc fusion region peptide or leader or secretory sequence or a sequence which is employed for purification of the mature polypeptide or a proprotein sequence. Such fragments, derivatives and analogs are deemed to be within the scope of those skilled in the art from the teachings herein.

Of particular interest are substitutions of charged amino acids with another charged amino acid and with neutral or negatively charged amino acids. The latter results in proteins with reduced positive charge to improve the characteristics of the TR2 proteins. The prevention of aggregation is highly desirable. Aggregation of proteins not only results in a loss of activity but can also be problematic when preparing pharmaceutical formulations, because they can be immunogenic. (Pinckard *et al.*, *Clin Exp. Immunol.* 2:331-340 (1967); Robbins *et al.*, *Diabetes* 36:838-845 (1987); Cleland *et al.* *Crit. Rev. Therapeutic Drug Carrier Systems* 10:307-377 (1993)).

The replacement of amino acids can also change the selectivity of binding to cell surface receptors. Ostade *et al.*, *Nature* 361:266-268 (1993) describes certain mutations resulting in selective binding of TNF- $\alpha$  to only one of the two known types of TNF receptors. Thus, the TR2 receptors of the present invention may include one or more amino acid substitutions, deletions or additions, either from natural mutations or human manipulation.

As indicated, changes are preferably of a minor nature, such as conservative amino acid substitutions that do not significantly affect the folding or activity of the protein (see Table 1).

**TABLE I**  
**CONSERVATIVE AMINO ACID SUBSTITUTIONS.**

Aromatic	Phenylalanine Tryptophan Tyrosine
Hydrophobic	Leucine Isoleucine Valine
Polar	Glutamine Asparagine
Basic	Arginine Lysine Histidine
Acidic	Aspartic Acid Glutamic Acid
Small	Alanine Serine Threonine Methionine Glycine

Amino acids in the TR2 proteins of the present invention that are essential for function can be identified by methods known in the art, such as site-directed mutagenesis or alanine-scanning mutagenesis (Cunningham and Wells, *Science* 244:1081-1085 (1989)). The latter procedure introduces single alanine mutations at every residue in the molecule. The resulting mutant molecules are then tested for biological activity such as receptor binding or *in vitro*, or *in vitro* proliferative activity. Sites that are critical for ligand-receptor binding can also be determined by structural analysis such as crystallization, nuclear magnetic resonance or photoaffinity labeling (Smith *et al.*, *J. Mol. Biol.* 224:899-904 (1992) and de Vos *et al. Science* 255:306-312 (1992)).

The polypeptides of the present invention are preferably provided in an isolated form. By "isolated polypeptide", is intended a polypeptide removed from its native environment. Thus, a polypeptide produced and contained within a



recombinant host cell would be considered "isolated" for purposes of the present invention. Also intended as an "isolated polypeptide" are polypeptides that have been purified, partially or substantially, from a recombinant host. For example, recombinantly produced versions of the TR2 receptors can be substantially purified by the one-step method described in Smith and Johnson, *Gene* 67:31-40 (1988).

The polypeptides of the present invention also include the polypeptide encoded by the deposited cDNAs including the leader; the polypeptide encoded by the deposited the cDNAs minus the leader (i.e., the mature protein); the polypeptides of FIG. 1A-1B (SEQ ID NO:2) or FIG. 4A-4B (SEQ ID NO:5) including the leader; the polypeptides of FIG. 1A-1B (SEQ ID NO:2) or FIG. 4A-4B (SEQ ID NO.5) including the leader but minus the N-terminal methionine; the polypeptides of FIG. 1A-1B (SEQ ID NO:2) or FIG. 4A-4B (SEQ ID NO:5) minus the leader; the polypeptide of FIG. 7A-7B (SEQ ID NO:8); the extracellular domain, the transmembrane domain, and the intracellular domain of the TR2 receptor shown in FIG. 1A-1B (SEQ ID NO:2); and polypeptides which are at least 80% identical, more preferably at least 90% or 95% identical, still more preferably at least 96%, 97%, 98% or 99% identical to the polypeptides described above, and also include portions of such polypeptides with at least 30 amino acids and more preferably at least 50 amino acids.

By a polypeptide having an amino acid sequence at least, for example, 95% "identical" to a reference amino acid sequence of a TR2 polypeptide is intended that the amino acid sequence of the polypeptide is identical to the reference sequence except that the polypeptide sequence may include up to five amino acid alterations per each 100 amino acids of the reference amino acid of a TR2 receptor. In other words, to obtain a polypeptide having an amino acid sequence at least 95% identical to a reference amino acid sequence, up to 5% of the amino acid residues in the reference sequence may be deleted or substituted with another amino acid, or a number of amino acids up to 5% of the total amino

acid residues in the reference sequence may be inserted into the reference sequence. These alterations of the reference sequence may occur at the amino or carboxy terminal positions of the reference amino acid sequence or anywhere between those terminal positions, interspersed either individually among residues in the reference sequence or in one or more contiguous groups within the reference sequence.

As a practical matter, whether any particular polypeptide is at least 90%, 95%, 96%, 97%, 98% or 99% identical to, for instance, the amino acid sequence shown in FIG. 1A-1B (SEQ ID NO:2), FIG. 4A-4B (SEQ ID NO:5), or FIG. 7A-7B (SEQ ID NO:8) or to the amino acid sequence encoded by one of the deposited cDNA clones can be determined conventionally using known computer programs such the Bestfit program (Wisconsin Sequence Analysis Package, Version 8 for Unix, Genetics Computer Group, University Research Park, 575 Science Drive, Madison, WI 53711). When using Bestfit or any other sequence alignment program to determine whether a particular sequence is, for instance, 95% identical to a reference sequence according to the present invention, the parameters are set, of course, such that the percentage of identity is calculated over the full length of the reference amino acid sequence and that gaps in homology of up to 5% of the total number of amino acid residues in the reference sequence are allowed.

The polypeptides of the present invention could be used as a molecular weight marker on SDS-PAGE gels or on molecular sieve gel filtration columns using methods well known to those of skill in the art.

In another aspect, the invention provides peptides or polypeptides comprising epitope-bearing portions of the polypeptides of the invention. The epitopes of these polypeptide portions are an immunogenic or antigenic epitopes of the polypeptides described herein. An "immunogenic epitope" is defined as a part of a protein that elicits an antibody response when the whole protein is the immunogen. On the other hand, a region of a protein molecule to which an

antibody can bind is defined as an "antigenic epitope." The number of immunogenic epitopes of a protein generally is less than the number of antigenic epitopes. See, for instance, Geysen *et al.*, *Proc. Natl. Acad. Sci. USA* 81:3998-4002 (1983).

5 As to the selection of peptides or polypeptides bearing an antigenic epitope (i.e., that contain a region of a protein molecule to which an antibody can bind), it is well known in that art that relatively short synthetic peptides that mimic part of a protein sequence are routinely capable of eliciting an antiserum that reacts with the partially mimicked protein. See, for instance, Sutcliffe, J. G., Shinnick, T. M., Green, N. and Learner, R.A. (1983) Antibodies that react with predetermined sites on proteins. *Science* 219:660-666. Peptides capable of eliciting protein-reactive sera are frequently represented in the primary sequence of a protein, can be characterized by a set of simple chemical rules, and are confined neither to immunodominant regions of intact proteins (i.e., immunogenic epitopes) nor to the amino or carboxyl terminals.

10 Antigenic epitope-bearing peptides and polypeptides of the invention are therefore useful to raise antibodies, including monoclonal antibodies, that bind specifically to a polypeptide of the invention. See, for instance, Wilson *et al.*, *Cell* 37:767-778 (1984) at 777. Antigenic epitope-bearing peptides and polypeptides of the invention preferably contain a sequence of at least seven, more preferably at least nine and most preferably between at least about 15 to about 30 amino acids contained within the amino acid sequence of a polypeptide of the invention.

20 Non-limiting examples of antigenic polypeptides or peptides that can be used to generate TR2 receptor-specific antibodies include: a polypeptide comprising amino acid residues from about 39 to about 70 in FIG. 1 (amino acid residues 3 to 34 in SEQ ID NO:2); a polypeptide comprising amino acid residues from about 106 to about 120 in FIG. 1A-1B (amino acid residues 70 to 84 in SEQ ID NO:2); a polypeptide comprising amino acid residues from about 142 to about 189 in FIG. 1A-1B (amino acid residues 106 to 153 in SEQ ID NO:2); a

polypeptide comprising amino acid residues from about 276 to about 283 in FIG. 1 (amino acid residues 240 to 247 in SEQ ID NO:2); a polypeptide comprising amino acid residues from about 39 to about 70 in FIG. 4A-4B (amino acid residues 3 to 34 in SEQ ID NO:5); a polypeptide comprising amino acid residues from about 99 to about 136 in FIG. 4A-4B (amino acid residues 63 to 100 in SEQ ID NO:5); a polypeptide comprising amino acid residues from about 171 to about 185 in FIG. 4A-4B (amino acid residues 135 to 149 in SEQ ID NO:5); a polypeptide comprising amino acid residues from about 56 to about 68 in FIG. 7A-7B (SEQ ID NO:8); and a polypeptide comprising amino acid residues from about 93 to about 136 in FIG. 7A-7B (SEQ ID NO:8). As indicated above, the inventors have determined that the above polypeptide fragments are antigenic regions of the TR2 receptor proteins.

The epitope-bearing peptides and polypeptides of the invention may be produced by any conventional means. Houghten, R. A. (1985) General method for the rapid solid-phase synthesis of large numbers of peptides: specificity of antigen-antibody interaction at the level of individual amino acids. *Proc. Natl. Acad. Sci. USA* 82:5131-5135. This "Simultaneous Multiple Peptide Synthesis (SMPS)" process is further described in U.S. Patent No. 4,631,211 to Houghten *et al.* (1986).

As one of skill in the art will appreciate, TR2 polypeptides of the present invention and the epitope-bearing fragments thereof described above can be combined with parts of the constant domain of immunoglobulins (IgG), resulting in chimeric polypeptides. These fusion proteins facilitate purification and show an increased half-life *in vivo*. This has been shown, e.g., for chimeric proteins consisting of the first two domains of the human CD4-polypeptide and various domains of the constant regions of the heavy or light chains of mammalian immunoglobulins (EPA 394,827; Trauneker *et al.*, *Nature* 331:84-86 (1988)). Fusion proteins that have a disulfide-linked dimeric structure due to the IgG part can also be more efficient in binding and neutralizing other molecules than the

monomeric TR2 receptor proteins or protein fragments alone (Fountoulakis *et al.*, *J. Biochem* 270:3958-3964 (1995)).

### *Detection of Disease States*

5 The TNF-family ligands induce various cellular responses by binding to  
TNF-family receptors, including the TR2 receptors of the present invention. TNF-  
10  $\beta$ , a potent ligand of the TNF receptor proteins, is known to be involved in a  
number of biological processes including lymphocyte development, tumor  
necrosis, induction of an antiviral state, activation of polymorphonuclear  
leukocytes, induction of class I major histocompatibility complex antigens on  
endothelial cells, induction of adhesion molecules on endothelium and growth  
hormone stimulation (Ruddle and Homer, *Prog. Allergy*, 40:162-182 (1988)).  
TNF- $\alpha$ , also a ligand of the TNF receptor proteins, has been reported to have a  
role in the rapid necrosis of tumors, immunostimulation, autoimmune disease,  
15 graft rejection, producing an anti-viral response, septic shock, cerebral malaria,  
cytotoxicity, protection against deleterious effects of ionizing radiation produced  
during a course of chemotherapy, such as denaturation of enzymes, lipid  
peroxidation and DNA damage (Nata *et al.*, *J. Immunol.* 136(7):2483 (1987);  
Porter, *Tibtech* 9 158-162 (1991)), growth regulation, vascular endothelium  
effects and metabolic effects. TNF- $\alpha$  also triggers endothelial cells to secrete  
20 various factors, including PAI-1, IL-1, GM-CSF and IL-6 to promote cell  
proliferation. In addition, TNF- $\alpha$  up-regulates various cell adhesion molecules  
such as E-Selectin, ICAM-1 and VCAM-1. TNF- $\alpha$  and the Fas ligand have also  
been shown to induce programmed cell death.

25 Cells which express the TR2 polypeptides and are believed to have a  
potent cellular response to TR2 receptor ligands include B lymphocytes (CD19<sup>+</sup>),  
both CD4<sup>+</sup> and CD8<sup>+</sup> T lymphocytes, monocytes, endothelial cells and other cell  
types shown in Tables 2 and 3. By "a cellular response to a TNF-family ligand"  
is intended any genotypic, phenotypic, and/or morphologic change to a cell, cell

line, tissue, tissue culture or patient that is induced by a TNF-family ligand. As indicated, such cellular responses include not only normal physiological responses to TNF-family ligands, but also diseases associated with increased cell proliferation or the inhibition of increased cell proliferation, such as by the inhibition of apoptosis. Apoptosis-programmed cell death-is a physiological mechanism involved in the deletion of peripheral T lymphocytes of the immune system, and its dysregulation can lead to a number of different pathogenic processes (Ameisen, J.C., *AIDS* 8:1197-1213 (1994); Krammer, P.H. *et al.*, *Curr. Opin. Immunol.* 6:279-289 (1994)).

It is believed that certain tissues in mammals with specific disease states associated with aberrant cell survival express significantly altered levels of the TR2 receptor protein and mRNA encoding the TR2 receptor protein when compared to a corresponding "standard" mammal, i.e., a mammal of the same species not having the disease state. Further, since some forms of this protein are secreted, it is believed that enhanced levels of the TR2 receptor protein can be detected in certain body fluids (e.g., sera, plasma, urine, and spinal fluid) from mammals with the disease state when compared to sera from mammals of the same species not having the disease state. Thus, the invention provides a diagnostic method useful during diagnosis of disease states, which involves assaying the expression level of the gene encoding the TR2 receptor protein in mammalian cells or body fluid and comparing the gene expression level with a standard TR2 receptor gene expression level, whereby an increase or decrease in the gene expression level over the standard is indicative of certain disease states associated with aberrant cell survival.

Where diagnosis of a disease state involving the TR2 receptors of the present invention has already been made according to conventional methods, the present invention is useful as a prognostic indicator, whereby patients exhibiting significantly aberrant TR2 receptor gene expression will experience a worse clinical outcome relative to patients expressing the gene at a lower level.

By "assaying the expression level of the gene encoding the TR2 receptor protein" is intended qualitatively or quantitatively measuring or estimating the level of the TR2 receptor protein or the level of the mRNA encoding the TR2 receptor protein in a first biological sample either directly (e.g., by determining or estimating absolute protein level or mRNA level) or relatively (e.g., by comparing to the TR2 receptor protein level or mRNA level in a second biological sample).

Preferably, the TR2 receptor protein level or mRNA level in the first biological sample is measured or estimated and compared to a standard TR2 receptor protein level or mRNA level, the standard being taken from a second biological sample obtained from an individual not having the disease state. As will be appreciated in the art, once a standard TR2 receptor protein level or mRNA level is known, it can be used repeatedly as a standard for comparison.

By "biological sample" is intended any biological sample obtained from an individual, cell line, tissue culture, or other source which contains TR2 receptor protein or mRNA. Biological samples include mammalian body fluids (such as sera, plasma, urine, synovial fluid and spinal fluid) which contain secreted mature TR2 receptor protein, and thymus, prostate, heart, placenta, muscle, liver, spleen, lung, kidney and other tissues. Methods for obtaining tissue biopsies and body fluids from mammals are well known in the art. Where the biological sample is to include mRNA, a tissue biopsy is the preferred source.

Diseases associated with increased cell survival, or the inhibition of apoptosis, include cancers (such as follicular lymphomas, carcinomas with p53 mutations, and hormone-dependent tumors); autoimmune disorders (such as systemic lupus erythematosus and immune-related glomerulonephritis rheumatoid arthritis) and viral infections (such as herpes viruses, pox viruses and adenoviruses), information graft v. host disease, acute graft rejection, and chronic graft rejection. Diseases associated with decreased cell survival, or increased apoptosis, include AIDS; neurodegenerative disorders (such as Alzheimer's disease, Parkinson's disease, Amyotrophic lateral sclerosis, Retinitis pigmentosa,

Cerebellar degeneration); myelodysplastic syndromes (such as aplastic anemia), ischemic injury (such as that caused by myocardial infarction, stroke and reperfusion injury), toxin-induced liver disease (such as that caused by alcohol), septic shock, cachexia and anorexia.

5           Assays available to detect levels of soluble receptors are well known to those of skill in the art, for example, radioimmunoassays, competitive-binding assays, Western blot analysis, and preferably an ELISA assay may be employed.

10           TR2 receptor-protein specific antibodies can be raised against the intact TR2 receptor protein or an antigenic polypeptide fragment thereof, which may presented together with a carrier protein, such as an albumin, to an animal system (such as rabbit or mouse) or, if it is long enough (at least about 25 amino acids), without a carrier.

15           As used herein, the term "antibody" (Ab) or "monoclonal antibody" (mAb) is meant to include intact molecules as well as antibody fragments (such as, for example, Fab and F(ab')<sub>2</sub> fragments) which are capable of specifically binding to TR2 receptor protein. Fab and F(ab')<sub>2</sub> fragments lack the Fc fragment of intact antibody, clear more rapidly from the circulation, and may have less non-specific tissue binding of an intact antibody (Wahl *et al.*, *J. Nucl. Med.* 24:316-325 (1983)). Thus, these fragments are preferred.

20           The antibodies of the present invention may be prepared by any of a variety of methods. For example, cells expressing the TR2 receptor protein or an antigenic fragment thereof can be administered to an animal in order to induce the production of sera containing polyclonal antibodies. In a preferred method, a preparation of TR2 receptor protein is prepared and purified to render it  
25           substantially free of natural contaminants. Such a preparation is then introduced into an animal in order to produce polyclonal antisera of greater specific activity.

          In the most preferred method, the antibodies of the present invention are monoclonal antibodies (or TR2 receptor protein binding fragments thereof). Such monoclonal antibodies can be prepared using hybridoma technology (Kohler *et al.*,



5 *Nature* 256:495 (1975); Kohler *et al.*, *Eur. J. Immunol.* 6:511 (1976); Kohler *et al.*, *Eur. J. Immunol.* 6:292 (1976); Hammerling *et al.*, In: *Monoclonal Antibodies and T-Cell Hybridomas*, Elsevier, N.Y., (1981) pp. 563-681). In general, such procedures involve immunizing an animal (preferably a mouse) with a TR2 receptor protein antigen or, more preferably, with a TR2 receptor protein-expressing cell. Suitable cells can be recognized by their capacity to bind anti-TR2 receptor protein antibody. Such cells may be cultured in any suitable tissue culture medium; however, it is preferable to culture cells in Earle's modified Eagle's medium supplemented with 10% fetal bovine serum (inactivated at about 56°C), and supplemented with about 10 g/l of nonessential amino acids, about 1,000 U/ml of penicillin, and about 100 µg/ml of streptomycin. The splenocytes of such mice are extracted and fused with a suitable myeloma cell line. Any suitable myeloma cell line may be employed in accordance with the present invention; however, it is preferable to employ the parent myeloma cell line (SP<sub>2</sub>O), available from the American Type Culture Collection, Rockville, Maryland. After fusion, the resulting hybridoma cells are selectively maintained in HAT medium, and then cloned by limiting dilution as described by Wands *et al.* (*Gastroenterology* 80:225-232 (1981)). The hybridoma cells obtained through such a selection are then assayed to identify clones which secrete antibodies capable of binding the TR2 receptor protein antigen.

#### *Agonists and Antagonists of TR2 Receptor Function*

25 In one aspect, the present invention is directed to a method for inhibiting an activity of TR2 induced by a TNF-family ligand (e.g., cell proliferation, hematopoietic development), which involves administering to a cell which expresses a TR2 polypeptide an effective amount of a TR2 receptor ligand, analog or an antagonist capable of decreasing TR2, receptor mediated signaling. Preferably, TR2 receptor mediated signaling is increased to treat a disease wherein increased cell proliferation is exhibited. An antagonist can include soluble forms

of the TR2 receptors and antibodies directed against the TR2 polypeptides which block TR2 receptor mediated signaling. Preferably, TR2 receptor mediated signaling is decreased to treat a disease.

5 In a further aspect, the present invention is directed to a method for increasing cell proliferation induced by a TNF-family ligand, which involves administering to a cell which expresses a TR2 polypeptide an effective amount of an agonist capable of increasing TR2 receptor mediated signaling. Preferably, TR2 receptor mediated signaling is increased to treat a disease wherein decreased cell proliferation is exhibited. Agonists of the present invention include  
10 monoclonal antibodies directed against the TR2 polypeptides which stimulate TR2 receptor mediated signaling. Preferably, TR2 receptor mediated signaling is increased to treat a disease.

15 By "agonist" is intended naturally occurring and synthetic compounds capable of enhancing cell proliferation and differentiation mediated by TR2 polypeptides. Such agonists include agents which increase expression of TR2 receptors or increase the sensitivity of the expressed receptor. By "antagonist" is intended naturally occurring and synthetic compounds capable of inhibiting TR2 mediated cell proliferation and differentiation. Such antagonists include agents which decrease expression of TR2 receptors or decrease the sensitivity of the  
20 expressed receptor. Whether any candidate "agonist" or "antagonist" of the present invention can enhance or inhibit cell proliferation and differentiation can be determined using art-known TNF-family ligand/receptor cellular response assays, including those described in more detail below.

25 One such screening technique involves the use of cells which express the receptor (for example, transfected CHO cells) in a system which measures extracellular pH changes caused by receptor activation, for example, as described in *Science* 246:181-296 (October 1989). For example, compounds may be contacted with a cell which expresses the receptor polypeptide of the present invention and a second messenger response, e.g., signal transduction or pH

changes, may be measured to determine whether the potential compound activates or inhibits the receptor.

Another such screening technique involves introducing RNA encoding the receptor into *Xenopus* oocytes to transiently express the receptor. The receptor oocytes may then be contacted with the receptor ligand and a compound to be screened, followed by detection of inhibition or activation of a calcium signal in the case of screening for compounds which are thought to inhibit activation of the receptor.

Another method involves screening for compounds which inhibit activation of the receptor polypeptide of the present invention antagonists by determining inhibition of binding of labeled ligand to cells which have the receptor on the surface thereof. Such a method involves transfecting a eukaryotic cell with DNA encoding the receptor such that the cell expresses the receptor on its surface and contacting the cell with a compound in the presence of a labeled form of a known ligand. The ligand can be labeled, e.g., by radioactivity. The amount of labeled ligand bound to the receptors is measured, e.g., by measuring radioactivity of the receptors. If the compound binds to the receptor as determined by a reduction of labeled ligand which binds to the receptors, the binding of labeled ligand to the receptor is inhibited.

Soluble forms of the polypeptides of the present invention may be utilized in the ligand binding assay described above. These forms of the TR2 receptors are contacted with ligands in the extracellular medium after they are secreted. A determination is then made as to whether the secreted protein will bind to TR2 receptor ligands.

Further screening assays for agonist and antagonist of the present invention are described in Tartaglia, L.A., and Goeddel, D.V., *J. Biol. Chem.* 267(7):4304-4307(1992).

Thus, in a further aspect, a screening method is provided for determining whether a candidate agonist or antagonist is capable of enhancing or inhibiting a

cellular response to a TNF-family ligand. The method involves contacting cells which express TR2 polypeptides with a candidate compound and a TNF-family ligand, assaying a cellular response, and comparing the cellular response to a standard cellular response, the standard being assayed when contact is made with the ligand in absence of the candidate compound, whereby an increased cellular response over the standard indicates that the candidate compound is an agonist of the ligand/receptor signaling pathway and a decreased cellular response compared to the standard indicates that the candidate compound is an antagonist of the ligand/receptor signaling pathway. By "assaying a cellular response" is intended qualitatively or quantitatively measuring a cellular response to a candidate compound and/or a TNF-family ligand (e.g., determining or estimating an increase or decrease in T cell proliferation or tritiated thymidine labeling). By the invention, a cell expressing a TR2 polypeptide can be contacted with either an endogenous or exogenously administered TNF-family ligand.

In an additional aspect, a thymocyte proliferation assay may be employed to identify both ligands and potential drug candidates. For example, thymus cells are disaggregated from tissue and grown in culture medium. Incorporation of DNA precursors such as  $^3\text{H}$ -thymidine or 5-bromo-2'-deoxyuridine (BrdU) is monitored as a parameter for DNA synthesis and cellular proliferation. Cells which have incorporated BrdU into DNA can be detected using a monoclonal antibody against BrdU and measured by an enzyme or fluorochrome-conjugated second antibody. The reaction is quantitated by fluorimetry or by spectrophotometry. Two control wells and an experimental well are set up as above and TNF- $\beta$  or cognate ligand is added to all wells while soluble receptor polypeptides of the present invention are added individually to the second control wells, with the experimental well containing a compound to be screened. The ability of the compound to be screened to stimulate or inhibit the above interaction may then be quantified.

Agonists according to the present invention include compounds such as, for example, TNF-family ligand peptide fragments, transforming growth factor  $\beta$ , and neurotransmitters (such as glutamate, dopamine, *N*-methyl-D-aspartate). Preferred agonist include polyclonal and monoclonal antibodies raised against TR2 polypeptide, or a fragment thereof. Such agonist antibodies raised against a TNF-family receptor are disclosed in Tartaglia, L.A., *et al.*, *Proc. Natl. Acad. Sci. USA* 88:9292-9296 (1991); and Tartaglia, L.A., and Goeddel, D.V., *J. Biol. Chem.* 267 (7):4304-4307 (1992). See, also, PCT Application WO 94/09137. Further preferred agonists include chemotherapeutic drugs such as, for example, cisplatin, doxorubicin, bleomycin, cytosine arabinoside, nitrogen mustard, methotrexate and vincristine. Others include ethanol and  $\beta$ -amyloid peptide. (*Science* 267:1457-1458 (1995)).

Antagonist according to the present invention include soluble forms of the TR2 receptors (e.g., fragments of the TR2 receptor shown in FIG. 1A-1B that include the ligand binding domain from the extracellular region of the full length receptor). Such soluble forms of the receptor, which may be naturally occurring or synthetic, antagonize TR2, TR2-SV1 or TR2-SV2 mediated signaling by competing with the cell surface bound forms of the receptor for binding to TNF-family ligands.. Antagonists of the present invention also include antibodies specific for TNF-family ligands and TR2-Fc fusion proteins such as the one described below in Examples 5 and 6.

By a "TNF-family ligand" is intended naturally occurring, recombinant, and synthetic ligands that are capable of binding to a member of the TNF receptor family and inducing the ligand/receptor signaling pathway. Members of the TNF ligand family include, but are not limited to, TNF- $\alpha$ , lymphotoxin- $\alpha$  (LT- $\alpha$ , also known as TNF- $\beta$ ), LT- $\beta$  (found in complex heterotrimer LT- $\alpha$ 2- $\beta$ ), FasL, CD40L, CD27L, CD30L, 4-1BBL, OX40L and nerve growth factor (NGF).

The experiments set forth in Example 6 demonstrate that the TR2 receptors of the present invention are capable of inducing the proliferation of

lymphocytes. Further, such proliferation can be inhibited by a TR2 protein fragment fused to an Fc antibody fragment.

TNF $\alpha$  has been shown to protect mice from infection with herpes simplex virus type 1 (HSV-1). Rossol-Voth, R. *et al.*, *J. Gen. Virol.* 72:143-147 (1991).  
5 The mechanism of the protective effect of TNF $\alpha$  is unknown but appears to involve neither interferons nor NK cell killing. One member of the TNFR family has been shown to mediate HSV-1 entry into cells. Montgomery, R. *et al.*, *Eur. Cytokine Newt.* 7:159 (1996). Further, antibodies specific for the extracellular domain of this TNFR block HSV-1 entry into cells. Thus, TR2 receptors of the present invention include both TR2 amino acid sequences and antibodies capable of preventing TNFR mediated viral entry into cells. Such sequences and antibodies can function by either competing with cell surface localized TNFR for binding to virus or by directly blocking binding of virus to cell surface receptors.

Antibodies according to the present invention may be prepared by any of a variety of methods using TR2 receptor immunogens of the present invention. Such TR2 receptor immunogens include the TR2 receptor protein shown in FIG. 1A-1B (SEQ ID NO:2) and the TR2-SV1 (FIG. 4A-4B (SEQ ID NO:5)) and TR2-SV2 (FIG. 7A-7B (SEQ ID NO:8)) polypeptides (any of which may or may not include a leader sequence) and polypeptide fragments of the receptors comprising the ligand binding, extracellular, transmembrane, the intracellular domains of the TR2 receptors, or any combination thereof.

Polyclonal and monoclonal antibody agonist or antagonist according to the present invention can be raised according to the methods disclosed in Tartaglia and Goeddel, *J. Biol. Chem.* 267(7):4304-4307(1992)); Tartaglia *et al.*, *Cell* 73:213-216 (1993)), and PCT Application WO 94/09137. The term "antibody" (Ab) or "monoclonal antibody" (mAb) as used herein is meant to include intact molecules as well as fragments thereof (such as, for example, Fab and F(ab')<sub>2</sub> fragments) which are capable of binding an antigen. Fab and F(ab')<sub>2</sub> fragments lack the Fc fragment of intact antibody, clear more rapidly from the circulation,

and may have less non-specific tissue binding of an intact antibody (Wahl *et al.*, *J. Nucl. Med.* 24:316-325 (1983)).

In a preferred method, antibodies according to the present invention are mAbs. Such mAbs can be prepared using hybridoma technology (Kohler and Millstein, *Nature* 256:495-497 (1975) and U.S. Patent No. 4,376,110; Harlow *et al.*, *Antibodies: A Laboratory Manual*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, 1988; *Monoclonal Antibodies and Hybridomas: A New Dimension in Biological Analyses*, Plenum Press, New York, NY, 1980; Campbell, "Monoclonal Antibody Technology," In: *Laboratory Techniques in Biochemistry and Molecular Biology*, Volume 13 (Burdon *et al.*, eds ), Elsevier, Amsterdam (1984)).

Proteins and other compounds which bind the TR2 receptor domains are also candidate agonist and antagonist according to the present invention. Such binding compounds can be "captured" using the yeast two-hybrid system (Fields and Song, *Nature* 340:245-246 (1989)). A modified version of the yeast two-hybrid system has been described by Roger Brent and his colleagues (Gyuris, J. *et al.*, *Cell* 75:791-803 (1993); Zervos, A.S. *et al.*, *Cell* 72:223-232 (1993)). Preferably, the yeast two-hybrid system is used according to the present invention to capture compounds which bind to the ligand binding, extracellular, intracellular, and transmembrane domains of the TR2 receptors. Such compounds are good candidate agonist and antagonist of the present invention.

Using the two-hybrid assay described above, the intracellular domain of the TR2 receptor, or a portion thereof, may be used to identify cellular proteins which interact with the receptor *in vivo*. Such an assay may also be used to identify ligands with potential agonistic or antagonistic activity of TR2 receptor function. This screening assay has previously been used to identify protein which interact with the cytoplasmic domain of the murine TNF-RII and led to the identification of two receptor associated proteins. Rothe, M. *et al.*, *Cell* 78:681 (1994). Such proteins and amino acid sequences which bind to the cytoplasmic

domain of the TR2 receptors are good candidate agonist and antagonist of the present invention.

Other screening techniques include the use of cells which express the polypeptide of the present invention (for example, transfected CHO cells) in a system which measures extracellular pH changes caused by receptor activation, for example, as described in *Science*, 246:181-296 (1989). In another example, potential agonists or antagonists may be contacted with a cell which expresses the polypeptide of the present invention and a second messenger response, e.g., signal transduction may be measured to determine whether the potential antagonist or agonist is effective.

The TR2 receptor agonists may be employed to stimulate ligand activities, such as inhibition of tumor growth and necrosis of certain transplantable tumors. The agonists may also be employed to stimulate cellular differentiation, for example, T-cell, fibroblasts and hemopoietic cell differentiation. Agonists to the TR2 receptor may also augment TR2's role in the host's defense against microorganisms and prevent related diseases (infections such as that from *Listeria monocytogenes*) and Chlamidiae. The agonists may also be employed to protect against the deleterious effects of ionizing radiation produced during a course of radiotherapy, such as denaturation of enzymes, lipid peroxidation, and DNA damage.

Agonists to the receptor polypeptides of the present invention may be used to augment TNF's role in host defenses against microorganisms and prevent related diseases. The agonists may also be employed to protect against the deleterious effects of ionizing radiation produced during a course of radiotherapy, such as denaturation of enzymes, lipid peroxidation, and DNA damage.

The agonists may also be employed to mediate an anti-viral response, to regulate growth, to mediate the immune response and to treat immunodeficiencies related to diseases such as HIV by increasing the rate of lymphocyte proliferation and differentiation.



The antagonists to the polypeptides of the present invention may be employed to inhibit ligand activities, such as stimulation of tumor growth and necrosis of certain transplantable tumors. The antagonists may also be employed to inhibit cellular differentiation, for example, T-cell, fibroblasts and hemopoietic cell differentiation. Antagonists may also be employed to treat autoimmune diseases, for example, graft versus host rejection and allograft rejection, and T-cell mediated autoimmune diseases such as AIDS. It has been shown that T-cell proliferation is stimulated via a type 2 TNF receptor. Accordingly, antagonizing the receptor may prevent the proliferation of T-cells and treat T-cell mediated autoimmune diseases.

The state of immunodeficiency that defines AIDS is secondary to a decrease in the number and function of CD4<sup>+</sup> T-lymphocytes. Recent reports estimate the daily loss of CD4<sup>+</sup> T cells to be between  $3.5 \times 10^7$  and  $2 \times 10^9$  cells (Wei X., *et al.*, *Nature* 373:117-122 (1995)). One cause of CD4<sup>+</sup> T cell depletion in the setting of HIV infection is believed to be HIV-induced apoptosis. Indeed, HIV-induced apoptotic cell death has been demonstrated not only *in vitro* but also, more importantly, in infected individuals (Ameisen, J.C., *AIDS* 8:1197-1213 (1994); Finkel, T.H., and Banda, N.K., *Curr. Opin. Immunol.* 6:605-615(1995); Muro-Cacho, C.A. *et al.*, *J. Immunol.* 154:5555-5566 (1995)). Furthermore, apoptosis and CD4<sup>+</sup> T-lymphocyte depletion is tightly correlated in different animal models of AIDS (Brunner, T., *et al.*, *Nature* 373:441-444 (1995); Gougeon, M.L., *et al.*, *AIDS Res. Hum. Retroviruses* 9:553-563 (1993)) and, apoptosis is not observed in those animal models in which viral replication does not result in AIDS (Gougeon, M.L. *et al.*, *AIDS Res. Hum. Retroviruses* 9:553-563 (1993)). Further data indicates that uninfected but primed or activated T lymphocytes from HIV-infected individuals undergo apoptosis after encountering the TNF-family ligand FasL. Using monocytic cell lines that result in death following HIV infection, it has been demonstrated that infection of U937 cells with HIV results in the *de novo* expression of FasL and that FasL mediates HIV-

induced apoptosis (Badley, A.D. *et al.*, *J. Virol.* 70:199-206 (1996)). Further the TNF-family ligand was detectable in uninfected macrophages and its expression was upregulated following HIV infection resulting in selective killing of uninfected CD4<sup>+</sup> T-lymphocytes (Badley, A.D *et al.*, *J. Virol.* 70:199-206 (1996)).

5 As shown in Example 6, the TR2 receptor shown in FIG. 1A-1B is expressed in CD4<sup>+</sup> T-lymphocytes and is capable of inducing lymphocyte proliferation. Thus, by the invention, a method for treating HIV<sup>+</sup> individuals is provided which involves administering an agonist of the present invention to increase the rate of proliferation and differentiation of CD4<sup>+</sup> T-lymphocytes. Such agonists include agents capable of inducing the expression of TR2 receptors (e.g., TNF $\alpha$ , PMA and DMSO) or enhancing the signal of such receptors which induces lymphocyte proliferation and differentiation. Modes of administration and dosages are discussed in detail below.

10 In rejection of an allograft, the immune system of the recipient animal has not previously been primed to respond because the immune system for the most part is only primed by environmental antigens. Tissues from other members of the same species have not been presented in the same way that, for example, viruses and bacteria have been presented. In the case of allograft rejection, immunosuppressive regimens are designed to prevent the immune system from reaching the effector stage. However, the immune profile of xenograft rejection may resemble disease recurrence more than allograft rejection. In the case of disease recurrence, the immune system has already been activated, as evidenced by destruction of the native islet cells. Therefore, in disease recurrence the immune system is already at the effector stage. Antagonists of the present invention are able to suppress the immune response to both allografts and xenografts by decreasing the rate of TR2 mediated lymphocyte proliferation and differentiation. Such antagonists include the TR2-Fc fusion protein described in Examples 5 and 6. Thus, the present invention further provides a method for suppression of immune responses.

In addition, TNF $\alpha$  has been shown to prevent diabetes in strains of animals which are prone to this affliction resulting from autoimmunity. *See* Porter, A., *Tibtech* 9:158-162 (1991). Thus, agonists and antagonists of the present invention may be useful in the treatment of autoimmune diseases such as type 1 diabetes.

In addition, the role played by the TR2 receptors in cell proliferation and differentiation indicates that agonist or antagonist of the present invention may be used to treat disease states involving aberrant cellular expression of these receptors. TR2 receptors may in some circumstances induce an inflammatory response, and antagonists may be useful reagents for blocking this response. Thus TR2 receptor antagonists (e.g., soluble forms of the TR2 receptors; neutralizing antibodies) may be useful for treating inflammatory diseases, such as rheumatoid arthritis, osteoarthritis, psoriasis, septicemia, and inflammatory bowel disease.

Antagonists to the TR2 receptor may also be employed to treat and/or prevent septic shock, which remains a critical clinical condition. Septic shock results from an exaggerated host response, mediated by protein factors such as TNF and IL-1, rather than from a pathogen directly. For example, lipopolysaccharides have been shown to elicit the release of TNF leading to a strong and transient increase of its serum concentration. TNF causes shock and tissue injury when administered in excessive amounts. Accordingly, it is believed that antagonists to the TR2 receptor will block the actions of TNF and treat/prevent septic shock. These antagonists may also be employed to treat meningococemia in children which correlates with high serum levels of TNF.

Among other disorders which may be treated by the antagonists to TR2 receptors, there are included, inflammation which is mediated by TNF receptor ligands, and the bacterial infections cachexia and cerebral malaria. The TR2 receptor antagonists may also be employed to treat inflammation mediated by ligands to the receptor such as TNF.

### *Modes of administration*

5 The agonist or antagonists described herein can be administered *in vitro*,  
*ex vivo*, or *in vivo* to cells which express the receptor of the present invention. By  
administration of an "effective amount" of an agonist or antagonist is intended an  
amount of the compound that is sufficient to enhance or inhibit a cellular response  
to a TNF-family ligand and include polypeptides. In particular, by administration  
of an "effective amount" of an agonist or antagonists is intended an amount  
effective to enhance or inhibit TR2 receptor mediated activity. Of course, where  
cell proliferation and/or differentiation is to be enhanced, an agonist according to  
the present invention can be co-administered with a TNF-family ligand. One of  
ordinary skill will appreciate that effective amounts of an agonist or antagonist can  
be determined empirically and may be employed in pure form or in  
pharmaceutically acceptable salt, ester or pro-drug form. The agonist or  
antagonist may be administered in compositions in combination with one or more  
pharmaceutically acceptable excipients.

10 It will be understood that, when administered to a human patient, the total  
daily usage of the compounds and compositions of the present invention will be  
decided by the attending physician within the scope of sound medical judgement  
The specific therapeutically effective dose level for any particular patient will  
depend upon factors well known in the medical arts

20 As a general proposition, the total pharmaceutically effective amount of  
a TR2 polypeptide administered parenterally per dose will be in the range of about  
1 µg/kg/day to 10 mg/kg/day of patient body weight, although, as noted above,  
this will be subject to therapeutic discretion. More preferably, this dose is at least  
0.01 mg/kg/day, and most preferably for humans between about 0.01 and 1  
mg/kg/day for the hormone. If given continuously, the TR2 polypeptide is  
typically administered at a dose rate of about 1 µg/kg/hour to about 50  
µg/kg/hour, either by 1-4 injections per day or by continuous subcutaneous

infusions, for example, using a mini-pump. An intravenous bag solution may also be employed.

Pharmaceutical compositions containing the TR2 receptor polypeptides of the invention may be administered orally, rectally, parenterally, intracisternally, intravaginally, intraperitoneally, topically (as by powders, ointments, drops or transdermal patch), buccally, or as an oral or nasal spray. By "pharmaceutically acceptable carrier" is meant a non-toxic solid, semisolid or liquid filler, diluent, encapsulating material or formulation auxiliary of any type. The term "parenteral" as used herein refers to modes of administration which include intravenous, intramuscular, intraperitoneal, intrasternal, subcutaneous and intraarticular injection and infusion

Having generally described the invention, the same will be more readily understood by reference to the following examples, which are provided by way of illustration and are not intended as limiting.

### *Example 1*

#### *Expression and Purification of TR2 in E. coli*

The bacterial expression vector pQE60 is used for bacterial expression in this example. (QIAGEN, Inc., 9259 Eton Avenue, Chatsworth, CA, 91311). pQE60 encodes ampicillin antibiotic resistance ("Amp<sup>r</sup>") and contains a bacterial origin of replication ("ori"), an IPTG inducible promoter, a ribosome binding site ("RBS"), six codons encoding histidine residues that allow affinity purification using nickel-nitrilo-tri-acetic acid ("Ni-NTA") affinity resin sold by QIAGEN, Inc., *supra*, and suitable single restriction enzyme cleavage sites. These elements are arranged such that a DNA fragment encoding a polypeptide may be inserted in such a way as to produce that polypeptide with the six His residues (i.e., a "6 X His tag") covalently linked to the carboxyl terminus of that polypeptide. However, in this example, the polypeptide coding sequence is inserted such that

translation of the six His codons is prevented and, therefore, the polypeptide is produced with no 6 X His tag.

The DNA sequence encoding the desired portion of the TR2 protein lacking the hydrophobic leader sequence is amplified from the deposited cDNA clone using PCR oligonucleotide primers which anneal to the amino terminal sequences of the desired portion of the TR2 protein and to sequences in the deposited construct 3' to the cDNA coding sequence. Additional nucleotides containing restriction sites to facilitate cloning in the pQE60 vector are added to the 5' and 3' sequences, respectively.

For cloning the mature protein, the 5' primer has the sequence:

5' CGCCCATGGCCCCAGCTCTGCCGTCCT 3' (SEQ ID NO:14) containing the underlined NcoI restriction site followed by 18 nucleotides complementary to the amino terminal coding sequence of the mature TR2 sequence in FIG. 1A-1B. One of ordinary skill in the art would appreciate, of course, that the point in the protein coding sequence where the 5' primer begins may be varied to amplify a desired portion of the complete protein shorter or longer than the mature form. The 3' primer has the sequence:

5' CGCAAAGCTTATTGTGGGAGCTGCTGGTCCC 3' (SEQ ID NO:15) containing the underlined HindIII restriction site followed by 18 nucleotides complementary to the 3' end of the nucleotide sequence shown in FIG. 1A-1B (SEQ ID NO:1) encoding the extracellular domain of the TR2 receptor.

The amplified TR2 DNA fragments and the vector pQE60 are digested with NcoI and HindIII and the digested DNAs are then ligated together. Insertion of the TR2 DNA into the restricted pQE60 vector places the TR2 protein coding region including its associated stop codon downstream from the IPTG-inducible promoter and in-frame with an initiating AUG. The associated stop codon prevents translation of the six histidine codons downstream of the insertion point.

The ligation mixture is transformed into competent *E. coli* cells using standard procedures such as those described in Sambrook *et al.*, *Molecular*

*Cloning: a Laboratory Manual, 2nd Ed.*; Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY (1989). *E. coli* strain MI5/rep4, containing multiple copies of the plasmid pREP4, which expresses the lac repressor and confers kanamycin resistance ("Kan"), is used in carrying out the illustrative example described herein. This strain, which is only one of many that are suitable for expressing TR2 protein, is available commercially from QIAGEN, Inc., *supra*. Transformants are identified by their ability to grow on LB plates in the presence of ampicillin and kanamycin. Plasmid DNA is isolated from resistant colonies and the identity of the cloned DNA confirmed by restriction analysis, PCR and DNA sequencing.

Clones containing the desired constructs are grown overnight ("O/N") in liquid culture in LB media supplemented with both ampicillin (100 µg/ml) and kanamycin (25 µg/ml). The O/N culture is used to inoculate a large culture, at a dilution of approximately 1:25 to 1:250. The cells are grown to an optical density at 600 nm ("OD600") of between 0.4 and 0.6. Isopropyl-b-D-thiogalactopyranoside ("IPTG") is then added to a final concentration of 1 mM to induce transcription from the lac repressor sensitive promoter, by inactivating the lacI repressor. Cells subsequently are incubated further for 3 to 4 hours. Cells then are harvested by centrifugation.

The cells are then stirred for 3-4 hours at 4°C in 6 M guanidine-HCl, pH 8. The cell debris is removed by centrifugation, and the supernatant containing the TR2 is dialyzed against 50 mM Na-acetate buffer pH 6, supplemented with 200 mM NaCl. Alternatively, the protein can be successfully refolded by dialyzing it against 500 mM NaCl, 20% glycerol, 25 mM Tris/HCl pH 7.4, containing protease inhibitors. After renaturation the protein can be purified by ion exchange, hydrophobic interaction and size exclusion chromatography. Alternatively, an affinity chromatography step such as an antibody column can be used to obtain pure TR2 protein. The purified protein is stored at 4°C or frozen at -80°C.

## Example 2

### *Example 2(a): Cloning and Expression of a Soluble Fragment of TR2 Protein in a Baculovirus Expression System*

5 In this example, the plasmid shuttle vector pA2 GP was used to insert the cloned DNA encoding the mature extracellular domain of the TR2 receptor protein shown in FIG. 1A-1B, lacking its naturally associated secretory signal (leader) sequence, into a baculovirus. This protein was expressed using a baculovirus leader and standard methods as described in Summers *et al.*, *A Manual of Methods for Baculovirus Vectors and Insect Cell Culture Procedures*, Texas Agricultural Experimental Station Bulletin No. 1555 (1987). This expression vector contains the strong polyhedrin promoter of the *Autographa californica* nuclear polyhedrosis virus (AcMNPV) followed by the secretory signal peptide (leader) of the baculovirus gp67 protein and convenient restriction sites such as BamHI, XbaI and Asp718. The polyadenylation site of the simian virus 40 ("SV40") is used for efficient polyadenylation. For easy selection of recombinant virus, the plasmid contains the beta-galactosidase gene from *E. coli* under control of a weak *Drosophila* promoter in the same orientation, followed by the polyadenylation signal of the polyhedrin gene. The inserted genes are flanked on both sides by viral sequences for cell-mediated homologous recombination with wild-type viral DNA to generate viable virus that expresses the cloned polynucleotide.

20 Many other baculovirus vectors could be used in place of the vector above, such as pAc373, pVL941 and pAcIM1, as one skilled in the art would readily appreciate, as long as the construct provides appropriately located signals for transcription, translation, secretion and the like, including a signal peptide and an in-frame AUG as required. Such vectors are described, for instance, in Luckow *et al.*, *Virology* 170:31-39.



The cDNA sequence encoding essentially the mature extracellular domain (amino acids 37 to 200 shown in FIG. 1A-1B) of the TR2 receptor protein in the deposited clone (ATCC Deposit Number 97059) was amplified using PCR oligonucleotide primers corresponding to the relevant 5' and 3' sequences of the gene. The 5' primer for each of the above has the sequence:

5' CGCGGATCCCGGAGCCCCCTGCTAC 3' (SEQ ID NO:16) containing the underlined BamHI restriction enzyme site, an efficient signal for initiation of translation in eukaryotic cells, as described by Kozak, M., *J. Mol. Biol.* 196:947-950 (1987), followed by 15 bases of the coding sequence of the TR2 protein shown in FIG. 1A-1B, beginning with the nucleotide 354. The 3' primer has the sequence:

5' CGCGGTACCATTGTGGGAGCTGCTGGTCCC 3' (SEQ ID NO:17) containing the underlined, Asp718 restriction sites followed by 17 nucleotides complementary to the coding sequences in FIG. 1A-1B.

The amplified fragment was isolated from a 1% agarose gel using a commercially available kit ("GeneClean," BIO 101 Inc., La Jolla, Ca.). The fragment was then digested with BamHI and Asp718 and purified on a 1% agarose gel. This fragment is designated herein "F1".

The plasmid was digested with the restriction enzymes BamHI and Asp718 dephosphorylated using calf intestinal phosphatase. The DNA was then isolated from a 1% agarose gel using a commercially available kit ("GeneClean" BIO 101 Inc., La Jolla, Ca.). This vector DNA was designated herein "V1".

Fragment F1 and the dephosphorylated plasmid V1 were ligated together with T4 DNA ligase. *E. coli* HB101 cells were transformed with the ligation mixture and spread on culture plates. Other suitable *E. coli* hosts such as XL-1 Blue (Stratagene Cloning Systems, La Jolla, CA) may also be used. Bacteria were identified that contain the plasmid with the human TR2 sequences using the PCR method, in which one of the primers that was used to amplify the gene and the second primer was from well within the vector so that only those bacterial

colonies containing TR2 gene fragments show amplification of the DNA. The sequence of the cloned fragment was confirmed by DNA sequencing. The plasmid was designated herein pBacTR2-T.

Five µg of pBacTR2-T was co-transfected with 1.0 µg of a commercially available linearized baculovirus DNA ("BaculoGold™ baculovirus DNA", Pharmingen, San Diego, CA.), using the lipofection method described by Felgner *et al.*, *Proc. Natl. Acad. Sci. USA* 84:7413-7417 (1987). 1 µg of BaculoGold™ virus DNA and 5 µg of plasmid pBacTR2-T were mixed in a sterile well of a microtiter plate containing 50 µl of serum-free Grace's medium (Life Technologies Inc., Gaithersburg, MD). Afterwards, 10 µl Lipofectin plus 90 µl Grace's medium were added, mixed and incubated for 15 minutes at room temperature. Then the transfection mixture was added drop-wise to Sf9 insect cells (ATCC CRL 1711) seeded in a 35 mm tissue culture plate with 1 ml Grace's medium without serum. The plate was rocked back and forth to mix the newly added solution. The plate was then incubated for 5 hours at 27°C. After 5 hours the transfection solution was removed from the plate and 1 ml of Grace's insect medium supplemented with 10% fetal calf serum was added. The plate was put back into an incubator and cultivation was continued at 27°C for four days.

After four days the supernatant was collected and a plaque assay was performed, as described by Summers and Smith, *supra*. An agarose gel with "Blue Gal" (Life Technologies Inc., Gaithersburg) was used to allow easy identification and isolation of gal-expressing clones, which produce blue-stained plaques. (A detailed description of a "plaque assay" of this type can also be found in the user's guide for insect cell culture and baculovirology distributed by Life Technologies Inc., Gaithersburg, page 9-10). After appropriate incubation, blue stained plaques were picked with the tip of a micropipettor (e.g., Eppendorf). The agar containing the recombinant viruses was then resuspended in a microcentrifuge tube containing 200 µl of Grace's medium and the suspension containing the recombinant baculovirus was used to infect Sf9 cells seeded in 35

mm dishes. Four days later the supernatants of these culture dishes were harvested and then they were stored at 4°C. The recombinant virus is called V-TR2-T.

To verify the expression of the gene used, Sf9 cells were grown in Grace's medium supplemented with 10% heat inactivated FBS. The cells were infected with the recombinant baculovirus V-TR2-T at a multiplicity of infection ("MOI") of about 2. Six hours later the medium was removed and replaced with SF900 II medium minus methionine and cysteine (available from Life Technologies Inc., Rockville, MD). Forty-two hours later, 5 µCi of <sup>35</sup>S-methionine and 5 µCi <sup>35</sup>S-cysteine (available from Amersham) were added to radiolabel proteins. The cells were further incubated for 16 hours and then they were harvested by centrifugation. The proteins in the supernatant as well as the intracellular proteins were analyzed by SDS-PAGE followed by autoradiography. Microsequencing of the amino acid sequence of the amino terminus of purified protein was used to determine the amino terminal sequence of the mature protein and thus the cleavage point and length of the secretory signal peptide.

***Example 2(b): Cloning and Expression of TR2 Protein in a Baculovirus Expression System***

Similarly to the cloning and expression of the truncated version of the TR2 receptor described in Example 2(a), recombinant baculoviruses were generated which express the full length TR2 receptor protein shown in FIG. 1A-1B (SEQ ID NO:2).

In this example, the plasmid shuttle vector pA2 was used to insert the cloned DNA encoding the complete protein, including its naturally associated secretory signal (leader) sequence, into a baculovirus to express the mature TR2 protein. Other attributes of the pA2 vector are as described for the pA2 GP vector used in Example 2(a).

The cDNA sequence encoding the full length TR2 protein in the deposited clone, including the AUG initiation codon and the naturally associated leader sequence shown in FIG. 1A-1B (SEQ ID NO:2), was amplified using PCR oligonucleotide primers corresponding to the 5' and 3' sequences of the gene. The 5' primer has the sequence:

5' GCGCGGATCCACCATGGAGCCTCCTGGAGACTGG 3' (SEQ ID NO:18) containing the underlined BamHI restriction enzyme site, an efficient signal for initiation of translation in eukaryotic cells, as described by Kozak, M., *J. Mol. Biol.* 196:947-950 (1987), followed by 21 bases of the sequence of the complete TR2 protein shown in FIG. 1A-1B, beginning with the AUG initiation codon. The 3' primer has the sequence:

5' GCGCGGTACCTCTACCCCAGCAGGGGCGCCA 3' (SEQ ID NO:19) containing the underlined, Asp718 restriction site followed by 21 nucleotides complementary to the 3' noncoding sequence in FIG. 1A-1B.

The amplified fragment was isolated and digested with restriction enzymes as described in Example 2(a) to produce plasmid pBacTR2

5 µg of pBacTR2 was co-transfected with 1 µg of BaculoGold™ (Pharmingen) viral DNA and 10 µl of Lipofectin™ (Life Technologies, Inc ) in a total volume of 200 µl serum free media. The primary viruses were harvested at 4-5 days post-infection (pi), and used in plaque assays. Plaque purified viruses were subsequently amplified and frozen, as described in Example 2(a).

For radiolabeling of expressed proteins, Sf9 cells were seeded in 12 well dishes with 2.0 ml of a cell suspension containing  $0.5 \times 10^6$  cells/ml and allowed to attach for 4 hours. Recombinant baculoviruses were used to infect the cells at an MOI of 1-2. After 4 hours, the media was replaced with 1.0 ml of serum free media depleted for methionine and cysteine (-Met/-Cys). At 3 days pi, the culture media was replaced with 0.5 ml -Met/-Cys containing 2 µCi each [ $^{35}$ S]-Met and [ $^{35}$ S]-Cys. Cells were labeled for 16 hours after which the culture media was removed and clarified by centrifugation (Supernatant). The cells were lysed in the

dish by addition of 0.2 ml lysis buffer (20 mM HEPES, pH 7.9; 130 mM NaCl; 0.2 mM EDTA; 0.5 mM DTT and 0.5% vol/vol NP-40) and then diluted up to 1.0 ml with dH<sub>2</sub>O (Cell Extract). 30 µl of each supernatant and cell extract were resolved by 15% SDS-PAGE. Protein gels were stained, destained, amplified, dried and autoradiographed. Labeled bands corresponding to the recombinant proteins were visible after 16-72 hours exposure.

### *Example 3*

#### *Cloning and Expression of TR2 in Mammalian Cells*

A typical mammalian expression vector contains the promoter element, which mediates the initiation of transcription of mRNA, the protein coding sequence, and signals required for the termination of transcription and polyadenylation of the transcript. Additional elements include enhancers, Kozak sequences and intervening sequences flanked by donor and acceptor sites for RNA splicing. Highly efficient transcription can be achieved with the early and late promoters from SV40, the long terminal repeats (LTRS) from Retroviruses, e.g., RSV, HTLVI, HIVI and the early promoter of the cytomegalovirus (CMV). However, cellular elements can also be used (e.g., the human actin promoter). Suitable expression vectors for use in practicing the present invention include, for example, vectors such as PSVL and PMSG (Pharmacia, Uppsala, Sweden), pRSVcat (ATCC 37152), pSV2dhfr (ATCC 37146) and pBC12MI (ATCC 67109). Mammalian host cells that could be used include, human HeLa 293, H9 and Jurkat cells, mouse NIH3T3 and C127 cells, Cos 1, Cos 7 and CV 1, quail QC1-3 cells, mouse L cells and Chinese hamster ovary (CHO) cells.

Alternatively, the gene can be expressed in stable cell lines that contain the gene integrated into a chromosome. The co-transfection with a selectable marker such as dhfr, gpt, neomycin, or hygromycin allows the identification and isolation of the transfected cells.

5 The transfected gene can also be amplified to express large amounts of the encoded protein. The DHFR (dihydrofolate reductase) marker is useful to develop cell lines that carry several hundred or even several thousand copies of the gene of interest. Another useful selection marker is the enzyme glutamine synthase (GS) (Murphy *et al.*, *Biochem J.* 227:277-279 (1991); Bebbington *et al.*, *Bio/Technology* 10:169-175 (1992)). Using these markers, the mammalian cells are grown in selective medium and the cells with the highest resistance are selected. These cell lines contain the amplified gene(s) integrated into a chromosome. Chinese hamster ovary (CHO) and NSO cells are often used for the production of proteins.

10 The expression vectors pC1 and pC4 contain the strong promoter (LTR) of the Rous Sarcoma Virus (Cullen *et al.*, *Molecular and Cellular Biology*, 438-447 (March, 1985)) plus a fragment of the CMV-enhancer (Boshart *et al.*, *Cell* 41:521-530 (1985)). Multiple cloning sites, e.g., with the restriction enzyme cleavage sites BamHI, XbaI and Asp718, facilitate the cloning of the gene of interest. The vectors contain in addition the 3' intron, the polyadenylation and termination signal of the rat preproinsulin gene.

### *Example 3(a): Cloning and Expression in COS Cells*

20 The expression plasmid, pTR2 HA, is made by cloning a cDNA encoding TR2 into the expression vector pcDNAI/Amp or pcDNAIII (which can be obtained from Invitrogen, Inc.).

25 The expression vector pcDNAI/amp contains: (1) an *E. coli* origin of replication effective for propagation in *E. coli* and other prokaryotic cells; (2) an ampicillin resistance gene for selection of plasmid-containing prokaryotic cells; (3) an SV40 origin of replication for propagation in eukaryotic cells; (4) a CMV promoter, a polylinker, an SV40 intron; (5) several codons encoding a hemagglutinin fragment (i.e., an "HA" tag to facilitate purification) followed by a termination codon and polyadenylation signal arranged so that a cDNA can be

conveniently placed under expression control of the CMV promoter and operably linked to the SV40 intron and the polyadenylation signal by means of restriction sites in the polylinker. The HA tag corresponds to an epitope derived from the influenza hemagglutinin protein described by Wilson *et al.*, *Cell* 37:767 (1984).  
5 The fusion of the HA tag to the target protein allows easy detection and recovery of the recombinant protein with an antibody that recognizes the HA epitope. pcDNAIII contains, in addition, the selectable neomycin marker.

A DNA fragment encoding a TR2 is cloned into the polylinker region of the vector so that recombinant protein expression is directed by the CMV promoter. The plasmid construction strategy is as follows. The TR2 cDNA of the deposited clone is amplified using primers that contain convenient restriction sites, much as described above for construction of vectors for expression of TR2 in *E. coli*. Suitable primers include the following, which are used in this example. The 5' primer, containing the underlined BamHI site, a Kozak sequence, an AUG start codon and 6 additional codons of the 5' coding region of the complete TR2 has the following sequence:

10 5' GCGCGGATCCACCATGGAGCCTCCTGGAGACTGG 3' (SEQ ID NO:20).

The 3' primer, containing the underlined XbaI site, a stop codon, HA tag, and 19 bp of 3' coding sequence has the following sequence (at the 3' end):

20 5' GCGCTCTAGATCAAGCGTAGTCTGGGACGTCGT  
ATGGGTAGTGGTTTGGGCTCCTCCC 3' (SEQ ID NO:21).

The PCR amplified DNA fragment and the vector, pcDNAI/Amp, are digested with BamHI and XbaI and then ligated. The ligation mixture is transformed into *E. coli* strain SURE (available from Stratagene Cloning Systems, 11099 North Torrey Pines Road, La Jolla, CA 92037), and the transformed culture is plated on ampicillin media plates which then are incubated to allow growth of ampicillin resistant colonies. Plasmid DNA is isolated from resistant colonies and examined by restriction analysis or other means for the presence of the TR2-encoding fragment.

For expression of recombinant TR2, COS cells are transfected with an expression vector, as described above, using DEAE-DEXTRAN, as described, for instance, in Sambrook *et al.*, *Molecular Cloning: a Laboratory Manual*, Cold Spring Laboratory Press, Cold Spring Harbor, New York (1989). Cells are incubated under conditions for expression of TR2 by the vector.

Expression of the TR2-HA fusion protein is detected by radiolabeling and immunoprecipitation, using methods described in, for example Harlow *et al.*, *Antibodies: A Laboratory Manual, 2nd Ed.*; Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York (1988). To this end, two days after transfection, the cells are labeled by incubation in media containing  $^{35}\text{S}$ -cysteine for 8 hours. The cells and the media are collected, and the cells are washed and lysed with detergent-containing RIPA buffer: 150 mM NaCl, 1% NP-40, 0.1% SDS, 0.5% DOC, 50 mM TRIS, pH 7.5, as described by Wilson *et al.* cited above. Proteins are precipitated from the cell lysate and from the culture media using an HA-specific monoclonal antibody. The precipitated proteins then are analyzed by SDS-PAGE and autoradiography. An expression product of the expected size is seen in the cell lysate, which is not seen in negative controls.

#### *Example 3(b): Cloning and Expression in CHO Cells*

The vector pC4 is used for the expression of TR2 protein. Plasmid pC4 is a derivative of the plasmid pSV2-dhfr (ATCC Accession No. 37146). The plasmid contains the mouse DHFR gene under control of the SV40 early promoter. Chinese hamster ovary- or other cells lacking dihydrofolate activity that are transfected with these plasmids can be selected by growing the cells in a selective medium (alpha minus MEM, Life Technologies) supplemented with the chemotherapeutic agent methotrexate. The amplification of the DHFR genes in cells resistant to methotrexate (MTX) has been well documented (see, e.g., Alt, F. W., Kellems, R. M., Bertino, J. R., and Schimke, R. T., 1978, *J Biol. Chem.* 253:1357-1370, Hamlin, J. L. and Ma, C. 1990, *Biochem. et Biophys. Acta*,



1097:107-143, Page, M. J. and Sydenham, M.A. 1991, *Biotechnology* 9:64-68).

Cells grown in increasing concentrations of MTX develop resistance to the drug by overproducing the target enzyme, DHFR, as a result of amplification of the DHFR gene. If a second gene is linked to the DHFR gene, it is usually co-amplified and over-expressed. It is known in the art that this approach may be used to develop cell lines carrying more than 1,000 copies of the amplified gene(s). Subsequently, when the methotrexate is withdrawn, cell lines are obtained which contain the amplified gene integrated into one or more chromosome(s) of the host cell.

Plasmid pC4 contains for expressing the gene of interest the strong promoter of the long terminal repeat (LTR) of the Rous Sarcoma Virus (Cullen, *et al.*, *Molecular and Cellular Biology*, March 1985:438-447) plus a fragment isolated from the enhancer of the immediate early gene of human cytomegalovirus (CMV) (Boshart *et al.*, *Cell* 41:521-530 (1985)). Downstream of the promoter are BamHI, XbaI, and Asp718 restriction enzyme cleavage sites that allow integration of the genes. Behind these cloning sites the plasmid contains the 3' intron and polyadenylation site of the rat preproinsulin gene. Other high efficiency promoters can also be used for the expression, e.g., the human  $\beta$ -actin promoter, the SV40 early or late promoters or the long terminal repeats from other retroviruses, e.g., HIV and HTLV. Clontech's Tet-Off and Tet-On gene expression systems and similar systems can be used to express the TR2 protein in a regulated way in mammalian cells (Gossen, M., & Bujard, H. 1992, *Proc. Natl. Acad. Sci. USA* 89: 5547-5551). For the polyadenylation of the mRNA other signals, e.g., from the human growth hormone or globin genes can be used as well. Stable cell lines carrying a gene of interest integrated into the chromosomes can also be selected upon co-transfection with a selectable marker such as gpt, G418 or hygromycin. It is advantageous to use more than one selectable marker in the beginning, e.g., G418 plus methotrexate.

The plasmid pC4 is digested with the restriction enzymes BamHI and Asp718 and then dephosphorylated using calf intestinal phosphatase by procedures known in the art. The vector is then isolated from a 1% agarose gel.

The DNA sequence encoding the complete TR2 protein including its leader sequence is amplified using PCR oligonucleotide primers corresponding to the 5' and 3' sequences of the gene. The 5' primer has the sequence:

5' GCGCGGATCCACCATGGAGCCTCCTGGAGACTGG 3' (SEQ ID NO:22) containing the underlined BamHI restriction enzyme site followed by an efficient signal for initiation of translation in eukaryotes, as described by Kozak, M., *J. Mol. Biol.* 196:947-950 (1987), and 21 bases of the coding sequence of TR2 protein shown in FIG. 1A-1B (SEQ ID NO:1). The 3' primer has the sequence: 5' GCGCGGTACCTCTACCCCAGCAGGGGCGCCA 3' (SEQ ID NO:19) containing the underlined Asp718 restriction site followed by 21 nucleotides complementary to the non-translated region of the TR2 gene shown in FIG. 1A-1B (SEQ ID NO:1).

The amplified fragment is digested with the endonucleases BamHI and Asp718 and then purified again on a 1% agarose gel. The isolated fragment and the dephosphorylated vector are then ligated with T4 DNA ligase. *E. coli* HB101 or XL-1 Blue cells are then transformed and bacteria are identified that contain the fragment inserted into plasmid pC4 using, for instance, restriction enzyme analysis.

Chinese hamster ovary cells lacking an active DHFR gene are used for transfection. 5 µg of the expression plasmid pC4 is cotransfected with 0.5 µg of the plasmid pSV2-neo using lipofectin (Felgner *et al.*, *supra*). The plasmid pSV2-neo contains a dominant selectable marker, the neo gene from Tn5 encoding an enzyme that confers resistance to a group of antibiotics including G418. The cells are seeded in alpha minus MEM supplemented with 1 mg/ml G418. After 2 days, the cells are trypsinized and seeded in hybridoma cloning plates (Greiner, Germany) in alpha minus MEM supplemented with 10, 25, or 50 ng/ml of methotrexate plus 1 mg/ml G418. After about 10-14 days single clones are

trypsinized and then seeded in 6-well petri dishes or 10 ml flasks using different concentrations of methotrexate (50 nM, 100 nM, 200 nM, 400 nM, 800 nM). Clones growing at the highest concentrations of methotrexate are then transferred to new 6-well plates containing even higher concentrations of methotrexate (1  $\mu$ M, 2  $\mu$ M, 5  $\mu$ M, 10 mM, 20 mM). The same procedure is repeated until clones are obtained which grow at a concentration of 100 - 200  $\mu$ M. Expression of the desired gene product is analyzed, for instance, by SDS-PAGE and Western blot or by reverse phase HPLC analysis.

#### *Example 4*

##### *Tissue distribution of TR2 mRNA expression*

Northern blot analysis is carried out to examine TR2 gene expression in human tissues, using methods described by, among others, Sambrook *et al.*, cited above. A cDNA probe containing the entire nucleotide sequence of the TR2 protein (SEQ ID NO:1) is labeled with  $^{32}$ P using the *rediprime*<sup>TM</sup> DNA labeling system (Amersham Life Science), according to manufacturer's instructions. After labeling, the probe is purified using a CHROMA SPIN- 100<sup>TM</sup> column (Clontech Laboratories, Inc.), according to manufacturer's protocol number PT1200-1. The purified labeled probe is then used to examine various human tissues for TR2 mRNA.

Multiple Tissue Northern (MTN) blots containing various human tissues (H) or human immune system tissues (IM) are obtained from Clontech and are examined with the labeled probe using ExpressHyb<sup>TM</sup> hybridization solution (Clontech) according to manufacturer's protocol number PT1190-1. Following hybridization and washing, the blots are mounted and exposed to film at -70°C overnight, and films developed according to standard procedures.

### Example 5

#### Example 5(a): Expression and Purification of TR2-Fc(TR2-Ig Fusion Protein) and Cleaved TR2

5 The putative transmembrane domain of translated TR2 receptor was determined by hydrophobicity using the method of Goldman *et al.* (*Ann. Rev. of Biophys. Biophys. Chem.* 15:321-353 (1986)) for identifying nonpolar transbilayer helices. The region upstream of this transmembrane domain, encoding the putative leader peptide and extracellular domain, was chosen for the production of an Fc fusion protein. Primers were designed to PCR the corresponding coding region from HTXBS40 with the addition of a BglII site (single underlined), a Factor Xa protease site and an Asp718I site (double underlined) at the 3' end. PCR with this primer pair (forward 35-mer:

10 5' CAGGAATTTCGCAGCCATGGAGCCTCCTGGAGACTG 3' (SEQ ID NO:23), and reverse primer 53-mer:

15 5' CCATACCCAGGTACCCCTTCCCTCGATAGATCT

20 TGCCTTCGTCACCAGCCAGC 3' (SEQ ID NO:24)), which contains 18 nucleotides of the TR2 coding sequence, resulted in one band of the expected size. This was cloned into COSFclink to give the TR2-Fclink plasmid. The PCR product was digested with EcoRI and Asp718I and ligated into the COSFclink plasmid (Johansen, *et al.*, *J. Biol. Chem.* 270:9459-9471 (1995)) to produce TR2-Fclink.

25 COS cells were transiently transfected with TR2-Fclink and the resulting supernatant was immunoprecipitated with protein A agarose. Western blot analysis of the immunoprecipitate using goat anti-human Fc antibodies revealed a strong band consistent with the expected size for glycosylated TR2-Fc (greater than 47.5 kD). A 15L transient COS transfection was performed and the resulting supernatant was purified (see below). The purified protein was used to immunize mice following DNA injection for the production of mAbs.

CHO cells were transfected with TR2-Fclink to produce stable cell lines. Five lines were chosen by dot blot analysis for expansion and were adapted to shaker flasks. The line with the highest level of TR2-Fc protein expression was identified by Western blot analysis. TR2-Fc protein purified from the supernatant of this line was used for cell binding studies by flow cytometry, either as intact protein or after factor Xa cleavage and biotinylation (see below).

Clone HTXBS40 is an allelic variant of TR2 which differs from the sequence shown in FIG. 1A-1B (SEQ ID NO:1) in that HTXBS40 contains guanine at nucleotide 314, thymine at nucleotide 386 and cytosine at nucleotide 627.

A plasmid suitable for expression of the extracellular domain of TR2 was constructed as follows to immunize mice for the production of anti-TR2 mAbs. The Fc fragment was removed from TR2-Fclink by a BglII/XbaI digestion, Klenow was used to fill in the overhangs, and the blunt ends of the plasmid were religated. The resulting frame shift introduced a stop codon immediately following the amino acids which had originally been introduced into TR2-Fclink by the addition of the BglII site. Thus, the C terminus of the extracellular domain of TR2 is followed by only 2 amino acids (RS) in this constructed (TR2exlink).

***Example 5(b): Purification of TR2-Fc from CHO E1A Conditioned Media Followed by Cleavage and Biotinylation of TR2.***

***Assays***

Product purity through the purification was monitored on 15% Laemmli SDS-PAGE gels run under reducing and non-reducing conditions. Protein concentration was monitored by  $A_{280}$  assuming an extinction coefficient of 0.7 for the receptor and 1.28 for the chimera, both calculated from the sequence. Extinction coefficients were confirmed by AAA.

*Protein G Chromatography of the TR2-Fc Fusion Protein*

5 All steps described below were carried out at 4°C. 15L of CHO conditioned media (CM) (0.2 µ filtered following harvest in cell culture) was applied to a 5 X 10 cm column of Protein G at a linear flow rate of 199 cm/h. The column had been washed with 100 mM glycine, pH 2.5 and equilibrated in 20 mM sodium phosphate, 150 mM sodium chloride, pH 7 prior to sample application. After the CM was loaded the column was washed with 5 column volumes of 20 mM sodium phosphate, 150 mM sodium chloride, pH 7 and eluted with 100 mM glycine, pH 2.5. 435 ml of eluate was immediately neutralized with 3 M Tris, pH 8.5 and 0.2 µ filtered. Based on A<sub>280</sub>, extinction coefficient 1.28, 65 mg of protein was recovered at 0.15 mg/ml.

*Concentration/Dialysis*

15 385 ml of Protein G eluate was concentrated in an Amicon stirred cell fitted with a 30K membrane to 34 ml at a final concentration of 1.7. The concentrate was dialyzed against buffer.

*Factor Xa Cleavage and Purification to Generate Free Receptor*

Six ml (10.2 mg) of TR2-Fc was added to 50 µg of Factor Xa resulting in a 1:200, e:s ratio. The mixture was incubated overnight at 4°C.

*Protein G Chromatography of the Free TR2 receptor*

20 A 1 ml column of Protein G was equilibrated in 20 mM sodium phosphate, 150 mM sodium chloride, pH 6.5 in a disposable column using gravity flow. The cleaved receptor was passed over the column 3 times after which the column was washed with 20 mM sodium phosphate, 150 mM sodium chloride, pH 6.5 until no A<sub>280</sub> absorbance was seen. The column was eluted with 2.5 ml of 100 mM glycine, pH 2.5 neutralized with 83 µl of 3 M Tris, pH 8.5. TR2 eluted in the nonbound fraction

25

### *Concentration*

The nonbound fraction from the Protein G column, about 12 ml, was concentrated in a Centricon 10K cell (Amicon) to about 1 ml to a final concentration of 3.5 mg/ml estimated by  $A_{280}$ , extinction coefficient 0.7.

### 5 *Mono S Chromatography*

The concentrated sample was diluted to 5 ml with 20 mM sodium phosphate, pH 6 and applied to a 0.5 X 5 cm Mono S column equilibrated in 20 mM sodium phosphate, pH 6 at a linear flow rate of 300 cm/h. The column was washed with 20 mM sodium phosphate, pH 6 and eluted with a 20 column volume linear gradient of 20 mM sodium phosphate, pH 6 to 20 mM sodium phosphate, 1 M sodium chloride, pH 6. TR2 protein eluted in the nonbound fraction.

### *Concentration/Dialysis*

The 3 ml nonbound fraction from the Mono S column was concentrated to 1 ml as above using a Centricon 10K cell and dialyze against 20 mM sodium phosphate, 150 mM sodium chloride, pH 7. The concentration following dialysis was 2.1 mg/ml.

### *Biotinylation*

0.5 mg of TR2 at 2.1 mg/ml was dialyzed against 100 mM borate, pH 8.5. A 20-fold molar excess of NHS-LC Biotin was added and the mixture was left on a rotator overnight at 4°C. The biotinylated TR2 was dialyzed against 20 mM sodium phosphate, 150 mM sodium chloride, pH 7, sterile filtered and stored at -70°C. Biotinylation was demonstrated on a Western blot probed with streptavidin HRP and subsequently developed with ECL reagent.

003030-059040

20

### Example 6

#### ***The Membrane Bound Form of the TR2 Receptor is a TNFR which Induces Lymphocytes Proliferation and Differentiation***

The members of the tumor necrosis factor (TNFR)/nerve growth factor receptor (NGFR) superfamily are characterized by the presence of three to six repeats of a cysteine-rich motif that consists of approximately 30 to 40 amino acids in the extracellular part of the molecule (Mallett, S. and Barclay, A.N., *Immunol. Today* 12:220 (1991)). The crystal structure of TNFR-I showed that the cysteine-rich motif (TNFR domain) was composed of three elongated strands of residues held together by a twisted ladder of disulfide bonds (Banner, D.W. *et al.*, *Cell* 73:431 (1993)). These receptors contain a hinge-like region immediately adjacent to the transmembrane domain, characterized by a lack of cysteine residues and a high proportion of serine, threonine, and proline, which are likely to be glycosylated with O-linked sugars. A cytoplasmic part of these molecules shows limited sequence similarities - a finding which may be the basis for diverse cellular signaling. At present, the members identified from human cells include CD40 (Stamenkovic, I. *et al.*, *EMBO J.* 8:1403 (1989)), 4-1BB (Kwon, B.S. and Weissman, S.M., *Proc. Natl. Acad. Sci. USA* 86:1963 (1989)), OX-40 (Mallett, S. *et al.*, *EMBO J.* 9:1063 (1990)), TNFR-I (Loetscher, H. *et al.*, *Cell* 61:351 (1990); Schall, T.J. *et al.*, *Cell* 61:361 (1990)), TNFR-II (Smith, C.A. *et al.*, *Science* 248:1019 (1990)), CD27 (Van Lier, R.A. *et al.*, *J. Immunol.* 139:1589 (1987)), Fas (Itoh, N. *et al.*, *Cell* 66:233 (1991)), NGFR (Johnson, D. *et al.*, *Cell* 47:545 (1986)), CD30 (Durkop, H. *et al.*, *Cell* 68:421 (1992)) and LTBR (Baens, M. *et al.*, *Genomics* 16:214 (1993)). Viral open reading frames encoding soluble TNFRs have also been identified, such as SFV-T2 (Smith, C.A. *et al.*, *Science* 248:1019 (1990)), Va53 (Howard, S.T. *et al.*, *Virology* 180:633 (1991)), G4RG (Hu, F.-Q. *et al.*, *Virology* 204:343 (1994)) and crmB (Smith, G.L., *J. Gen. Virol.* 74:1725 (1993)).



Recent intensive studies have shown that these molecules are involved in diverse biological activities such as immunoregulation (Armitage, R.J., *Curr. Opin. Immunol.* 6:407 (1994); Smith, C.A. *et al.*, *Cell* 75:959 (1994)), by regulating cell proliferation (Banchereau, J. *et al.*, *Science* 251:70 (1991); Pollok, K.E. *et al.*, *J. Immunol.* 150:771 (1993); Baum, P.R. *et al.*, *EMBO J.* 13:3992 (1994)), cell survival (Grass, H.-J. *et al.*, *Blood* 83:2045 (1994); Torcia, M. *et al.*, *Cell* 85:345-356 (1996)), and cell death (Tartaglia, L.A. *et al.*, *Cell* 74:845 (1993); Gillette-Ferguson, I. and Sidman, C.L., *Eur. J. Immunol.* 24:1181 (1994); Krammer, P.H. *et al.*, *Curr. Opin. Immunol.* 6:279 (1994)).

Because of their biological significance and the diverse membership of this superfamily, we predicted that there would be further members of the superfamily. By searching an EST-data base, we have identified a new member of the TNFR superfamily. We report here the initial characterization of the molecule called TR2.

## ***Material and Methods***

### ***Identification and Cloning of New Members of the TNFR Superfamily***

An expressed sequence tag (EST) cDNA data base, obtained over 500 different cDNA libraries (Adams, M.D. *et al.*, *Science* 252:1651 (1991); Adams, M.D. *et al.*, *Nature* 355:632 (1992)), was screened for sequence similarity with cysteine-rich motif of the TNFR superfamily, using the blastn and tblastn algorithms (Altschul, S.F. *et al.*, *J. Mol. Biol.* 215:403 (1990)). One EST (HT1SB52 - ATCC Accession No. 97059) was identified in a human T cell line library which showed significant identity to TNFR-II at the amino acid level. This sequence was used to clone the missing 5' end by RACE (rapid amplification of cDNA ends) using a 5'-RACE-ready cDNA of human leukocytes (Clontech, PT1155-1. Cat. #7301-1). This sequence matched four further ESTs (HTOBH42, HTOAU65, HLHA49 and HTXBS40). Complete sequencing of these and other cDNAs indicated that they contained an identical open reading frame homologous

to the TNFR superfamily and was named TR2. Analysis of several other ESTs and cDNAs indicated that some cDNAs had additional sequences inserted in the open reading frame identified above, and might represent various partially-spliced mRNAs.

## Cells

The myeloid and B-cell lines studied represent cell types at different stages of the differentiation pathway. KG1a and PLB 985 (Koeffler, H. *et al.*, *Blood* 56:265 (1980); Tucker, K. *et al.*, *Blood* 70:372 (1987)) were obtained from Phillip Koeffler (UCLA School of Medicine), BJA-B was from Z. Jonak (SmithKline Beecham), and TF 274, a stromal cell line exhibiting osteoblastic features, was generated from the bone marrow of a healthy male donor (Tan & Jonak, unpublished). All of the other cell lines were obtained from the American Type Culture Collection (Rockville, MD). Monocytes were prepared by differential centrifugation of peripheral blood mononuclear cells (PBMC) and adhesion to tissue culture dish. CD19<sup>+</sup>, CD4<sup>+</sup> and CD8<sup>+</sup> were isolated from PBMC by immunomagnetic beads (Dynal, Lake Success, NY). Endothelial cells from human coronary artery were purchased from clonetics (Clonetics, CA).

## RNA and DNA Blot Hybridization

Total RNA of adult tissues was purchased from Clontech (Palo Alto, CA), or extracted from primary cells and cell lines with TriReagent (Molecular Research Center, Inc., Cincinnati, OH). 5 to 7.5 µg of total RNA was fractionated in a 1% agarose gel containing formaldehyde, as described (Sambrook *et al.*, *Molecular Cloning*, Cold Springs Harbor (1989)) and transferred quantitatively to Zeta-probe nylon membrane (Biorad, Hercules, CA) by vacuum-blotting. The blots were prehybridized, hybridized with <sup>32</sup>P-labeled XhoI/EcoRI fragment of TR2 or OX-40 probe, washed under stringent conditions and exposed to X-ray films.

High molecular weight human DNA was digested with various restriction enzymes and fractionated in 0.8% agarose gel. The DNA was denatured, neutralized and transferred to nylon membrane and hybridized to  $^{32}\text{P}$ -labeled TR-2 or its variant cDNA.

5

### ***In Situ Hybridization and FISH Detection***

The *in situ* hybridization and FISH detection of TR2 location in human chromosome were performed as previously described (Heng, H.H.Q. *et al.*, *Proc. Natl. Acad. Sci. USA* 89:9509 (1992); Heng, H.H.Q. *et al.*, *Human Molecular Genetics* 3:61 (1994)). FISH signals and the DAPI banding pattern were recorded separately by taking photographs, and the assignment of the FISH mapping data with chromosomal bands was achieved by superimposing FISH signals with DAPI banded chromosome (Heng, H.H.Q. and Tsui, L.-C., *Chromosoma*. 102:325 (1993)).

### ***Production of Recombination TR2-Fc Fusion Proteins***

The 5' portion of the TR2 containing the entire putative open reading frame of extracellular domain was amplified by polymerase chain reaction (Saiki, R.K. *et al.*, *Science* 239:487 (1988)). For correctly oriented cloning, a HindIII site on the 5' end of the forward primer and a BglII site on the 5' end of the reverse primer were created. The Fc portion of human IgG<sub>1</sub> was PCR-amplified from ARH-77 (ATCC) cell RNA and cloned in SmaI site of pGem7 vector (Promega). The Fc fragment including hinge, CH<sub>2</sub>, and CH<sub>3</sub> domain sequences contained a BglII site at its 5' end and an XhoI site at its 3' end. The HindIII-BglII fragment of TR2 cDNA was inserted into the upstream of human IgG<sub>1</sub>Fc and an in frame fusion was confirmed by sequencing. The TR2-Fc fragment was released by digesting the plasmid with HindIII-XhoI and cloned it into pcDNA3 expression plasmid

20

25

The TR2-Fc plasmid, linearized with PvuI, was transferred into NIH 3T3 by the calcium phosphate co-precipitation method. After selection in 400 µg/ml G418, neomycin-resistant colonies were picked and expanded. ELISA with anti-human IgG<sub>1</sub> and Northern analysis with <sup>32</sup>P-labeled TR2 probe were used to select clones that produce high levels of TR2-Fc in the supernatant. In some experiments, a slightly different engineered TR2-Fc produced in Chinese hamster ovary (CHO) cells was used. The TR2-Fc was purified by protein G chromatography, and the amino acid sequence of N-terminus of the TR2-Fc fusion protein was determined by automatic peptide sequencer (ABI). TR2-Fc was used to produce polyclonal rabbit anti-TR2 antibodies.

#### ***Blocking MLR-Mediated PBMC Proliferation***

PBMC were isolated from three healthy adult volunteers by Ficoll gradient centrifugation at 400 x g for 30 minutes. PBMCs were recovered, washed in RPMI 1640 (GIBCO-BRL) supplemented with 10% FBS, 300 µg/ml L-glutamine and 50 µg/ml genetomycin, and adjusted to 1 x 10<sup>6</sup> cells/ml for two donors and to 2 x 10<sup>5</sup> cells/ml for the third donor.

Fifty µl of each cell suspension was added to 96-well (round bottom) plates (Falcon, Franklin Lakes, NS) together with 50 µl of TR2-Fc, IL-5R-Fc, anti-CD4 mAb or control mAb. Plates were incubated at 37°C in 5% CO<sub>2</sub> for 96 hours. One µCi of [<sup>3</sup>H]-methylthymidine (ICN Biomedicals, Costa Mesa, CA) was then added for an additional 16 hours. Cells were harvested and radioactivity was counted.

### ***Results and Discussion***

#### ***TR2 is a New Member of the TNFR Superfamily***

FIG. 1A-1B (SEQ ID NO:2) shows the amino acid sequence of TR2 deduced from the longest open reading frame of one of the isolated cDNAs

(HLHAB49). Comparison with other sequenced cDNAs and ESTs in the database indicated potential allelic variants which resulted in amino acid changes at positions 17 (either Arg or Lys) and 41 (either Ser or Phe) of the protein sequence shown in FIG. 1A-1B (amino acid residues -20 and 5 in SEQ ID NO:2).

The open reading frame encodes 283 amino acids with a calculated molecular weight of 30,417. The TR2 protein was expected to be a receptor. Therefore, the potential signal sequence and transmembrane domain were sought. A hydrophobic stretch of 23 amino acids towards the C terminus (amino acids 201-225) (FIG. 1A-1B) was assigned as a transmembrane domain because it made a potentially single helical span, but the signal sequence was less obvious. The potential ectodomain TR2 was expressed in NIH 3T3 and CHO cells as a Fc-fusion protein, and the N-terminal amino acid sequence of the recombinant TR2-Fc protein was determined in both cases. The N-terminal sequence of the processed mature TR2 started from amino acid 37, indicating that the first 36 amino acids constituted the signal sequence (FIG. 1A-1B).

Using a polyclonal rabbit antibody raised to TR2, the molecular size of natural TR2 was determined to be 38 kD by Western analysis. Since the protein backbone of processed TR2 would be composed of 247 amino acids with an Mr of 26,000, the protein must be modified post-translationally. Two potential asparagine-linked glycosylation sites are located at amino acid positions 110 and 173 (FIG. 1A-1B). Along with the other members of the TNFR family, TR2 contains the characteristic cysteine-rich motifs which have been shown by X-ray crystallography (Banner *et al.*, *Cell* 73:431 (1993)) to represent a repetitive structural unit (Banner, D.W. *et al.*, *Cell* 73:431 (1993)). FIG. 16 shows the potential TNFR domain aligned among TR2 (SEQ ID NO:2), TNFR-I (SEQ ID NO:10), TNFR-II (SEQ ID NO:11), CD40 (SEQ ID NO:12) and 4-1BB (SEQ ID NO:13). TR2 contained two perfect TNFR domain and two imperfect ones.

The TR2 cytoplasmic tail (TR2 cy) appears to be more closely related to those of CD40cy and 4-1BBcy, and does not contain the death domain seen in the

5

**TABLE 2**  
**GENE EXPRESSION OF TR2 AND OX40 IN TISSUES AND CELLS**

SOURCE	GENE	
	TR2	OX-40
<b>TISSUES (adult)</b>		
Brain	+/-	-
Heart	+	-
Lung	+	-
Thymus	++	-
Spleen	+++	-
Liver	+	-
Kidney	+	-
Small Intestine	+++	-
Prostate	++	-
Skeletal Muscle	+/-	-
Ovary	+	-
Pancreas	+	-
Colon	+	-
Thyroid	+	-
Spinal Cord	+	-
Trachea	+	-
Adrenal Gland	+	-
Lymph Node	+++	-
<b>PRIMARY CELLS</b>		
PBL, CD19+	++	-
PBL, CD8+	++	-
PBL, CD8+ (activated)	++	++
PBL	+++	-
PBL, CD4+ (activated)	++	++
Bone Marrow	+	-
Monocyte	++	-
Endothelial	+	-
<b>HEMATOPOIETIC CELL LINES</b>		
<b>Erythroid</b>		
K562	-	-
HEL	+	-
<b>Myeloid</b>		
KG1a (Promyeloblast)	+	+
KG1 (Myeloblast)	±	±
PLB985 (Late myeloblast)	-	-
HL60 (Promyelocyte)	±	-
U937 (Promonocyte)	±	-
THP-1 (Monocyte)	+	-
<b>B-Lymphocyte</b>		
REH (Pre-preB)	±	-
BJA-B (Early B, IgM)	+	-
Raji (Mature B, IgM)	+	-
TM-9 (Mature B, IgG)	-	-
<b>T-Lymphocyte</b>		
Sup-T1 (CD4+)	-	-
Molt-3 (CD4+)	±	-
H9 (CD4+)	+	-
Jurkat (CD4+)	+	+
no entry = not tested, - = not detected, ± to ++ = increasing amounts of RNA detected		

### *TR2 RNA Expression*

A human tissue RNA blot was used to determine tissue distribution of TR2 RNA expression. TR2 RNA was detected in several tissues with a relatively high level in the lung, spleen and thymus (Table 2) but was not detected by this method in the brain, liver or skeletal muscle (Table 2). TR-2 was also expressed in monocytes, CD19<sup>+</sup> B cells, and resting or PMA plus PHA-treated CD4<sup>+</sup> or CD8<sup>+</sup> T cells. It was only weakly expressed in bone marrow and endothelial cells (Tables 2 and 3), although expression was observed in the hematopoietic cell line KG1a (Table 2). For comparison, the tissue distribution of OX-40, another member of the TNFR superfamily, was examined (Table 2). Unlike TR2, OX-40 was not detected in any tissues examined, and was detected only in activated T-cells and KG1a. Several cell lines were negative for TR2 expression, including TF 274 (bone marrow stromal), MG 63 and TE 85 (osteosarcoma), RL 95-2 (endometrial sarcoma), MCF-7 and T-47D (breast cancer cells), BE, HT 29 (colon cancer cells), HTB-11 and IMR-32 (neuroblastoma), although TR2 was found in the rhabdomyosarcoma HTB-82 (data not shown).

Several cell lines were examined for inducible TR2 expression. HL60, U937 and THP1, which belong to the myelomonocytic lineage, all increased TR2 expression in response to the differentiation agents PMA or DMSO. Increases in expression in response to these agents were observed in KG1a and Jurkat cells. In contrast, PMA did not induce TR2 expression in MG63, but unexpectedly TNF- $\alpha$  did.

In almost all cases, the predominant mRNA was approximately 1.7 kb in size, although several higher molecular weight species could be detected in some tissues. While many cDNAs and ESTs which were sequenced contained insertions in the coding region indicative of partial splicing, we only detected one major protein by Western blot, suggesting that if these encode alternate proteins they are not evident in the cells we examined. The abundance of higher MW mRNAs raises the possibility that TR2 may in part be regulated at the level of mRNA maturation.



**TABLE 3**  
**RELATIVE ABUNDANCE (RA) OF TR2 RNA IN**  
**VARIOUS TISSUE AND CELL TYPES**

Tissue or Cell Type	RA	Tissue or Cell Type	RA
Activated Macrophage (LPS)	22	Fetal Heart	1
Breast Lymph Node	5	Fetal Lung	2
B Cell Lymphoma	5	Glioblastoma	1
Activated Monocytes	2	Hypothalamus, Schizophrenia	1
Activated T Cells	3	Infant Brain	2
Activated Neutrophil	2	Lung	2
Tonsils	5	Osteosarcoma	1
Thymus	3	Pancreas Tumor	1
Anergic T-cell	1	Placenta	2
Jurkat T-Cell	3	Small Intestine	1
Raji Cells (Cycloheximide Treated)	3	Smooth Muscle	1
Atrophic Endometrium	1	Stomach	2
Bone Marrow	1	T-Cell Lymphoma	1
Brain	1	T-Cells	1
Breast	1	Testes	3
CD34 Depleted Buffy Coat (Cord Blood)	1	Testes Tumor	2
Cerebellum	1	Tongue	1
Corpus Colosum	1	Umbilical Vein Endothelial Cells	2
Caco-2 Cells (adenocarcinoma, colon)	1	White Fat	3
Fetal Dura Mater	1		

***TR2 Maps at 1P36.2-P36.3***

The FISH mapping procedure was applied to localize the TR2 gene to a specific human chromosomal region. The assignment of a hybridization signal to the short arm of chromosome 1 was obtained with the aid of DAPI banding. A total of 10 metaphase figures were photographed which indicated that the TR2 gene is located on the chromosome 1 region p36.2-p36.3. The TR2 position is in close

proximity with CD30 (Smith, C.A. *et al.*, *Cell* 73:1349-1360 (1993), 4-1BB (Kwon, B.S. *et al.*, *J. Immunol.* 152:2256-2262 (1994); Goodwin, R.G. *et al.*, *Eur. J. Immunol.* 23:2631-2641 (1993), OX-40 (Birkeland, M.L. *et al.*, *Eur. J. Immunol.* 25:926-930 (1995), and TNFR-II (Baker, E. *et al.*, *Cytogenet. & Cell Genet.* 57:117-118 (1991), suggesting that it evolved through a localized gene duplication event. Interestingly, all of these receptors have stimulatory phenotypes in T cells in response to cognate ligand binding, in contrast to Fas and TNFR-I which stimulate apoptosis. This prompted us to test if TR2 might be involved in lymphocyte stimulation.

#### ***TR2-Fc Interfaces with MLR-Mediated Proliferation of PBMC***

To determine the possible involvement of cell surface TR2 with its ligand in lymphocyte proliferation, we examined allogeneic MLR proliferative responses. When TR-2-Fc was added to the culture, a significant reduction of maximal responses was observed ( $p < 0.05$ ). The addition of TR2-Fc at 100  $\mu\text{g/ml}$  inhibited the proliferation up to 53%. No significant inhibition of proliferation was observed with the control IL-5R-Fc. Surprisingly, at high concentrations (10-100  $\mu\text{g/ml}$ ) IL-5R-Fc was shown to enhance proliferation. An anti-CD4 mAb assayed simultaneously inhibited MLR-mediated proliferation up to 60%, whereas a control anti-IL-5 mAb failed to inhibit the proliferation. It is well known that a major component of the MLR proliferative response is T cell-dependent; hence, it would appear that inhibiting the interaction of TR2 with its ligand prevents optimal T lymphocyte activation and proliferation. The inhibition of MLR proliferation by TR2-Fc at concentrations of 1-100  $\mu\text{g/ml}$  compares favorably with biological effects seen with other TNFR-Fc superfamily members such as CD40-Fc (unpublished results, Jeremy Harrop).

Hence, we have identified an additional member of the TNF receptor superfamily which either plays a direct role in T cell stimulation or binds to a ligand which can stimulate T cell proliferation through one or more receptors

which may include TR2. Consistent with a direct role for TR2 is the similarity of the cytoplasmic domain with CD40 and 4-1BB. We are currently trying to identify this ligand to which TR2 binds in order to clarify its role.

It will be clear that the invention may be practiced otherwise than as particularly described in the foregoing description and examples.

Numerous modifications and variations of the present invention are possible in light of the above teachings and, therefore, are within the scope of the appended claims.

The entire disclosure of all publications (including patents, patent applications, journal articles, laboratory manuals, books, or other documents) cited herein are hereby incorporated by reference.

***What Is Claimed Is:***

1. An isolated nucleic acid molecule comprising a polynucleotide having a nucleotide sequence at least 95% identical to a sequence selected from the group consisting of:

5 (a) a nucleotide sequence encoding the TR2 receptor having the amino acid sequence at positions from about -36 to about 247 in SEQ ID NO:2;

(b) a nucleotide sequence encoding the TR2 receptor polypeptide having the amino acid sequence at positions from about -35 to about 247 in SEQ ID NO:2;

(c) a nucleotide sequence encoding the TR2 receptor polypeptide having the amino acid sequence at positions from about 1 to about 247 in SEQ ID NO:2;

10 (d) a nucleotide sequence encoding the TR2 receptor having the amino acid sequence encoded by the cDNA clone contained in ATCC Deposit Number 97059;

15 (e) a nucleotide sequence encoding the mature TR2 receptor having the amino acid sequence encoded by the cDNA clone contained in ATCC Deposit Number 97059;

20 (f) a nucleotide sequence encoding the TR2 extracellular domain;

(g) a nucleotide sequence encoding the TR2 transmembrane domain;

25 (h) a nucleotide sequence encoding the TR2 intracellular domain;

(i) a nucleotide sequence encoding the TR2-SV1 receptor having the amino acid sequence at positions from about -36 to about 149 in SEQ ID NO:5;

(j) a nucleotide sequence encoding the TR2-SV1 receptor having the amino acid sequence at positions from about -35 to about 149 in SEQ ID NO:5;

(k) a nucleotide sequence encoding the TR2-SV1 receptor having the amino acid sequence at positions from about 1 to about 149 in SEQ ID NO:5;

(l) a nucleotide sequence encoding the TR2-SV1 receptor having the amino acid sequence encoded by the cDNA clone contained in ATCC Deposit Number 97058;

(m) a nucleotide sequence encoding the mature TR2-SV1 receptor having the amino acid sequence encoded by the cDNA clone contained in ATCC Deposit Number 97058;

(n) a nucleotide sequence encoding the TR2-SV2 receptor having the amino acid sequence in SEQ ID NO:8;

(o) a nucleotide sequence encoding the TR2-SV2 receptor having the amino acid sequence at positions from about 2 to about 136 in SEQ ID NO:8;

(p) a nucleotide sequence encoding the TR2-SV2 receptor having the amino acid sequence encoded by the cDNA clone contained in ATCC Deposit Number 97057; and

(q) a nucleotide sequence complementary to any of the nucleotide sequences in (a), (b), (c), (d), (e), (f), (g), (h), (i), (j), (k), (l), (m), (n), (o) or (p).

2. The nucleic acid molecule of claim 1 wherein said polynucleotide has the nucleotide sequence in SEQ ID NO:1, SEQ ID NO:4 or SEQ ID NO:7.

3. The nucleic acid molecule of claim 1 wherein said polynucleotide has the nucleotide sequence in SEQ ID NO:1 encoding a polypeptide having the

amino acid sequence in SEQ ID NO:2, the nucleotide sequence in SEQ ID NO:4 encoding a polypeptide having the amino acid sequence in SEQ ID NO:5, or the nucleotide sequence in SEQ ID NO:7 encoding a polypeptide having the amino acid sequence in SEQ ID NO:8.

5

4. The nucleic acid molecule of claim 1 wherein said polynucleotide has the nucleotide sequence in SEQ ID NO:1 encoding the mature TR2 receptor having the amino acid sequence in SEQ ID NO:2 or the nucleotide sequence in SEQ ID NO:4 encoding the mature TR2-SV1 receptor having the amino acid sequence in SEQ ID NO:5.

10

5. The nucleic acid molecule of claim 1 wherein said polynucleotide has the complete nucleotide sequence of the cDNA clone contained in any one of ATCC Deposit Numbers 97059, 97058 or 97057.

15

6. The nucleic acid molecule of claim 1 wherein said polynucleotide has the nucleotide sequence encoding TR2 receptors having the amino acid sequence encoded by the cDNA clone contained in any one of ATCC Deposit Numbers 97059, 97058 or 97057.

20

7. The nucleic acid molecule of claim 1 wherein said polynucleotide has the nucleotide sequence encoding mature TR2 receptors having the amino acid sequence encoded by the cDNA clone contained in any one of ATCC Deposit Numbers 97059 or 97058.

8. An isolated nucleic acid molecule comprising a polynucleotide which hybridizes under stringent hybridization conditions to a polynucleotide having a nucleotide sequence identical to a nucleotide sequence in (a), (b), (c), (d), (e), (f), (g), (h), (i), (j), (k), (l), (m), (n), (o), (p) or (q) of claim 1 wherein said

polynucleotide which hybridizes does not hybridize under stringent hybridization conditions to a polynucleotide having a nucleotide sequence consisting of only A residues or of only T residues.

5 9. An isolated nucleic acid molecule comprising a polynucleotide which encodes the amino acid sequence of an epitope-bearing portion of a TR2 receptor having an amino acid sequence in (a), (b), (c), (d), (e), (f), (g), (h), (i), (j), (k), (l), (m), (n), (o), (p) or (q) of claim 1.

10 10. The isolated nucleic acid molecule of claim 9, which encodes an epitope-bearing portion of a TR2 receptor selected from the group consisting of: a polypeptide comprising amino acid residues from about 3 to about 34 in SEQ ID NO:2; a polypeptide comprising amino acid residues from about 70 to about 84 in SEQ ID NO:2; a polypeptide comprising amino acid residues from about 106 to about 153 in SEQ ID NO:2; a polypeptide comprising amino acid residues from about 240 to about 247 in SEQ ID NO:2; a polypeptide comprising amino acid residues from about 3 to about 34 in SEQ ID NO:5; a polypeptide comprising amino acid residues from about 63 to about 100 in SEQ ID NO:5; a polypeptide comprising amino acid residues from about 135 to about 149 in SEQ ID NO:5; a polypeptide comprising amino acid residues from about 56 to about 68 in SEQ ID NO:8; and a polypeptide comprising amino acid residues from about 93 to about 136 in SEQ ID NO:8.

11. The isolated nucleic acid molecule of claim 1, which encodes a TR2 receptor extracellular domain.

12. The isolated nucleic acid molecule of claim 1, which encodes a TR2 receptor transmembrane domain

13. The isolated nucleic acid molecule of claim 1, which encodes a TR2 receptor intracellular domain.

14. A method for making a recombinant vector comprising inserting an isolated nucleic acid molecule of claim 1 into a vector.

15. A recombinant vector produced by the method of claim 14.

16. A method of making a recombinant host cell comprising introducing the recombinant vector of claim 15 into a host cell.

17. A recombinant host cell produced by the method of claim 16.

18. A recombinant method for producing a TR2 polypeptide, comprising culturing the recombinant host cell of claim 17 under conditions such that said polypeptide is expressed and recovering said polypeptide.

19. An isolated nucleic acid molecule comprising a polynucleotide having a nucleotide sequence in SEQ ID NO:1 wherein nucleotide 314 is either guanine or adenine, nucleotide 386 is either thymine or cytosine, and nucleotide 627 is either thymine or cytosine.

20. An isolated nucleic acid molecule comprising a polynucleotide having a nucleotide sequence encoding the TR2 receptor having the amino acid sequence in SEQ ID NO:2 wherein amino acid number -20 is either lysine or arginine and amino acid number 5 is either serine or phenylalanine.

21. An isolated TR2 polypeptide having an amino acid sequence at least 95% identical to a sequence selected from the group consisting of



(a) the amino acid sequence of the TR2 polypeptide having the amino acid sequence at positions from about -36 to about 247 in SEQ ID NO:2;

(b) the amino acid sequence of the TR2 polypeptide having the amino acid sequence at positions from about -35 to about 247 in SEQ ID NO:2;

(c) the amino acid sequence of the TR2 polypeptide having the amino acid sequence at positions from about 1 to about 247 in SEQ ID NO:2;

(d) the amino acid sequence of the TR2 polypeptide having the amino acid sequence encoded by the cDNA clone contained in ATCC Deposit Number 97059;

(e) the amino acid sequence of the mature TR2 polypeptide having the amino acid sequence encoded by the cDNA clone contained in ATCC Deposit Number 97059;

(f) the amino acid sequence of the TR2 receptor extracellular domain;

(g) the amino acid sequence of the TR2 receptor transmembrane domain;

(h) the amino acid sequence of the TR2 receptor intracellular domain;

(i) the amino acid sequence encoding the TR2-SV1 receptor having the amino acid sequence at positions from about -36 to about 149 in SEQ ID NO:5;

(j) the amino acid sequence encoding the TR2-SV1 receptor having the amino acid sequence at positions from about -35 to about 149 in SEQ ID NO:5;

(k) the amino acid sequence encoding the TR2-SV1 receptor having the amino acid sequence at positions from about 1 to about 149 in SEQ ID NO:5;

(l) the amino acid sequence encoding the TR2-SV1 receptor having the complete amino acid sequence encoded by the cDNA clone contained in ATCC Deposit Number 97058;

(m) the amino acid sequence encoding the mature TR2-SV1 receptor having the amino acid sequence encoded by the cDNA clone contained in ATCC Deposit Number 97058;

(n) the amino acid sequence encoding the TR2-SV2 receptor having the amino acid sequence in SEQ ID NO:8;

(o) the amino acid sequence encoding the TR2-SV2 receptor having the amino acid sequence at positions from about 2 to about 136 in SEQ ID NO:8;

(p) the amino acid sequence encoding the TR2-SV2 receptor having the amino acid sequence encoded by the cDNA clone contained in ATCC Deposit Number 97057; and

(q) the amino acid sequence of an epitope-bearing portion of any one of the polypeptides of (a), (b), (c), (d), (e), (f), (g), (h), (i), (j), (k), (l), (m), (n), (o) or (p).

22. An isolated polypeptide comprising an epitope-bearing portion of a TR2 receptor protein, wherein said portion is selected from the group consisting of: a polypeptide comprising amino acid residues from about 3 to about 34 in SEQ ID NO:2; a polypeptide comprising amino acid residues from about 70 to about 84 in SEQ ID NO:2; a polypeptide comprising amino acid residues from about 106 to about 153 in SEQ ID NO:2; a polypeptide comprising amino acid residues from about 240 to about 247 in SEQ ID NO:2; a polypeptide comprising amino acid residues from about 3 to about 34 in SEQ ID NO:5; a polypeptide comprising amino acid residues from about 63 to about 100 in SEQ ID NO:5; a polypeptide comprising amino acid residues from about 135 to about 149 in SEQ ID NO:5; a polypeptide comprising amino acid residues from about 56 to about 68 in SEQ

ID NO:8; or a polypeptide comprising amino acid residues from about 93 to about 136 in SEQ ID NO:8.

23. An isolated antibody that binds specifically to a TR2 receptor polypeptide of claim 21.

24. A method of treating herpes simplex viral infection comprising introducing an effective amount of a soluble fragment of a TR2 polypeptide into an individual to be treated in admixture with a pharmaceutically acceptable carrier.

25. A method of treating a disease state associated with aberrant cell survival comprising introducing an effective amount of a TR2 protein, or agonist or antagonist thereof, into an individual to be treated in admixture with a pharmaceutically acceptable carrier.

26. A method of screening for agonists and antagonists of TR2 activity comprising:

- (a) contacting cells which express TR2 polypeptides with a candidate compound,
- (b) assaying a cellular response, and
- (c) comparing the cellular response to a standard cellular response made in absence of the candidate compound; whereby, an increased cellular response over the standard indicates that the compound is an agonist and a decreased cellular response over the standard indicates that the compound is an antagonist.

## Human Tumor Necrosis Factor Receptor-Like 2

### *Abstract*

5 The present invention relates to novel members of the Tumor Necrosis  
Factor family of receptors. The invention provides isolated nucleic acid molecules  
encoding a human TR2 receptor and two splice variants thereof. TR2  
polypeptides are also provided as are vectors, host cells and recombinant methods  
for producing the same. The invention further relates to screening methods for  
identifying agonists and antagonists of TR2 receptor activity. Also provided are  
diagnostic methods for detecting disease states related to the aberrant expression  
of TR2 receptors. Further provided are therapeutic methods for treating disease  
states related to aberrant proliferation and differentiation of cells which express  
the TR2 receptors.  
10

P:\USERS\STEPHENW\NHGS\1488.077\A78-05.WPD

10 30 50  
 GCACGAGCTGCCTCCCGCAGGCGCCACCTGTGTCCCCAGCGCCGCTCCACCCAGCAGGC  
 70 90 110  
 CTGAGCCCCCTCTCTGCTGCCAGACACCCCTGCTGCCCCTCTCTCTGCTGCTCGGGTTCT  
 130 150 170  
 GAGGCACAGCTTGTACACCGAGGCGGATTCTCTTTCTCTTTCTCTTTCTCTGGCCC  
 190 210 230  
 ACAGCCGCGAGCAATGGCGCTGAGTTCCTCTGCTGGAGTTCATCTGCTAGCTGGGTTCCT  
 250 270 290  
 GAGCTGCCGGTCTGAGCCTGAGGCATGGAGCCTCCTGGAGACTGGGGGCTCCTCCCTGG  
 M E P P G D W G P P P W  
 310 330 350  
 AGATCCACCCCCAAAACCGACGTCTTGAGGCTGGTGCTGTATCTCACCTTCTGGGAGCC  
 R S T P K T D V L R L V L Y L T F L G A  
 370 390 410  
 CCCTGCTACGCCCCAGCTCTGCCGTCTGCAAGGAGGACGAGTACCCAGTGGGCTCCGAG  
 P C Y A P A L P S C K E D E Y P V G S E  
 430 450 470  
 TGCTGCCCCAAGTGCAGTCCAGGTTATCGTGTGAAGGAGGCTGCGGGGAGCTGACGGGC  
 C C P K C S P G Y R V K E A C G E L T G  
 490 510 530  
 ACAGTGTGTGAACCCTGCCCTCCAGGCACCTACATTGCCCACCTCAATGGCCTAAGCAAG  
 T V C E P C P P G T Y I A H L N G L S K  
 550 570 590  
 TGTCTGCAGTGCCAAATGTGTGACCCAGCCATGGGCCTGCGCGGAGCCGGAAGTCTCC  
 C L Q C Q M C D P A M G L R A S R N C S  
 610 630 650  
 AGGACAGAGAACGCCGTGTGTGGTGTGAGCCAGGCCACTTCTGCATCGTCCAGGACGGG  
 R T E N A V C G C S P G H F C I V Q D G  
 670 690 710  
 GACCACTGCGCCGCGTGCCGCGCTTACGCCACCTCCAGCCCGGGCCAGAGGGTGCAGAAG  
 D H C A A C R A Y A T S S P G Q R V Q K  
 730 750 770  
 GGAGGCACCGAGAGTCAGGACACCCCTGTGTGAGAACTGCCCCCGGGGACCTTCTCTCCC  
 G G T E S Q D T L C Q N C P P G T F S P  
 790 810 830  
 AATGGGACCCTGGAGGAATGTCAGCACCAGACCAAGTGCAGCTGGCTGGTGACGAAGGCC  
 N G T L E E C Q H Q T K C S W L V T K A  
 850 870 890  
 GGAGCTGGGACCAGCAGCTCCCACTGGGTATGGTGGTTTCTCTCAGGGAGCCTCGTCATC  
 G A G T S S S H W V W W F L S G S L V I  
 910 930 950  
 GTCATTGTTTGTCTCCACAGTTGGCCTAATCATATGTGTGAAAAGAAGAAAGCCAAGGGGT  
 V I V C S T V G L I I C V K R R K P R G  
 970 990 1010

Figure 1A

GATGTAGTCAAGGTGATCGTCTCCGTCCAGCGGAAAAGACAGGAGGCAGAAGGTGAGGCC  
 D V V K V I V S V Q R K R Q E A E G E A  
 1030 1050 1070  
 ACAGTCATTGAGGCCCTGCAGGCCCCCTCCGGACGTACCACGGTGGCCGTGGAGGAGACA  
 T V I E A L Q A P P D V T T V A V E E T  
 1090 1110 1130  
 ATACCCTCATTACGGGGAGGAGCCCAAACCACTGACCCACAGACTCTGCACCCCGACGC  
 I P S F T G R S P N H \*  
 1150 1170 1190  
 CAGAGATACCTGGAGCGACGGCTGAATGAAAGAGGCTGTCCACCTGGCGGAACCACCGGA  
 1210 1230 1250  
 GCGCGGAGGCTTGGGGCTCCACCCTGGACTGGCTTCCGTCTCCTCCAGTGGAGGGAGAG  
 1270 1290 1310  
 GTGGCGCCCCCTGCTGGGGTAGAGCTGGGGACGCCACGTGCCATTCCCATGGGCCAGTGAG  
 1330 1350 1370  
 GGCCTGGGGCCTCTGTTCTGCTGTGGCCTGAGCTCCCCAGAGTCTTGAGGAGGAGCGCCA  
 1390 1410 1430  
 GTTGCCCCTCGCTCACAGACCACACACCCAGCCCTCCTGGGCCAACCAGAGGGCCTTCA  
 1450 1470 1490  
 GACCCAGCTGTGTGCGCGTCTGACTCTTGTGGCCTCAGCAGGACAGGCCCCGGGCACTG  
 1510 1530 1550  
 CCTCACAGCCAAGGCTGGACTGGGTGGCTGCAGTGTGGTGTTTAGTGGATAACCACATCG  
 1570 1590 1610  
 GAAGTGATTTTCTAAATTGGATTGAATTCGGCTCCTGTTTTCTATTGTCATGAAACAG  
 1630 1650 1670  
 TGATTTTGGGGAGATGCTGTGGGAGGATGTAAATATCTTGTTCCTCAAAAAAAAAA  
 1690  
 AAAAAAAAAAAAAAAAAAAAAA

Figure 1B

Percent Similarity: 46.591    Percent Identity: 28.788

00340690 062999

```
1 MEPPGDWGGPPWRSTPKTDVLRRLVLYLTFLGAPCYAPALPSCKEDEYPVG 50
      ...: ||. ... | .... .. ..|...:| :
1 .....MVSLPRLCALWGCLLTAVHLGQCVTCSDKQYLHD 34

51 SECCPKCSPGYRVKEACGELTGTVCEPCPPGTYYIAHLNGLSKCLQCQMCD 100
      ::||. | || |... |... | |...|...: |::| :| | . |:
35 GQCCDLCQPGSRLTSHCTALEKTQCHPCDSGEFSAQWNREIRCHQHRHCE 84

101 PAMGLRASRNCSRTENAVCGCSPGHFCIVQGDHCAACRAYATISSPGQRV 150
      |. |||..... :...|||...|: |. .| |...| ..... || |
85 PNQGLRVKKEGTAESDTVCTCKEGQHCTSKD...CEACAQHTPCIPGFGV 131

151 QKGGTESQDTLCQNCPPGTFSPNGTL.EECQHQTKE.SWLVTKAGAGTSS 198
      .:|... ||::|: ||. | || .:| |... |... |... |... |||
132 MEMATETTTDTVCHPCPVGFFSNQSSLFKCYPWTSCEDKNLEVLQKGTSQ 181

199 SH.....WVWWFLSGSLVIVIVCSTVGLIICVKR..RKPRGDVVKVIV 239
      .: ..... :::: | .:|...|...|...|...: |...|...: .
182 TNVICGLKSRMRALLVIPVVMGILITIFGVFLYIKKVVKPKDNEMLPPA 231

240 SVQRKRQEAEG.....EATVIEALQAPPDVTTVAVEETIPSFIGRSPNH 283
      . ....|| | : ..... |. |...|...| |...| . | |...
232 ARRQDPQEMEDYPGHNTAAPVQETLHGCQPVTQEDGKESRISVQERQVTD 281
```

Figure 2

4




Figure 3



10 30 50  
 CCCCCTTCTACAGGAAACCCGGAGTGGACTGGAACGGTGCAGGGGAGAACTCGCCCCCTC  
 70 90 110  
 CCATCGGGCGCCTCCTTCATACCGGCCCTTCCCCCTCGGCTTTGCCTGGACAGCTCCTGCC  
 130 150 170  
 TCAGGCAGCGCCACCTGTGTGCGCCAGCGCCGCTCCACCCAGCAGGCCTGAGCCCCCTCTC  
 190 210 230  
 TGCTGCCAGACACCCCTGCTGCCCCACTACTCCTGCTGCTCGGGTTCTGAGGCACAGCTT  
 250 270 290  
 GTCACACCGAGGCGGATTCTCTTTCTCTTTCTCTTTCTCTTCTGGCCACAGCCGAGCA  
 310 330 350  
 ATGGCGCTGAGTTCTCTGCTGGAGTTCATCCTGCTAGCTGGGTTCCTGAGCTGCCGGTC  
 370 390 410  
 TGAGCCTGAGTCATGGAGCCTCCTGGAGACTGGGGCCTCCTCCCTGGAGATCCACCCCC  
 M E P P G D W G P P P W R S T P  
 430 450 470  
 AGAACCGACGTCTTGAGGCTGGTGTGTATCTCACCTTCCTGGGAGCCCCCTGCTACGCC  
 R T D V L R L V L Y L T F L G A P C Y A  
 490 510 530  
 CCAGCTCTGCCGTCTGCAAGGAGGACGAGTACCCAGTGGGCTCCGAGTGCTGCCCAAG  
 P A L P S C K E D E Y P V G S E C C P K  
 550 570 590  
 TGCAGTCCAGGTTATCGTGTGAAGGAGCCTGCGGGGAGCTGACGGGCACAGTGTGTGAA  
 C S P G Y R V K E A C G E L T G T V C E  
 610 630 650  
 CCCTGCCCTCCAGGCACCTACATTGCCACCTCAATGGCCTAAGCAAGTGTCTGCAAGTGC  
 P C P P G T Y I A H L N G L S K C L Q C  
 670 690 710  
 CAAATGTGTGACCCAGATATTGGTTCCCCCTGTGACCTCAGGGGAAGAGGTCACCTGGAG  
 Q M C D P D I G S P C D L R G R G H L E  
 730 750 770  
 GCTGGTGCCACCTGAGTCCAGGCAGACAGAAAGGGGAACCAGACCCAGAGGTGGCCTTT  
 A G A H L S P G R Q K G E P D P E V A F  
 790 810 830  
 GAGTCACTGAGCGCAGAGCCTGTCCATGCGCCAACGGCTCTGTCCCCTTGGAGCCTCAT  
 E S L S A E P V H A A N G S V P L E P H  
 850 870 890  
 GCCAGGCTCAGCATGGCCAGTGCTCCCTGCGGCCAGGCAGGACTGCACCTGCGGGACAGG  
 A R L S M A S A P C G Q A G L H L R D R  
 910 930 950  
 GCTGACGGCACACCTGGGGGCAGGGCCTGAGCCTACAGGGAGGCACAGGGCAGGTGGGCT  
 A D G T P G G R A \*  
 970 990 1010  
 AGCCATGAACAGAAGAGGAAGCTGGAGTGCTTTGGGGGTTTCATGCATGTAGGCTGGGATT

Figure 4A

1030	1050	1070
TGGGGCTCACACCTCAACCTGCATGCCCAGTTCCATGCCCCCTCCCCTCTTGTGAAAGCAC		
1090	1110	1130
CTGTCTACTTGGGCTGAGGATGTGGGGGCACAGGTGGCAGGTGAGGCTGCCCTCAGGAGG		
1150	1170	1190
GGCCCAGGCCAGCTTGTACCCACCTCCACCAGTACCTGAAGAAGTGGGGCTCTCACCC		
1210	1230	1250
TACCTGCCTCTGCCATTGGAATGGCCTGGTTTGCACAGATGGGAAACCCGTTTGAGGGGT		
1270	1290	1310
GGGTGTCTGGGTGGGCACGTGGGGCGAGGACCTGCCTGAGGGACCCTGCCCTGGAAGTGA		
1330	1350	1370
CAGTGCAAGCTCGGCGTCTTGCCATCTGGGCAGAAGGCTGGTTTCTCCCATCAACGAAG		
1390	1410	1430
CCCTCCCAGGACCTTCTGCAAGCCCTCGTCCCACACGCAGCTCTGCCGTCCCTTGGTGT		
1450	1470	1490
CCCTCCCGGCTCAGGTCTCCATGCTGGGTACCTCTGGGCACCTCGTTTGGCTGAGCCA		
1510	1530	1550
GGGGTTCAGCCTGGCAGGGCGCCCTGGCAGCAGTCTTGGCCTGTGGATGCTGTCTTGGC		
1570	1590	1610
CTGTGGATGGTGTCCCGCCCTCCACGTACCCCTCTCACCCCTCCTCTTGGACTCCAGCC		
1630	1650	1670
ATGGGCCTGCGCGCGAGCCGGAAGTGTCCAGGACAGAGAAGCCGTGTGTGGCTGCAGC		
1690	1710	1730
CCAGGCCACTTCTGCATCGTCCAGGACGGGAGCCACTGCGCCGCGTGCCGCGCTTACGCC		
1750	1770	1790
ACCTCCAGCCCCGGGCCAGAGGGTGCAGAAGGGAGGCACCGAGAGTCAGGACACCCTGTGT		
1810	1830	1850
CAGAACTGCCCCCGGGGACCTTCTCTCCCAATGGGACCCTGGAGGAATGTCAGCACCAG		
1870	1890	1910
ACCAATTGGCCTAATCATATGTGTGAAAAGAAGAAAGCCAAGGGGTGAGCACACGGTGGC		
1930	1950	1970
CCCATCAGGGTTCATGTCCCCAGCCGTCACCTCTTGGAGCTCTGTCAACCCCAAGCCTGGG		
1990	2010	2030
AGGTGGCCCCAGAGCTTTTCCAGGATCCGCGGCTCCTCCCAGGGCAGCCACTGCAGGCTG		
2050	2070	2090
GGGCAGGTGTATGTAGTCAAGGTGATCGTCTCCGTCCAGCGGTAAAAGACAGGAGGCAGA		
2110	2130	2150
AGGTGAGGCCACAGTCAATTGAGCCCTGCAGGCCCTCCGGACGTACCCACGGTGGCCGTG		
2170	2190	2210
GAGGAGACAATACCCCTCATTACGGGGAGGAGCCCAAACCACTGACCCACAGACTCTGCA		
2230	2250	2270
CCCCGACGCCAGAGATACCTGGAGAGACGGCTGCTGATAGAGGCTGTCCACCTGGCGAAA		
2290	2310	2330
CCACCGGAGCCCCGAGGCTTGGGGGCTCCGCCCTGGGCTGGTTTCCGTCTCCTCCAGTGG		
2350	2370	2390
AGGGAGAGGTGGTGGCCCTGCTGGTGGTAGAGCTGGGGACGCCACGTGCCATTCCCATGG		
2410	2430	2450
TTCACTGAGGGGCTGGTGGCCTCTGTTCTGCTGTGGCCTGAGCTCCCCAGAGTCTTGAGG		
2470	2490	2510
AGGAGCCCCAGTTGCCCCCTCGCTCACAGACCACACCCAGCCCTCCTGGGGCCAACCCAG		
2530	2550	2570
AGGCCCTTCAGACCCCAGCTGTCTGCGCGTCTGACTCTTGTGGCCTCAGCAGGACAGGC		
2590	2610	2630
CCCGGGCACTGCCTCACAGCCAAGGCTGGAATGGGTGGCTGCAGTGTGGTGTGTTAGTGG		
2650	2670	2690
ATACCACATCGGAAGTGATTTTCTAAAAATTGGATTGAATTCGGAAAAAA		

Figure 4B






Figure 6

```

10          30          50
AAAGCTCGGGCTCCACCGGGGACGACCGCTCCTAGAACTGAGTGGTATCCCCCGGGCCT
70          90          110
GCAGGAATTCCAACCTGCCTGAAGGGACCCCTGCCCTGGAAGTACAGTGAAGCTCGGCG
130         150         170
TCCTGCCCATCTGGGAAGAAGGCTGGTTTCTCCCATCAACGAAGCCCTCCCAGGACCTTC
190         210         230
CTGCAAGCCCTCGTCCCACACGCAGCTCTGCCGTCCCTTGGTGTCCCTCCCGGCTCAGG
250         270         290
TCCTCCATGCTGGGTACCTCTGGGCACCTCGTTTGGCTGAGCCAGGGGTTTACGCTGGCA
M L G T S G H L V W L S Q G F S L A
310         330         350
GGGCGCCCTGGCAGCAGTCCTTGGCCTGTGGATGCTGTCTGGCCTGTGGATGGTGTCCC
G R P G S S P W P V D A V L A C G W C P
370         390         410
GGCCTCCACGTACCCCTCTCAGCCCTCCTCTTGGACTCCAGCCATGGGCCTGCGCGCG

G L H V P P L S P S S W T P A M G L R A
430         450         470
AGCCGGAAGTGTCCAGGACAGAGAACGCCGTGTGTGGCTGCAGCCCAGGCCACTTCTGC
S R N C S R T E N A V C G C S P G H F C
490         510         530
ATCGTCCAGGACGGGGACCACTGCGCCGCGTGCCGCGCTTACGCCACCTCCAGCCCGGGG
I V Q D G D H C A A C R A Y A T S S P G
550         570         590
CAGAGGGTGCAGAAGGGAGGCACCGAGAGTCAGGACACCCTGTGTGAGAAGTGCACCCCGG
Q R V Q K G G T E S Q D T L C Q N C P R
610         630         650
GGACCTTCTCTCCCAATGGGACCCTGGAGGAATGTCAGCACCAGACCAAGTAAGTGAACC
G P S L P M G P W R N V S T R P S K *
670         690         710
CGGGGGAGGCCAGCTCTGTGCCCTGGGGAGGGGGCTCCACGTTGCTTCCCTGGGAGATGA
730         750         770
CCGTCTTCTCCAGCAGAAAGGTTGAAGGTCCCACCCTGAGCGGCACCCCTGGTCACATGCC
790         810         830
TGCGTCCAGGAGAGCTGCAGGGTGAAGCCTGTGTGCCCCAGATAACCCCTTCCATGGGCC
850         870         890
CAGACAAAGCCTCATCAGATCTGAGCTTCCCTGGAGGCTCAGGATGGGCCTTCCCAGAAGC
910         930         950
AGGCCCAGAGGGAGGCTGCCTCCAGATCCCTGTCCCTGGGGCTGTGGGTGTCCCTGAA
970         990         1010
TGTCAGGGCCATGGGAGGGCCCTGGGCTTACAGGGTTGGGGAAAGTGAACACTCTGCTC
1030        1050        1070
TTTGTCCACCTTCGGGAGGACAACCTTCAAATGCTGACCCTGGGCCCCCTAACTGACCTGA
1090        1110        1130
GACTTCAGAGCTTCTTGGGAGGAGCTGGGGTCCCCCAGCGGAGCCTGGGATGGAGCAGGG
1150        1170        1190
ATGGCTGCCCCAGGGAGGGGGCGGTGGGGCCTTCCATCCTGCTCTGCCCTCCTCGTCCTC
1210        1230        1250
TGGCCCCAGCTCAGTCCTGTCCATCTCCAGCTCTAACCATTTGTGGCCCCGACACTGGCTC
1270        1290        1310
TCCCTCTACCTTCTGTCTTGTCTGACACTGGTCTCCCGTCTCTGGGGTCTCTGCACTG
1330        1350        1370
ATGGCTGCCTCCCGCTTCTCTCCCTCTCCCTCTGCCGTCTGTCTCCTGTGGCCAGTCT

```

Figure 7A

1390	1410	1430
CTCCTTGTTTCTCTTCTCCTCCTTCCTTCTCTCCACCTCCCCATAGCCGAGCTTGGA		
1450	1470	1490
GTCAGACAGACCTCTGAGGTCTCATCTGGAGCTGCCACCAGCCCAGCCTCCCTGGGACC		
1510	1530	1550
TGTCTTCACTGCCTGGGGCCCTGGGAGCCAGGGAGGCTCCCTGAGGCTGAGTGAACACTG		
1570	1590	1610
GGCGCTGCACCTGCCTCTCCACGTCTCGGCCCCACTCCCGCAGGTGCAGCTGGCTGGT		
1630	1650	1670
GACGAAGCCCGGAGCTGGGACCAGCAGCTCCCACTGGGTATGGTGGTTTCTCTCAGGGAG		
1690	1710	1730
CCTCGTCATCGTCATTGTTTGCTCCACAGTTGGCCTAATCATATGTGTGAAAAGAAGAAA		
1750	1770	1790
GCCAAGGGGTGATGTAGTCAAGGTGATCGTCTCCGTCCAGGTATTGATCCTCCTCCCCCT		
1810	1830	1850
CTCCCTCCCCCTCCACCTTCCACCTCCCTCTCCCCGCTGGGGCTGGTGTTCCTGGTG		
1870	1890	1910
TACATGGTGGGGGCTCCCAGTTCTCTGAGGGTCTGAGTCTTTCAAGTACAGCCACGGTA		
1930	1950	1970
GCTCAGGAAAGAACCACCCCCCTCAAAGTAAAGCAGTAAATGAACCCGAGAACCCTGGA		
1990	2010	2030
GTCCCAGGGGGCCTGAGCAGGCAGGGTCTCCACGATTTCGTGTGCTCACAGCGGAAAAG		
2050	2070	2090
ACAGGAGGCAGAAGGTGAGGCCACAGTCATTGAGGCCCTGCAGGCCCTCCGGACGTCAC		
2110	2130	2150
CACGGTGGCCGTGGAGGAGACAATACCCTCATTACGGGGGAGGAGCCCAAACCACTGAC		
2170	2190	2210
CCACAGACTCTGCACCCCGACGCCAGAGATACCTGGAGCGACGGCTGCTGAAAGAGGCTG		
2230	2250	2270
TCCACCTGGCGAAACCACCGGAGCCCGAGGTTTGGGGGCTCCGCCCTGGGCTGGTTTCC		
2290	2310	2330
GTCTCCTCCAGTGGAGGGAGAGGTGGGGCCCCCTGCTGGGGTAGAGCTGGGGACGCCACGT		
2350	2370	2390
GCCATTCCCAGTGGGCCAGTGAGGGCCTGGGGCCTCTGTTCTGCTGTGGCCTGAGCTCCCC		
2410	2430	2450
AGAGTCTTGAGGAGGAGCGCCAGTTGCCCTCGCTCACAGACCACACCCAGCCCTCCT		
2470	2490	2510
GGGTCCAGCCCAGAGGGCCCTTCAGACCCAGCTGTCTGCGCGTCTGACTCTTGTTGGCCT		
2530	2550	2570
CAGCAGGACAGGCCCCGGGCACTGCCTTCAAGCCAAGGCTGGACTGGGTGGCTGCAGTG		
2590	2610	2630
TGGTGTTTAGTGGATACCACATCGGAAGTGATTTTCTAAATTGGATTTGAAAAAAA		

Figure 7B

Percent Similarity: 45.522    Percent Identity: 26.866

```

1  MLGTSGHLVWLSQGFSL.....AGRPGSSPWPVD..... 29
   :. : : . |. |:.|      |. |||...: :
1  .MAPVAVWAALAVGLELWAAHALPAQVAFTPYAPEPGSTCRLREYYDQT 49

30  AVLACGWC.PGLHV.....PPLSPSSW 50
   | :. |:.| || |.      ..|||.:.|:
50  AQMCCSKCSPGQHAKVFCTKTSDTVCDSCEDSTYTQLWNWVPECLSCGSR 99

51  TPAMGLRASRNC SRTE NAVCGCSPGHFCIVQGDHCAACRAYATSSPGQR 100
   ... :. ....|. |:.| :|. |. || :| : ..: | | :. ...||
100  CSSDQV.ETQACTREQNRIC TCRPGWYCALSKQEGCRLCAPLRKCRPGFG 148

101  VQKGGTESQDTLCQNCPRGPSLPMGPWRNVSTRPSK..... 136
   |:.| |||. |:.|. |:.|.      ... ::: ..
149  VARPGTETSDVVCKPCAPGTFSNTTSSTDICRPHQICNVVAIPGNASMDA 198

```

Figure 8

09340690 06904660




Figure 9



Percent Similarity: 73.370      Percent Identity: 59.783

[illegible]

666290-06904E50

Percent Similarity: 70.588    Percent Identity: 60.294

```
1 MEPPGDWGWPPWRSTPKTDVLRVLVLYLTFLGAPCYAPALPSCKEDEYPVG 50
      . . . . . : : : | . . . . : : :
1 . . . . . MLGTSGHLVWLSQGFSLAGRPGSSP . . . . . WPVD 29
51 SECCPKCSPGYRVKEACGELTGTVCEPCPPGTYIAHLNGLSKCLQCQMCD 100
      . . . : | : : | . . . : : :
30 AVLACGWCPGLHV . . . . . PPLSPSSW . . . . . T 51
101 PAMGLRASRNC SRTENAVCGCSPGHFCIVQGDHCAACRAYATSSPGQRV 150
      |||
52 PAMGLRASRNC SRTENAVCGCSPGHFCIVQGDHCAACRAYATSSPGQRV 101
151 QKGGTESQDTLCQNCPPGTFSPNGTLEECQHQTCSWLVTKAGAGTSSSH 200
      |||
102 QKGGTESQDTLCQNCPRGPSLPMGPWRNV . . STRPSK . . . . . 136
```

Figure 11

00340690-06999

Percent Similarity: 37.984    Percent Identity: 20.155

```
1 MEPPGDWGPWPWRSTPRTDVLRLVLYLTFLGAPCYAP.....ALPCK 43
      . . . . . : | : : | | . . | | . . |
1 . . . . . MLGTSGHLVWLSQGFSLAGRPGSSPWPVDAVLACGWCP 38
44 EDEYPVGSECCPKCSPGYRVKEACGELTGTVCEPCPPGTYIAHLNGLSKC 93
      : . | . | : : . . | . | . . | : . . | | : : | . : | : : . . |
39 GLHVPPLSPSSWTPAMGLRASRNCSTENAVCGCSPGHFCI..VQGDHC 86
94 LQCMCDPDIGSPCDLRGRGHLEAGAHLSPPGRQKGEPDPEVAFESLSAEP 143
      . | . : . . : | | : : . : | | . . | : . . | . : : . : | . |
87 AACRAYAT..SSPGQRVQKGGTESQDTLCQNCPRGPSLPMGPWRNVSTRP 134
144 VHAANGSVPLEPHARLSMASAPCGQAGLHLRDRADGTPGGRA. 185
      |
135 SK..... 136
```

Figure 12





```

1309 TGGGCCAGTGAGGGCCTGG.GGCCTCTGTTCTGCTGTGGCCTGAGCTCCC 1357
    ||| ||||| ||||| ||||| ||||| ||||| ||||| ||||| |||||
2398 TGGTTCAGTGAGGGGCTGGTGGCCTCTGTTCTGCTGTGGCCTGAGCTCCC 2447
    .
1358 CAGAGTCCTGAGGAGGAGCGCCAGTTGCCCTCGCTCACAGACCACACAC 1407
    ||||| ||||| ||||| ||||| ||||| ||||| ||||| |||||
2448 CAGAGTCCTGAGGAGGAGCCCCAGTTGCCCTCGCTCACAGACCACACAC 2497
    .
1408 CCAGCCCTCCTGGGCCAACCCAGAGG.GCCTTCAGACCCCAGCTGTGTGC 1456
    ||||| ||||| ||||| ||||| ||||| ||||| ||||| |||||
2498 CCAGCCCTCCTGGGCCAACCCAGAGGCCCTTCAGACCCCAGCTGTCTGC 2547
    .
1457 GCGTCTGACTCTTGTGGCCTCAGCAGGACAGGCCCCGGGCACTGCCTCAC 1506
    ||||| ||||| ||||| ||||| ||||| ||||| ||||| |||||
2548 GCGTCTGACTCTTGTGGCCTCAGCAGGACAGGCCCCGGGCACTGCCTCAC 2597
    .
1507 AGCCAAGGCTGGACTGGGTTGGCTGCAGTGTGGTGTGTTAGTGGATAACCAC 1556
    ||||| ||||| ||||| ||||| ||||| ||||| ||||| |||||
2598 AGCCAAGGCTGGAATGGGTTGGCTGCAGTGTGGTGTGTTAGTGGATAACCAC 2647
    .
1557 ATCGGAAGTGATTTTCT..AAATTGGATTGAATTTCGGCTCCTGTTTTCT 1604
    ||||| ||||| ||||| ||||| ||||| ||||| ||||| |||||
2648 ATCGGAAGTGATTTTCTAAAAATTGATTGAATTTCGGAACAAAAA..... 2692

```

Figure 13C

```

1 .....GCACGAGCTGCCTCCCGCAGGCGC 24
    | | | | |
701 GTTGCTTCCCTGGGAGATGACCGTCTTCTCCAGCAGAAAGGTTGAAGGTC 750
    | | | | |
25 CACCTGTGTCCCCAGCGCCGCTCCACCCAGCAGGCCTGAGCCCCTCTCT 74
    | | | | |
751 CCACCCTGAGCGGCACCCTGGTCACATGCCTGCGTCCAGGAGAGCTGCAG 800
    | | | | |
75 GCTGCCAGACACCCCTGCTGCCCCTCTCCTGCTGCTCGGGTTCTGAGG 124
    | | | | |
801 GGTGAAGCCTGTGTGCCCCAGATAACCCCTTCCATGGGCCCCAGACAAAGC 850
    | | | | |
125 CACAGCTTGTACACCGAGGCGGATTCTCTTTCTCTTTCTCTTTCTCTTC 174
    | | | | |
851 CTCATCAGATCTGAGCTTCTGGAGGCTCAGGATGGGCCTTCCCAGAAGC 900
    | | | | |
175 TGGCCCACA.....GCCGCAGCAATGGCGCTGAGTTCCTCTGCTGGAGTT 219
    | | | | |
901 AGGCCCAGAGGGAGGCTGCCTCCAGATCCCCTGTCCCCTGGGGCTGTGGG 950
    | | | | |
220 CATCCTGCTAGCTGGGTTCCCGAGCTGCCGGTCTGAGCCTGAGGCATGGA 269
    | | | | |
951 TGTCCCTGAATGTCAGGGCCATGGGAGGGCCCCTGGGCTTCAGGGGTGG 1000
    | | | | |
270 GCCTCCTGGAGACTGGGGGCCTCCTCC.....CTGGAGATCCACCCCCAA 314
    | | | | |
1001 GGAAAGTGAACACTCTGCTCTTTGTCCACCTTCGGGAGGACAACCTTCAA 1050
    | | | | |
315 A.....ACCGACGTCTTGAGGCTGGTGTGTATCTCACCTTCCTGGGA 357
    | | | | |
1051 ATGCTGACCCTGGGCCCCCTAACTGACCTGAGACTTCAGAGCTTCTTGGGA 1100

```

Figure 14A

06904E60 06904E60 06904E60

```

358 G C C C C C T G C T A C G C C C C A G C T C T G C C G T C C T G C A A G G A G G A C G A G T A C C C 407
    | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
1101 G G A G C T G G G G T C C C C C A G C G G A G C C T G G G A T G G A G C A G G G A T G G C T G C C C 1150
    . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . .
408 A G T G G G C T C C G A G T G C T G C C C C A A G T G C A G T C C A G G T T A T C G T G T G A A G G 457
    | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
1151 C A G G G A G G G G G C G G T G G G G C C T T C C A T C C T G C T C T G C C C T C C T C G T C C T C 1200
    . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . .
458 A G G . . C C T G C G G G G A G C T G A C G G G C A C A G T G T G T G A A C C C T G C C C T C C A G 505
    | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
1201 T G G C C C C A G C T C A G T C C T G T C C A T C T C C A G C T C T A A C C A T T T G T G G C C C G 1250
    . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . .
506 G C A C C T A C A T T G C C C A C C T C A A T G G C C T A A G C A A G T G T C T G C A G T G C C . . 553
    | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
1251 A C A C T G G C T C T C C C T C T A C C T T C T G T C C T T G T C T G A C A C T G G T C T C C C G T 1300
    . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . .
554 . A A T G T G T G A C C C A G C C A T G G G C C T G C G C G C A G C C G G A A C T G C T C C A G 602
    | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
1301 G C T C T G G G G T C T C T G C A C T G A T G G C T G C C T C C C G C T T C T C T C C C C T C T C C 1350
    . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . .
603 G A C A G A G A A C G C C G T G T G T G G T T G C A G C C C A G G C C A C T T C T G C A T C G T C C 652
    | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
1351 C T C T G C C G T C C T G T C T C C T G T G G C C A G T C T C T C C T T G T T T C T T C T C C T 1400
    . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . .
653 A G G A C G G G G A C C A C T G C G C C G C G T G C C G C G C T T A C G C C A C C T C C A G C C C G 702
    | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
1401 C C T T C C T T C T C T C C A C C T C C C C A T A G C C G A G C T T G G A A A G T C A G A C A G A 1450
    . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . .
703 G G C C A G A G G G T G C A G A A G G G A G G C A C C G A G A G T C A G G A C A C C C T G T G T C A 752
    | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
1451 C C T C T G A G G T C T C A C C T G G A G C T G C C A C C A G C C C A G C C T C C C T G G G A C C 1500
    . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . .
753 G A A C T G C C C . . C C C G G G G A C C T T C T C T C C C A A T G G G A C C C T G G A G G . . . . 796
    | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
1501 T G T C T T C A C T G C C T G G G G C C C T G G G A G C C A G G G A G G C T C C C T G A G G C T G A 1550
    . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . .
797 . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . .
    | | | | |
1551 G T G A A C A C T G G G C G C T G C A C C T G C C T C T C C C A C G T C C T C G G C C C C A C T C C 1600
    . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . .
811 A C C A A G T G C A G C T G G C T G G T G A C G A A G G C C G A G C T G G G A C C A G C A G C T C 860
    | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
1601 C G C A G G T G C A G C T G G C T G G T G A C G A A G C C C G A G C T G G G A C C A G C A G C T C 1650
    . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . .
861 C C A C T G G G T A T G G T G G T T T C T C T C A G G G A G C C T C G T C A T C G T C A T T G T T T 910
    | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
1651 C C A C T G G G T A T G G T G G T T T C T C T C A G G G A G C C T C G T C A T C G T C A T T G T T T 1700
    . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . .
911 G C T C C A C A G T T G G C C T A A T C A T A T G T G T G A A A G A A G A A G C C A A G G G G T 960
    | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
1701 G C T C C A C A G T T G G C C T A A T C A T A T G T G T G A A A G A A G A A G C C A A G G G G T 1750
    . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . .
961 G A T G T A G T C A A G G T G A T C G T C T C C G T C C A G . . . . . . . . . . . . . . . . . . 990
    | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
1751 G A T G T A G T C A A G G T G A T C G T C T C C G T C C A G G T A T T G A T C C T C C T C C C C C T 1800
    . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . .
991 . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . .
    C G G A A A A G A C A G G A G G C A 1008

```

Figure 14B



```

2001 GGCAGGGTCTCCACGATTCGTGTGCTCACAGCGGAAAAGACAGGAGGCA 2050
1009 GAAGGTGAGGCCACAGTCATTGAGGCCCTGCAGGCCCTCCGGACGTCAC 1058
2051 GAAGGTGAGGCCACAGTCATTGAGGCCCTGCAGGCCCTCCGGACGTCAC 2100
1059 CACGGTGGCCGTGGAGGAGACAATACCCTCATTAC.GGGGAGGAGCCCA 1107
2101 CACGGTGGCCGTGGAGGAGACAATACCCTCATTACGCGGGGAGGAGCCCA 2150
1108 AACCCTGAGCCACAGACTCTGCACCCCGACGCCAGAGATACCTGGAGCG 1157
2151 AACCCTGAGCCACAGACTCTGCACCCCGACGCCAGAGATACCTGGAGCG 2200
1158 ACGGCTGAATGAAAGAGGCTGTCCACCTGGCGGAACCACCGGAGCCCGGA 1207
2201 ACGGCTG.CTGAAAGAGGCTGTCCACCTGGCGAAACCACCGGAGCCCGGA 2249
1208 GGCTTGGGGGCTCCACCCTGGACTGGCTTCCGTCTCCTCCAGTGGAGGGA 1257
2250 GGTTTGGGGGCTCCGCCCTGGGCTGGTTTCCGTCTCCTCCAGTGGAGGGA 2299
1258 GAGGTGGCGCCCCCTGCTGGGGTAGAGCTGGGGACGCCACGTGCCATTCCC 1307
2300 GAGGTGGGGCCCCCTGCTGGGGTAGAGCTGGGGACGCCACGTGCCATTCCC 2349
1308 ATGGGCCAGTGAGGGCCTGGGGCCTCTGTTCTGCTGTGGCCTGAGCTCCC 1357
2350 ATGGGCCAGTGAGGGCCTGGGGCCTCTGTTCTGCTGTGGCCTGAGCTCCC 2399
1358 CAGAGTCCTGAGGAGGAGCGCCAGTTGCCCTCGCTCACAGACCACACAC 1407
2400 CAGAGTCCTGAGGAGGAGCGCCAGTTGCCCTCGCTCACAGACCACACAC 2449
1408 CCAGCCCTCCTGGG.CCAACCCAGAGGG.CCTTCAGACCCAGCTGTGTG 1455
2450 CCAGCCCTCCTGGGTCCAGCCAGAGGGCCCTTCAGACCCAGCTGTCTG 2499
1456 CGCGTCTGACTCTTGTGGCCTCAGCAGGACAGGCCCCGGGCACTGCCTCA 1505
2500 CGCGTCTGACTCTTGTGGCCTCAGCAGGACAGGCCCCGGGCACTGCCTTC 2549
1506 CAGCCAAGGCTGGACTGGGTGGCTGCAGTGTGGTGTGTTAGTGGATACCA 1555
2550 AAGCCAAGGCTGGACTGGGTGGCTGCAGTGTGGTGTGTTAGTGGATACCA 2599
1556 CATCGGAAGTGATTTTCTAAATTGGATTTGAATTCGGCTCCTGTTTTCTA 1605
2600 CATCGGAAGTGATTTTCTAAATTGGATTTGAAAAAAA..... 2637

```

Figure 14C

06904660

Percent Similarity: 53.479    Percent Identity: 53.479

1 CCCCCTTCTACAGGAAACCCGGAGTGGACTGGAACGGTGCAGGGGGAGAA 50  
| |    |    | | | | | | |    |    | | |

Figure 15A



749 AGAAAGG.....GGAACCAGACCCAGAGGTGGCCTTTGAGTCACTGAGCG 793  
 ||||| || ||| | ||| ||| |||  
 735 AGAAAGGTTGAAGGTCCCACCCTGAGCGGCACCCTGGTCACATGCCTGCG 784  
 794 CAGAGCCTGTCCATGCGGCCAACGGCTCTGTCCCCTTGGAGCCTCATGCC 843  
 || || | ||| ||| ||| ||| ||| |||  
 785 TCCAGGAGAGCTGCAGGGTGAAGCCTGTGTGCCCCAGATAACCCCTTCCA 834  
 844 AGGCTCAGCATGGCCAGTGCTCCCTGCGGCCAGGCAGGACTGCACCTGCG 893  
 || || | ||| ||| ||| ||| ||| |||  
 835 TGGGCCCAGACAAAGCCTCATCAGATCTGAGCTTCCTGGAGGCTCAGGAT 884  
 894 GGACAGGGCTGACGGCACACCTGGGGGCAGGGCCTGAGCCTACAGGGAGG 943  
 || || | ||| ||| ||| ||| ||| |||  
 885 GGGCCTTCCCAGAAGCAGGCCAGAGGGAGGCTGCCTCCAGATCCCCTGT 934  
 944 CACAGGGCAGGTGGGCTAGCCATGAACAGAAGAGGAAGCTGGAGTGCTTT 993  
 || || | ||| ||| ||| ||| ||| |||  
 935 CCCCTGGGGCTGTGGGTGTCCTGAATGTCAGGGCCATGGGAGGGCCCT 984  
 994 GGGGGTTTCATGCATGTAGGCTGGGATTTGGGGCTCACACCTCAACCTGCA 1043  
 || || | ||| ||| ||| ||| ||| |||  
 985 GGGCTTCAGGGGTGGGGAAAGTGAACACTCTGCTCTTTGTCCACCTTCG 1034  
 1044 TGCCCAGTTCCATGCCCTCCCCTCTTGTGAAAGCACCTGTCTACTTGGG 1093  
 || || | ||| ||| ||| ||| ||| |||  
 1035 GGAGGACAACCTTCAAATGCTGACCCTGGGCCCTAACT.....GA 1075  
 1094 CTGAGGATGTGGGGGCACAGGTGGCAGGTGAGGCTGCCCTCAGGAGGGGC 1143  
 || || | ||| ||| ||| ||| ||| |||  
 1076 CCTGAGACTTCAGAGCTTCTTGGGAGGAGCTGGGGTCCCCCAGCGGAGCC 1125  
 1144 CCAGGGCCAGCTTGTACCCACCTCCACCAGTACCTGAAGAAGTGGGGCT 1193  
 || || | ||| ||| ||| ||| ||| |||  
 1126 TGGGATGGAGCAGGGATGGCTGCCCCA.....GGGAGGGGGCGGTGG 1167  
 1194 CTCACCTACCTGCCTCTGCCATTGGAATGGCCTGGTTTGCACAGATGGG 1243  
 || || | ||| ||| ||| ||| ||| |||  
 1168 GGCCTTCCATCCTGCTCTGCCCTCCTCGTCTCTGGCCCCAGCTCAGTCC 1217  
 1244 AAACCCGTTTGAGGGGTGGGTGTCTGGGTGGGCACGTGGGGCGAGGACCT 1293  
 || || | ||| ||| ||| ||| ||| |||  
 1218 TGTCCATCTCCAGCTCTAACCATTTGTGGCCCGACACTGGCTCTCCCTCT 1267  
 1294 GCCTGAGGGACCTTGCCCTGGAAGTACAGTGCAAGCTCGGCGTCTGCTGCC 1343  
 || || | ||| ||| ||| ||| ||| |||  
 1268 ACCTTCTGTCTTGTCTGACACTGGTCTCCCGTGTCTGGGGTCTCTGCA 1317  
 1344 CATCTGGGCAGAAGGCTGGTTTCTCCCATCAACGAAGCCCTCCCAGGACC 1393  
 || || | ||| ||| ||| ||| ||| |||  
 1318 CTGATGGCTGCCTCCCGCTTCTCTCCCTCTCCCTCTGCGTCTCTGCTC 1367  
 1394 TTCCTGCAAGCCCTCGTCCCACACGCAGCTCTGCCGTCCCTTGGTGTCCC 1443  
 || || | ||| ||| ||| ||| ||| |||  
 1368 CTGTGGCCAGTCTCTCCTTGTCTCTCTCTCTCTCTCTCTCTCTCTCTCT 1417  
 1444 TCCCGGCCTCAGGTCCTCCA....TGCTGGGTACCTCTGGGCACCTCGTT 1489  
 |||| | | | | | ||||| | |

Figure 15C



2182 AC.GGGGAGGAGCCCAAACCACTGACCCACAGACTCTGCACCCCGACGCC 2230  
 || |||||  
 2135 ACGGGGGAGGAGCCCAAACCACTGACCCACAGACTCTGCACCCCGACGCC 2184  
 2231 AGAGATACCTGGAGAGACGGCTGCTGATAGAGGCTGTCCACCTGGCGAAA 2280  
 |||||  
 2185 AGAGATACCTGGAGCGACGGCTGCTGAAAGAGGCTGTCCACCTGGCGAAA 2234  
 2281 CCACCGGAGCCCGGAGGCTTGGGGGCTCCGCCCTGGGCTGGTTTCCGTCT 2330  
 |||||  
 2235 CCACCGGAGCCCGGAGGTTTGGGGGCTCCGCCCTGGGCTGGTTTCCGTCT 2284  
 2331 CCTCCAGTGGAGGGAGAGGTGGTGCCCTGCTGGTGGTAGAGCTGGGGAC 2380  
 |||||  
 2285 CCTCCAGTGGAGGGAGAGGTGGGGCCCTGCTGG.GGTAGAGCTGGGGAC 2333  
 2381 GCCACGTGCCATTCCCATGGTTCAGTGAGGGGCTGGTGGCCTCTGTTCTG 2430  
 |||||  
 2334 GCCACGTGCCATTCCCATGGGCCAGTGAGGGCCTGG.GGCCTCTGTTCTG 2382  
 2431 CTGTGGCCTGAGCTCCCCAGAGTCTTGAGGAGGAGCCCCAGTTGCCCTC 2480  
 |||||  
 2383 CTGTGGCCTGAGCTCCCCAGAGTCTTGAGGAGGAGCGCCAGTTGCCCTC 2432  
 2481 GCTCACAGACCACACACCCAGCCCTCCTGGG.CCAACCCAGAGGCCCTT 2529  
 |||||  
 2433 GCTCACAGACCACACACCCAGCCCTCCTGGGTCCAGCCAGAGGGCCCTT 2482  
 2530 CAGACCCAGCTGTCTGCGCGTCTGACTCTTGTGGCCTCAGCAGGACAGG 2579  
 |||||  
 2483 CAGACCCAGCTGTCTGCGCGTCTGACTCTTGTGGCCTCAGCAGGACAGG 2532  
 2580 CCCCAGGCACTGCCTCACAGCCAAGGCTGGAATGGGTGGCTGCAGTGTG 2629  
 |||||  
 2533 CCCCAGGCACTGCCTTCAAGCCAAGGCTGGACTGGGTGGCTGCAGTGTG 2582  
 2630 GTGTTTAGTGGATACACATCGGAAGTGATTTTCTAAAAATTGGATTTGA 2679  
 |||||  
 2583 GTGTTTAGTGGATACACATCGGAAGTGATTTTCTAAA.....TTGG 2624  
 2680 ATTCGGAAAAAAA 2692  
 |||||  
 2625 ATTTGAAAAAAA 2637

Figure 15E

TNFR-I	V	C	PQGKYIHPQNNSI	C	C	TK	C	HKGYLYND	C	PGPGQDTD	C	R
TNFR-II	T	C	RLREYYDQTAQM	C	C	SK	C	SPGQHAKVF	C	TKTSDTV	C	D
CD40	A	C	REKQYLINSQ	C	C	SL	C	QPGQKLVSD	C	TEPTEIE	C	L
4-1BB	-	-	-	-	-	SN	C	PAGTF	C	DNNRNQI	C	S
TR-2	S	C	KEDEYFPVGSE	C	C	PK	C	SPGYRVKEA	C	GELTGTV	C	E

TNFR-I	E	C	ESGSFTASENHJRH	C	LS	C	SK	C	RKEMQVEISS	C	TVDRDTV	C	G
TNFR-II	S	C	EDSTYTQLMNWVPE	C	LS	C	GSR	C	SSDQVETQA	C	TREQNRI	C	T
CD40	P	C	GESEFLDTWNRETH	C	HQ	H	KY	C	DPNLGLRVQOK	G	TSETDTI	C	T
4-1BB	P	C	PENSFSAGGQRT	C	DI	C	RQ	C	KGVFTRKE	C	SSTSNAE	C	D
TR-2	P	C	PPGTYYAHLNGLSK	C	LQ	C	QM	C	DPAMGLRASRN	C	SRTENAV	C	G

TNFR-I	C	R	QYRHYWSENLFQ	C	FN	C	SL	C	LNCTVHLS	C	QEKQNTV	C	T		
TNFR-II	C	R	PGWY	C	ALSKQEG	C	RL	C	APLRK	C	RPFGVARP	G	TETSDV	C	K
CD40	C	EEGWH	C	TSEA	C	ES	C	VLHRS	C	SPGFGVKQIAT	G	VSDTI	C	E	
4-1BB	C	TPGFH	C	LGAG	C	SM	C	EQD	C	KQSQELTKKG	C	KD	C	-	
TR-2	C	SPGHF	C	IVQDGDH	C	AA	C	RAYAT	S	SPQRVQXG	G	TESQDTL	C	Q	

TNFR-I	-	C	HAGFFLRENE	C	VS	C	SN	C	KKSLE	C	TKL	C	L
TNFR-II	P	C	APGTFSTTSSTDI	C	RP	H	QI	C	NVVAIP	G	NASMDAV	C	T
CD40	P	C	FVGFFSNVSSAFEK	C	HP	N	TS	C	ETKDLVQQA	G	TNKTDTV	C	G
4-1BB	-	C	F-GTFNKQKRG	C	RP	N	TN	C	SLDGKSVLVN	G	TKERDVV	C	G
TR-2	N	C	PPGTFSPNGTLEE	C	QH	Q	TK	C	SNLVTKA	G	AGTSSSH	W	V

Figure 16

## Combined Declaration and Power of Attorney for Patent Application

Docket Number: 1488.0770004

As a below named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below next to my name.

I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter that is claimed and for which a patent is sought on the invention entitled **Human Tumor Necrosis Factor Receptor-Like 2**, the specification of which is attached hereto unless the following box is checked:

- ☒ was filed on **October 30, 1996**;  
as United States Application Number or PCT International Application Number **08/741,095**; and  
was amended on \_\_\_\_\_ (if applicable).

I hereby state that I have reviewed and understand the contents of the above identified specification, including the claims, as amended by any amendment referred to above.

I acknowledge the duty to disclose information that is material to patentability as defined in 37 C.F.R. § 1.56.

I hereby claim foreign priority benefits under 35 U.S.C. § 119(a)-(d) or § 365(b) of any foreign application(s) for patent or inventor's certificate, or § 365(a) of any PCT international application, which designated at least one country other than the United States listed below, and have also identified below any foreign application for patent or inventor's certificate, or PCT international application having a filing date before that of the application on which priority is claimed.

Prior Foreign Application(s)	Priority Claimed
_____ (Application No.)      _____ (Country)      _____ (Day/Month/Year Filed)	<input type="checkbox"/> Yes <input type="checkbox"/> No
_____ (Application No.)      _____ (Country)      _____ (Day/Month/Year Filed)	<input type="checkbox"/> Yes <input type="checkbox"/> No

I hereby claim the benefit under 35 U.S.C. § 119(e) of any United States provisional application(s) listed below.

_____ (Application No.)	_____ (Filing Date)
_____ (Application No.)	_____ (Filing Date)

I hereby claim the benefit under 35 U.S.C. § 120 of any United States application(s), or under § 365(c) of any PCT international application designating the United States, listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States or PCT international application in the manner provided by the first paragraph of 35 U.S.C. § 112, I acknowledge the duty to disclose information that is material to patentability as defined in 37 C.F.R. § 1.56 that became available between the filing date of the prior application and the national or PCT international filing date of this application.

<u>PCT/US95/05058</u> (Application No.)	<u>April 27, 1995</u> (Filing Date)	<u>Pending</u> (Status - patented, pending, abandoned)
<u>08/462,315</u> (Application No.)	<u>June 5, 1995</u> (Filing Date)	<u>Pending</u> (Status - patented, pending, abandoned)
<u>08/462,962</u> (Application No.)	<u>June 5, 1995</u> (Filing Date)	<u>Pending</u> (Status - patented, pending, abandoned)
<u>08/464,595</u> (Application No.)	<u>June 5, 1995</u> (Filing Date)	<u>Pending</u> (Status - patented, pending, abandoned)



I hereby appoint the following attorney(s) and/or agent(s) to prosecute this application and to transact all business in the Patent and Trademark Office connected therewith:

Robert G. Sterne, Esq., Reg. No. 28,912; Edward J. Kessler, Esq., Reg. No. 25,688; Jorge A. Goldstein, Esq., Reg. No. 29,021; Samuel L. Fox, Esq., Reg. No. 30,353; David K.S. Cornwell, Esq., Reg. No. 31,944; Robert W. Esmond, Esq., Reg. No. 32,893; Tracy-Gene G. Durkin, Esq., Reg. No. 32,831; Michele A. Cimbala, Esq., Reg. No. 33,851; Michael B. Ray, Esq., Reg. No. 33,997; Robert E. Sokohl, Esq., Reg. No. 36,013; Eric K. Steffe, Esq., Registration No. 36,688; Andrea G. Reister, Esq., Registration No. 36,253; Daniel N. Yannuzzi, Esq., Registration No. 36,727; Robert H. Benson, Registration No. 30,446; Paul C. Kimball, Registration No. 34,610; and A. Anders Brookes, Registration No. 36,373.

Send Correspondence to:

STERNE, KESSLER, GOLDSTEIN & FOX P.L.L.C.  
1100 New York Avenue, N.W.  
Suite 600  
Washington, D.C. 20005-3934

Direct Telephone Calls to:

(202) 371-2600

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

Full name of sole or first inventor <b>Jian NI</b>	
Inventor's signature	Date
Residence <b>U.S.A.</b>	
Citizenship <b>China</b>	
Post Office Address <b>5502 Manorfield Road, Rockville, MD 20853</b>	
Full name of second inventor <b>Craig A. ROSEN</b>	
Second inventor's signature	Date
Residence <b>U.S.A.</b>	
Citizenship <b>U.S.A.</b>	
Post Office Address <b>22400 Rolling Hill Road, Laytonsville, MD 20882</b>	

Full name of third inventor <b>Reiner L. GENTZ</b>	
Third inventor's signature	Date
Residence <b>U.S.A.</b>	
Citizenship <b>Germany</b>	
Post Office Address <b>13404 Fairland Park Drive, Silver Spring, MD 20904</b>	
Full name of fourth inventor <b>Sally Doreen Patricia LYN</b>	
Fourth inventor's signature ✓ <i>Sally Lyn</i>	April 2, 97 <sup>X</sup> ✓ Date
Residence <b>U.S.A.</b>	
Citizenship <b>Canada</b>	
Post Office Address <b>134-B Hampton Court, West Chester, PA 19380</b>	
Full name of fifth inventor <b>Mark Robert HURLE</b>	
Fifth inventor's signature ✓ <i>Mark Robert Hurle</i>	3/28/97 ✓ Date
Residence <b>U.S.A.</b>	
Citizenship <b>U.S.A.</b>	
Post Office Address <b>105 Stonybrook Drive, Norristown, PA 19403</b>	

## Combined Declaration and Power of Attorney for Patent Application

Docket Number: 1488.0770004

As a below named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below next to my name.

I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter that is claimed and for which a patent is sought on the invention entitled **Human Tumor Necrosis Factor Receptor-Like 2**, the specification of which is attached hereto unless the following box is checked:

- ☒ was filed on **October 30, 1996**,  
as United States Application Number or PCT International Application Number **08/741,095**; and  
was amended on \_\_\_\_\_ (if applicable).

I hereby state that I have reviewed and understand the contents of the above identified specification, including the claims, as amended by any amendment referred to above.

I acknowledge the duty to disclose information that is material to patentability as defined in 37 C.F.R. § 1.56.

I hereby claim foreign priority benefits under 35 U.S.C. § 119(a)-(d) or § 365(b) of any foreign application(s) for patent or inventor's certificate, or § 365(a) of any PCT international application, which designated at least one country other than the United States listed below, and have also identified below any foreign application for patent or inventor's certificate, or PCT international application having a filing date before that of the application on which priority is claimed.

Prior Foreign Application(s)			Priority Claimed
_____	_____	_____	<input type="checkbox"/> Yes <input type="checkbox"/> No
(Application No.)	(Country)	(Day/Month/Year Filed)	
_____	_____	_____	<input type="checkbox"/> Yes <input type="checkbox"/> No
(Application No.)	(Country)	(Day/Month/Year Filed)	

I hereby claim the benefit under 35 U.S.C. § 119(e) of any United States provisional application(s) listed below.

_____	_____
(Application No.)	(Filing Date)
_____	_____
(Application No.)	(Filing Date)

I hereby claim the benefit under 35 U.S.C. § 120 of any United States application(s), or under § 365(c) of any PCT international application designating the United States, listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States or PCT international application in the manner provided by the first paragraph of 35 U.S.C. § 112, I acknowledge the duty to disclose information that is material to patentability as defined in 37 C.F.R. § 1.56 that became available between the filing date of the prior application and the national or PCT international filing date of this application.

<u>PCT/US95/05058</u>	<u>April 27, 1995</u>	<u>Pending</u>
(Application No.)	(Filing Date)	(Status - patented, pending, abandoned)
<u>08/462,315</u>	<u>June 5, 1995</u>	<u>Pending</u>
(Application No.)	(Filing Date)	(Status - patented, pending, abandoned)
<u>08/462,962</u>	<u>June 5, 1995</u>	<u>Pending</u>
(Application No.)	(Filing Date)	(Status - patented, pending, abandoned)
<u>08/464,595</u>	<u>June 5, 1995</u>	<u>Pending</u>
(Application No.)	(Filing Date)	(Status - patented, pending, abandoned)

I hereby appoint the following attorney(s) and/or agent(s) to prosecute this application and to transact all business in the Patent and Trademark Office connected therewith:

Robert G. Sterne, Esq., Reg. No. 28,912; Edward J. Kessler, Esq., Reg. No. 25,688; Jorge A. Goldstein, Esq., Reg. No. 29,021; Samuel L. Fox, Esq., Reg. No. 30,353; David K.S. Cornwell, Esq., Reg. No. 31,944; Robert W. Esmond, Esq., Reg. No. 32,893; Tracy-Gene G. Durkin, Esq., Reg. No. 32,831; Michele A. Cimbala, Esq., Reg. No. 33,851; Michael B. Ray, Esq., Reg. No. 33,997; Robert E. Sokohl, Esq., Reg. No. 36,013; Eric K. Steffe, Esq., Registration No. 36,688; Andrea G. Reister, Esq., Registration No. 36,253; Daniel N. Yannuzzi, Esq., Registration No. 36,727; Robert H. Benson, Registration No. 30,446; Paul C. Kimball, Registration No. 34,610; and A. Anders Brookes, Registration No. 36,373.



Send Correspondence to:

STERNE, KESSLER, GOLDSTEIN & FOX P.L.L.C.  
1100 New York Avenue, N.W.  
Suite 600  
Washington, D.C. 20005-3934

Direct Telephone Calls to:

(202) 371-2600

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

Full name of sole or first inventor <b>Jian NI</b>	
Inventor's signature 	3/27/97 <sup>x</sup> Date
Residence <b>U.S.A.</b>	
Citizenship <b>China</b>	
Post Office Address <b>5502 Manorfield Road, Rockville, MD 20853</b>	
Full name of second inventor <b>Craig A. ROSEN</b>	
Second inventor's signature 	4/4/97 Date
Residence <b>U.S.A.</b>	
Citizenship <b>U.S.A.</b>	
Post Office Address <b>22400 Rolling Hill Road, Laytonsville, MD 20882</b>	

Full name of third inventor <b>Reiner L. GENTZ</b>	
Third inventor's signature <i>Reiner Gentz</i>	<i>3-27-97</i> <sup>X</sup> Date
Residence <b>U.S.A.</b>	
Citizenship <b>Germany</b>	
Post Office Address <b>13404 Fairland Park Drive, Silver Spring, MD 20904</b>	
Full name of fourth inventor <b>Sally Doreen Patricia LYN</b>	
Fourth inventor's signature	Date
Residence <b>U.S.A.</b>	
Citizenship <b>Canada</b>	
Post Office Address <b>134-B Hampton Court, West Chester, PA 19380</b>	
Full name of fifth inventor <b>Mark Robert HURLE</b>	
Fifth inventor's signature	Date
Residence <b>U.S.A.</b>	
Citizenship <b>U.S.A.</b>	
Post Office Address <b>105 Stonybrook Drive, Norristown, PA 19403</b>	

SEQUENCE LISTING

(1) GENERAL INFORMATION:

- (i) APPLICANT: Ni, Jian  
Rosen, Craig A.  
Gentz, Reiner L.  
Lyn, Sally Doreen Patricia  
Hurle, Mark Robert
- (ii) TITLE OF INVENTION: Human Tumor Necrosis Factor  
Receptor-Like 2
- (iii) NUMBER OF SEQUENCES: 24
- (iv) CORRESPONDENCE ADDRESS:  
(A) ADDRESSEE: Sterne, Kessler, Goldstein & Fox, P.L.L.C.  
(B) STREET: 1100 New York Ave, Suite 600  
(C) CITY: Washington  
(D) STATE: DC  
(E) COUNTRY: USA  
(F) ZIP: 20005-3934
- (v) COMPUTER READABLE FORM:  
(A) MEDIUM TYPE: Floppy disk  
(B) COMPUTER: IBM PC compatible  
(C) OPERATING SYSTEM: PC-DOS/MS-DOS  
(D) SOFTWARE: PatentIn Release #1.0, Version #1.30
- (vi) CURRENT APPLICATION DATA:  
(A) APPLICATION NUMBER:  
(B) FILING DATE:  
(C) CLASSIFICATION:
- (vii) PRIOR APPLICATION DATA:  
(A) APPLICATION NUMBER: US 08/464,595  
(B) FILING DATE: 05-JUN-1995
- (viii) PRIOR APPLICATION DATA:  
(A) APPLICATION NUMBER: US 08/462,962  
(B) FILING DATE: 05-JUN-1995
- (ix) PRIOR APPLICATION DATA:  
(A) APPLICATION NUMBER: US 08/462,315  
(B) FILING DATE: 05-JUN-1995
- (x) PRIOR APPLICATION DATA:  
(A) APPLICATION NUMBER: WO PCT/US95/05058  
(B) FILING DATE: 27-APR-1995
- (xi) ATTORNEY/AGENT INFORMATION:  
(A) NAME: Steffe, Eric K.  
(B) REGISTRATION NUMBER: 36,688  
(C) REFERENCE/DOCKET NUMBER: 1488.0770004/EKS/SGW
- (xii) TELECOMMUNICATION INFORMATION:  
(A) TELEPHONE: 202-271-2600  
(B) TELEFAX: 202-271-2540

06904E60 06290

(2) INFORMATION FOR SEQ ID NO:1:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 1704 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: double  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

- (ix) FEATURE:  
 (A) NAME/KEY: CDS  
 (B) LOCATION: 265..1113

- (ix) FEATURE:  
 (A) NAME/KEY: sig\_peptide  
 (B) LOCATION: 265..372

- (ix) FEATURE:  
 (A) NAME/KEY: mat\_peptide  
 (B) LOCATION: 373..1113

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

GCACGAGCTG CCTCCCGCAG GCGCCACCTG TGTCCCCCAG CGCCGCTCCA CCCAGCAGGC	60
CTGAGCCCCT CTCTGCTGCC AGACACCCCC TGCTGCCCAC TCTCCTGCTG CTCGGGTTCT	120
GAGGCACAGC TTGTCACACC GAGGCGGATT CTCTTTCTCT TTCTCTTTCT CTTCTGGCCC	180
ACAGCCGCAG CAATGGCGCT GAGTTCCTCT GCTGGAGTTC ATCCTGCTAG CTGGGTTCCC	240
GAGCTGCCGG TCTGAGCCTG AGGC ATG GAG CCT CCT GGA GAC TGG GGG CCT	291
Met Glu Pro Pro Gly Asp Trp Gly Pro	
-36 -35 -30	
CCT CCC TGG AGA TCC ACC CCC AAA ACC GAC GTC TTG AGG CTG GTG CTG	339
Pro Pro Trp Arg Ser Thr Pro Lys Thr Asp Val Leu Arg Leu Val Leu	
-25 -20 -15	
TAT CTC ACC TTC CTG GGA GCC CCC TGC TAC GCC CCA GCT CTG CCG TCC	387
Tyr Leu Thr Phe Leu Gly Ala Pro Cys Tyr Ala Pro Ala Leu Pro Ser	
-10 -5 1 5	
TGC AAG GAG GAC GAG TAC CCA GTG GGC TCC GAG TGC TGC CCC AAG TGC	435
Cys Lys Glu Asp Glu Tyr Pro Val Gly Ser Glu Cys Cys Pro Lys Cys	
10 15 20	
AGT CCA GGT TAT CGT GTG AAG GAG GCC TGC GGG GAG CTG ACG GGC ACA	483
Ser Pro Gly Tyr Arg Val Lys Glu Ala Cys Gly Glu Leu Thr Gly Thr	
25 30 35	
GTG TGT GAA CCC TGC CCT CCA GGC ACC TAC ATT GCC CAC CTC AAT GGC	531
Val Cys Glu Pro Cys Pro Pro Gly Thr Tyr Ile Ala His Leu Asn Gly	
40 45 50	

555230-06904660

CTA AGC AAG TGT CTG CAG TGC CAA ATG TGT GAC CCA GCC ATG GGC CTG	579
Leu Ser Lys Cys Leu Gln Cys Gln Met Cys Asp Pro Ala Met Gly Leu	
55 60 65	
CGC GCG AGC CGG AAC TGC TCC AGG ACA GAG AAC GCC GTG TGT GGT TGC	627
Arg Ala Ser Arg Asn Cys Ser Arg Thr Glu Asn Ala Val Cys Gly Cys	
70 75 80 85	
AGC CCA GGC CAC TTC TGC ATC GTC CAG GAC GGG GAC CAC TGC GCC GCG	675
Ser Pro Gly His Phe Cys Ile Val Gln Asp Gly Asp His Cys Ala Ala	
90 95 100	
TGC CGC GCT TAC GCC ACC TCC AGC CCG GGC CAG AGG GTG CAG AAG GGA	723
Cys Arg Ala Tyr Ala Thr Ser Ser Pro Gly Gln Arg Val Gln Lys Gly	
105 110 115	
GGC ACC GAG AGT CAG GAC ACC CTG TGT CAG AAC TGC CCC CCG GGG ACC	771
Gly Thr Glu Ser Gln Asp Thr Leu Cys Gln Asn Cys Pro Pro Gly Thr	
120 125 130	
TTC TCT CCC AAT GGG ACC CTG GAG GAA TGT CAG CAC CAG ACC AAG TGC	819
Phe Ser Pro Asn Gly Thr Leu Glu Glu Cys Gln His Gln Thr Lys Cys	
135 140 145	
AGC TGG CTG GTG ACG AAG GCC GGA GCT GGG ACC AGC AGC TCC CAC TGG	867
Ser Trp Leu Val Thr Lys Ala Gly Ala Gly Thr Ser Ser Ser His Trp	
150 155 160 165	
GTA TGG TGG TTT CTC TCA GGG AGC CTC GTC ATC GTC ATT GTT TGC TCC	915
Val Trp Trp Phe Leu Ser Gly Ser Leu Val Ile Val Ile Val Cys Ser	
170 175 180	
ACA GTT GGC CTA ATC ATA TGT GTG AAA AGA AGA AAG CCA AGG GGT GAT	963
Thr Val Gly Leu Ile Ile Cys Val Lys Arg Arg Lys Pro Arg Gly Asp	
185 190 195	
GTA GTC AAG GTG ATC GTC TCC GTC CAG CGG AAA AGA CAG GAG GCA GAA	1011
Val Val Lys Val Ile Val Ser Val Gln Arg Lys Arg Gln Glu Ala Glu	
200 205 210	
GGT GAG GCC ACA GTC ATT GAG GCC CTG CAG GCC CCT CCG GAC GTC ACC	1059
Gly Glu Ala Thr Val Ile Glu Ala Leu Gln Ala Pro Pro Asp Val Thr	
215 220 225	
ACG GTG GCC GTG GAG GAG ACA ATA CCC TCA TTC ACG GGG AGG AGC CCA	1107
Thr Val Ala Val Glu Glu Thr Ile Pro Ser Phe Thr Gly Arg Ser Pro	
230 235 240 245	
AAC CAC TGACCCACAG ACTCTGCACC CCGACGCCAG AGATACCTGG AGCGACGGCT	1163
Asn His	
GAATGAAAGA GGCTGTCCAC CTGGCGGAAC CACCGGAGCC CGGAGGCTTG GGGGCTCCAC	1223
CCTGGACTGG CTTCGGTCTC CTCCAGTGGA GGGAGAGGTG GCGCCCTGC TGGGGTAGAG	1283
CTGGGGACGC CACGTGCCAT TCCCATGGGC CAGTGAGGGC CTGGGGCCTC TGTTCCTGCTG	1343
TGGCCTGAGC TCCCCAGAGT CCTGAGGAGG AGCGCCAGTT GCCCCTCGCT CACAGACCAC	1403

00904E60



(2) INFORMATION FOR SEO ID NO:2:

(A) LENGTH: 283 amino acids

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

Met -36	Glu -35	Pro	Pro	Gly	Asp	Trp -30	Gly	Pro	Pro	Pro	Trp -25	Arg	Ser	Thr	Pro
Lys -20	Thr	Asp	Val	Leu	Arg -15	Leu	Val	Leu	Tyr	Leu -10	Thr	Phe	Leu	Gly	Ala -5
Pro	Cys	Tyr	Ala	Pro 1	Ala	Leu	Pro	Ser 5	Cys	Lys	Glu	Asp	Glu 10	Tyr	Pro
Val	Gly	Ser 15	Glu	Cys	Cys	Pro	Lys 20	Cys	Ser	Pro	Gly	Tyr 25	Arg	Val	Lys
Glu	Ala 30	Cys	Gly	Glu	Leu	Thr 35	Gly	Thr	Val	Cys	Glu 40	Pro	Cys	Pro	Pro
Gly 45	Thr	Tyr	Ile	Ala	His 50	Leu	Asn	Gly	Leu	Ser 55	Lys	Cys	Leu	Gln	Cys 60
Gln	Met	Cys	Asp	Pro 65	Ala	Met	Gly	Leu	Arg 70	Ala	Ser	Arg	Asn 75	Cys	Ser
Arg	Thr	Glu	Asn 80	Ala	Val	Cys	Gly	Cys 85	Ser	Pro	Gly	His 90	Phe	Cys	Ile
Val	Gln	Asp 95	Gly	Asp	His	Cys 100	Ala	Ala	Cys	Arg	Ala	Tyr 105	Ala	Thr	Ser
Ser	Pro 110	Gly	Gln	Arg	Val	Gln 115	Lys	Gly	Gly	Thr	Glu 120	Ser	Gln	Asp	Thr
Leu 125	Cys	Gln	Asn	Cys	Pro 130	Pro	Gly	Thr	Phe	Ser 135	Pro	Asn	Gly	Thr	Leu 140
Glu	Glu	Cys	Gln	His 145	Gln	Thr	Lys	Cys	Ser 150	Trp	Leu	Val	Thr	Lys 155	Ala

Gly Ala Gly Thr Ser Ser Ser His Trp Val Trp Trp Phe Leu Ser Gly  
160 165 170  
Ser Leu Val Ile Val Ile Val Cys Ser Thr Val Gly Leu Ile Ile Cys  
175 180 185  
Val Lys Arg Arg Lys Pro Arg Gly Asp Val Val Lys Val Ile Val Ser  
190 195 200  
Val Gln Arg Lys Arg Gln Glu Ala Glu Gly Glu Ala Thr Val Ile Glu  
205 210 215 220  
Ala Leu Gln Ala Pro Pro Asp Val Thr Thr Val Ala Val Glu Glu Thr  
225 230 235  
Ile Pro Ser Phe Thr Gly Arg Ser Pro Asn His  
240 245

(2) INFORMATION FOR SEQ ID NO:3:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 281 amino acids
  - (B) TYPE: amino acid
  - (C) STRANDEDNESS: not relevant
  - (D) TOPOLOGY: not relevant

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

Met Val Ser Leu Pro Arg Leu Cys Ala Leu Trp Gly Cys Leu Leu Thr  
1 5 10 15  
Ala Val His Leu Gly Gln Cys Val Thr Cys Ser Asp Lys Gln Tyr Leu  
20 25 30  
His Asp Gly Gln Cys Cys Asp Leu Cys Gln Pro Gly Ser Arg Leu Thr  
35 40 45  
Ser His Cys Thr Ala Leu Glu Lys Thr Gln Cys His Pro Cys Asp Ser  
50 55 60  
Gly Glu Phe Ser Ala Gln Trp Asn Arg Glu Ile Arg Cys His Gln His  
65 70 75 80  
Arg His Cys Glu Pro Asn Gln Gly Leu Arg Val Lys Lys Glu Gly Thr  
85 90 95  
Ala Glu Ser Asp Thr Val Cys Thr Cys Lys Glu Gly Gln His Cys Thr  
100 105 110  
Ser Lys Asp Cys Glu Ala Cys Ala Gln His Thr Pro Cys Ile Pro Gly  
115 120 125  
Phe Gly Val Met Glu Met Ala Thr Glu Thr Thr Asp Thr Val Cys His  
130 135 140

655250" 06904660

(2) INFORMATION FOR SEQ ID NO:4:

(A) LENGTH: 2692 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: double  
(D) TOPOLOGY: linear

(ix) FEATURE:

(A) NAME/KEY: CDS  
(B) LOCATION: 373..927

(ix) FEATURE:

```
(A) NAME/KEY: sig_peptide
(B) LOCATION: 373..480
```

(ix) FEATURE:

```
(A) NAME/KEY: mat_peptide
(B) LOCATION: 481..927
```

(xi) SEQUENCE DESCRIPTION: SEO ID NO:4:

CCCCCTTCTA	CAGGAACCC	GGAGTGGACT	GGAACGGTGC	AGGGGGAGAA	CTCGCCCCCTC	60
CCATCGGGCG	CCTCCTTCAT	ACCGGCCCTT	CCCCTCGGCT	TTGCCTGGAC	AGCTCCTGCC	120
TCAGGCAGCG	CCACCTGTGT	CGCCAGCGC	CGCTCCACCC	AGCAGGCCCTG	AGCCCCCTCTC	180

TGCTGCCAGA CACCCCTGCTG TGCCCACTAC TCCTGCTGCT CGGGTTCTGA GGCACAGCTT 240  
 GTCACACCGA GCGGATTCT CTTTCTCTTT CTCTTTCTCT TCTGGCCAC AGCCGCAGCA 300  
 ATGGCGCTGA GTTCCTCTGC TGGAGTTCAT CCTGCTAGCT GGGTTCCCGA GCTGCCGGTC 360  
 TGAGCCTGAG TC ATG GAG CCT CCT GGA GAC TGG GGG CCT CCT CCC TGG 408  
 Met Glu Pro Pro Gly Asp Trp Gly Pro Pro Pro Trp  
 -36 -35 -30 -25  
 AGA TCC ACC CCC AGA ACC GAC GTC TTG AGG CTG GTG CTG TAT CTC ACC 456  
 Arg Ser Thr Pro Arg Thr Asp Val Leu Arg Leu Val Leu Tyr Leu Thr  
 -20 -15 -10  
 TTC CTG GGA GCC CCC TGC TAC GCC CCA GCT CTG CCG TCC TGC AAG GAG 504  
 Phe Leu Gly Ala Pro Cys Tyr Ala Pro Ala Leu Pro Ser Cys Lys Glu  
 -5 1 5  
 GAC GAG TAC CCA GTG GGC TCC GAG TGC TGC CCC AAG TGC AGT CCA GGT 552  
 Asp Glu Tyr Pro Val Gly Ser Glu Cys Cys Pro Lys Cys Ser Pro Gly  
 10 15 20  
 TAT CGT GTG AAG GAG GCC TGC GGG GAG CTG ACG GGC ACA GTG TGT GAA 600  
 Tyr Arg Val Lys Glu Ala Cys Gly Glu Leu Thr Gly Thr Val Cys Glu  
 25 30 35 40  
 CCC TGC CCT CCA GGC ACC TAC ATT GCC CAC CTC AAT GGC CTA AGC AAG 648  
 Pro Cys Pro Pro Gly Thr Tyr Ile Ala His Leu Asn Gly Leu Ser Lys  
 45 50 55  
 TGT CTG CAG TGC CAA ATG TGT GAC CCA GAT ATT GGT TCC CCC TGT GAC 696  
 Cys Leu Gln Cys Gln Met Cys Asp Pro Asp Ile Gly Ser Pro Cys Asp  
 60 65 70  
 CTC AGG GGA AGA GGT CAC CTG GAG GCT GGT GCC CAC CTG AGT CCA GGC 744  
 Leu Arg Gly Arg Gly His Leu Glu Ala Gly Ala His Leu Ser Pro Gly  
 75 80 85  
 AGA CAG AAA GGG GAA CCA GAC CCA GAG GTG GCC TTT GAG TCA CTG AGC 792  
 Arg Gln Lys Gly Glu Pro Asp Pro Glu Val Ala Phe Glu Ser Leu Ser  
 90 95 100  
 GCA GAG CCT GTC CAT GCG GCC AAC GGC TCT GTC CCC TTG GAG CCT CAT 840  
 Ala Glu Pro Val His Ala Ala Asn Gly Ser Val Pro Leu Glu Pro His  
 105 110 115 120  
 GCC AGG CTC AGC ATG GCC AGT GCT CCC TGC GGC CAG GCA GGA CTG CAC 888  
 Ala Arg Leu Ser Met Ala Ser Ala Pro Cys Gly Gln Ala Gly Leu His  
 125 130 135  
 CTG CGG GAC AGG GCT GAC GGC ACA CCT GGG GGC AGG GCC TGAGCCTACA 937  
 Leu Arg Asp Arg Ala Asp Gly Thr Pro Gly Gly Arg Ala  
 140 145  
 GGGAGGCACA GGGCAGGTGG GCTAGCCATG AACAGAAGAG GAAGCTGGAG TGCTTTGGGG 997  
 GTTCATGCAT GTAGGCTGGG ATTTGGGGCT CACACCTCAA CCTGCATGCC CAGTTCCATG 1057  
 CCCCTCCCTT CTTGTGAAAG CACCTGTCTA CTTGGGCTGA GGATGTGGGG GCACAGGTGG 1117

00940690 06904E60

CAGGTGAGGC	TGCCCTCAGG	AGGGGCCCAG	GCCCAGCTTG	TACCCACCT	CCACCAGTAC	1177
CTGAAGAAGT	GGGGCTCTCA	CCCTACCTGC	CTCTGCCATT	GGAATGGCCT	GGTTTGCACA	1237
GATGGGAAAC	CCGTTTGAGG	GGTGGGTGTC	TGGGTGGGCA	CGTGGGGCGA	GGACCTGCCT	1297
GAGGGACCCT	GCCCTGGAAC	TGACAGTGCA	AGCTCGGCGT	CCTGCCCCATC	TGGGCAGAAG	1357
GCTGGTTTCT	CCCATCAACG	AAGCCCTCCC	AGGACCTTCC	TGCAAGCCCT	CGTCCCACAC	1417
GCAGCTCTGC	CGTCCCTTGG	TGTCCCTCCC	GGCCTCAGGT	CCTCCATGCT	GGGTACCTCT	1477
GGGCACCTCG	TTTGGCTGAG	CCAGGGGTTC	AGCCTGGCAG	GGCGCCCTGG	CAGCAGTCCT	1537
TGGCCTGTGG	ATGCTGTCCT	GGCCTGTGGA	TGGTGTCCCG	CCCTCCACGT	ACCCCTCTCA	1597
CCCCCTCCTC	TTGGACTCCA	GCCATGGGCC	TGCGCGCGAG	CCGGAAGTGC	TCCAGGACAG	1657
AGAACGCCGT	GTGTGGCTGC	AGCCCAGGCC	ACTTCTGCAT	CGTCCAGGAC	GGGGACCACT	1717
GCGCCGCGTG	CCGCGCTTAC	GCCACCTCCA	GCCC GGCCA	GAGGGTGCAG	AAGGGAGGCA	1777
CCGAGAGTCA	GGACACCCTG	TGTCAGAACT	GCCCCCGGG	GACCTTCTCT	CCCAATGGGA	1837
CCCTGGAGGA	ATGTCAGCAC	CAGACCAATT	GGCCTAATCA	TATGTGTGAA	AAGAAGAAAG	1897
CCAAGGGGTG	AGCACACGGT	GGCCCCATCA	GGGTTTATGT	CCCCAGCCGT	CACCTCTTGG	1957
AGCTCTGTCA	CCCCAAGCCT	GGGAGGTGGC	CCCAGAGCTT	TTCCAGGATC	CGCGGCTCCT	2017
CCCAGGGCAG	CCACTGCAGG	CTGGGGCAGG	TGTATGTAGT	CAAGGTGATC	GTCTCCGTCC	2077
AGCGGTAAAA	GACAGGAGGC	AGAAGGTGAG	GCCACAGTCA	TTGAGCCCTG	CAGGCCCCCTC	2137
CGGACGTCAC	CACGGTGGCC	GTGGAGGAGA	CAATACCCTC	ATTCACGGGG	AGGAGCCCCAA	2197
ACCACTGACC	CACAGACTCT	GCACCCCGAC	GCCAGAGATA	CCTGGAGAGA	CGGCTGCTGA	2257
TAGAGGCTGT	CCACCTGGCG	AAACCACCGG	AGCCCGGAGG	CTTGGGGGCT	CCGCCCTGGG	2317
CTGGTTTCCG	TCTCTCCAG	TGGAGGGAGA	GGTGGTGCCC	CTGCTGGTGG	TAGAGCTGGG	2377
GACGCCACGT	GCCATTCCCA	TGGTTCAGTG	AGGGGCTGGT	GGCCTCTGTT	CTGCTGTGGC	2437
CTGAGCTCCC	CAGAGTCCTG	AGGAGGAGCC	CCAGTTGCCC	CTCGCTCACA	GACCACACAC	2497
CCAGCCCTCC	TGGGCCAACC	CAGAGGCCCC	TTCAGACCCC	AGCTGTCTGC	GCGTCTGACT	2557
CTTGTGGCCT	CAGCAGGACA	GGCCCCGGGC	ACTGCCTCAC	AGCCAAGGCT	GGAATGGGTT	2617
GGCTGCAGTG	TGGTGTTTAG	TGGATACCAC	ATCGGAAGTG	ATTTTCTAAA	AATTGGATTT	2677
GAATTGCGAA	AAAAA					2692

(2) INFORMATION FOR SEQ ID NO:5:

(1) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 185 amino acids

06940690-062999

(B) TYPE: amino acid  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(x1) SEQUENCE DESCRIPTION: SEQ ID NO:5:

```

Met Glu Pro Pro Gly Asp Trp Gly Pro Pro Pro Trp Arg Ser Thr Pro
-36 -35                               -30           -25

Arg Thr Asp Val Leu Arg Leu Val Leu Tyr Leu Thr Phe Leu Gly Ala
-20                               -15           -10           -5

Pro Cys Tyr Ala Pro Ala Leu Pro Ser Cys Lys Glu Asp Glu Tyr Pro
                               1             5             10

Val Gly Ser Glu Cys Cys Pro Lys Cys Ser Pro Gly Tyr Arg Val Lys
                               15             20             25

Glu Ala Cys Gly Glu Leu Thr Gly Thr Val Cys Glu Pro Cys Pro Pro
                               30             35             40

Gly Thr Tyr Ile Ala His Leu Asn Gly Leu Ser Lys Cys Leu Gln Cys
45                               50           55           60

Gln Met Cys Asp Pro Asp Ile Gly Ser Pro Cys Asp Leu Arg Gly Arg
                               65           70           75

Gly His Leu Glu Ala Gly Ala His Leu Ser Pro Gly Arg Gln Lys Gly
                               80           85           90

Glu Pro Asp Pro Glu Val Ala Phe Glu Ser Leu Ser Ala Glu Pro Val
                               95           100          105

His Ala Ala Asn Gly Ser Val Pro Leu Glu Pro His Ala Arg Leu Ser
110                               115          120

Met Ala Ser Ala Pro Cys Gly Gln Ala Gly Leu His Leu Arg Asp Arg
125                               130          135          140

Ala Asp Gly Thr Pro Gly Gly Arg Ala
                               145

```

(2) INFORMATION FOR SEQ ID NO:6:

(i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 248 amino acids  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: not relevant  
 (D) TOPOLOGY: not relevant

(ii) MOLECULE TYPE: protein

(x1) SEQUENCE DESCRIPTION: SEQ ID NO:6:

Met Ala Pro Val Ala Val Trp Ala Ala Leu Ala Val Gly Leu Glu Leu

05940690-062999

1		5		10		15									
Trp	Ala	Ala	Ala	His	Ala	Leu	Pro	Ala	Gln	Val	Ala	Phe	Thr	Pro	Tyr
		20						25					30		
Ala	Pro	Glu	Pro	Gly	Ser	Thr	Cys	Arg	Leu	Arg	Glu	Tyr	Tyr	Asp	Gln
		35					40					45			
Thr	Ala	Gln	Met	Cys	Cys	Ser	Lys	Cys	Ser	Pro	Gly	Gln	His	Ala	Lys
	50					55					60				
Val	Phe	Cys	Thr	Lys	Thr	Ser	Asp	Thr	Val	Cys	Asp	Ser	Cys	Glu	Asp
65					70					75					80
Ser	Thr	Tyr	Thr	Gln	Leu	Trp	Asn	Trp	Val	Pro	Glu	Cys	Leu	Ser	Cys
				85					90					95	
Gly	Ser	Arg	Cys	Ser	Ser	Asp	Gln	Val	Glu	Thr	Gln	Ala	Cys	Thr	Arg
			100					105					110		
Glu	Gln	Asn	Arg	Ile	Cys	Thr	Cys	Arg	Pro	Gly	Trp	Tyr	Cys	Ala	Leu
		115					120					125			
Ser	Lys	Gln	Glu	Gly	Cys	Arg	Leu	Cys	Ala	Pro	Leu	Arg	Lys	Cys	Arg
	130					135					140				
Pro	Gly	Phe	Gly	Val	Ala	Arg	Pro	Gly	Thr	Glu	Thr	Ser	Asp	Val	Val
145					150					155					160
Cys	Lys	Pro	Cys	Ala	Pro	Gly	Thr	Phe	Ser	Asn	Thr	Thr	Ser	Ser	Thr
				165					170					175	
Asp	Ile	Cys	Arg	Pro	His	Gln	Ile	Cys	Asn	Val	Val	Ala	Ile	Pro	Gly
			180					185					190		
Asn	Ala	Ser	Met	Asp	Ala	Val	Cys	Thr	Ser	Thr	Ser	Pro	Thr	Arg	Ser
		195					200					205			
Met	Ala	Pro	Gly	Ala	Val	His	Leu	Pro	Gln	Pro	Val	Ser	Thr	Arg	Ser
	210					215					220				
Gln	His	Thr	Gln	Pro	Thr	Pro	Glu	Pro	Ser	Thr	Ala	Pro	Ser	Thr	Ser
225					230					235					240
Phe	Leu	Leu	Pro	Met	Gly	Pro	Ser								
				245											

(2) INFORMATION FOR SEQ ID NO:7:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 2637 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: double
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(ix) FEATURE:

09340690 06904E60

—

—

AAAGCTCGGG	CTCCACCGGG	GACGACCGCT	CCTAGAAACT	GAGTGGTATC	CCCCGGGCCT		60
GCAGGAATTC	CAACCTGCCT	GAAGGGACCC	TGCCCTGGAA	CTGACAGTGC	AAGCTCGGCG		120
TCCTGCCCAT	CTGGGAAGAA	GGCTGGTTTC	TCCCATCAAC	GAAGCCCTCC	CAGGACCTTC		180
CTGCAAGCCC	TCGTCCCACA	CGCAGCTCTG	CCGTCCCTTG	GTGTCCCTCC	CGGCCTCAGG		240
TCCTCC	ATG CTG GGT ACC TCT GGG CAC CTC GTT TGG CTG AGC CAG GGG						288
	Met Leu Gly Thr Ser Gly His Leu Val Trp Leu Ser Gln Gly						
	150		155		160		
TTC AGC CTG GCA GGG CGC CCT GGC AGC AGT CCT TGG CCT GTG GAT GCT							336
Phe Ser Leu Ala Gly Arg Pro Gly Ser Ser Pro Trp Pro Val Asp Ala							
	165		170		175		
GTC CTG GCC TGT GGA TGG TGT CCC GGC CTC CAC GTA CCC CCT CTC AGC							384
Val Leu Ala Cys Gly Trp Cys Pro Gly Leu His Val Pro Pro Leu Ser							
	180		185		190		195
CCC TCC TCT TGG ACT CCA GCC ATG GGC CTG CGC GCG AGC CGG AAC TGC							432
Pro Ser Ser Trp Thr Pro Ala Met Gly Leu Arg Ala Ser Arg Asn Cys							
		200			205		210
TCC AGG ACA GAG AAC GCC GTG TGT GGC TGC AGC CCA GGC CAC TTC TGC							480
Ser Arg Thr Glu Asn Ala Val Cys Gly Cys Ser Pro Gly His Phe Cys							
		215			220		225
ATC GTC CAG GAC GGG GAC CAC TGC GCC GCG TGC CGC GCT TAC GCC ACC							528
Ile Val Gln Asp Gly Asp His Cys Ala Ala Cys Arg Ala Tyr Ala Thr							
		230			235		240
TCC AGC CCG GGC CAG AGG GTG CAG AAG GGA GGC ACC GAG AGT CAG GAC							576
Ser Ser Pro Gly Gln Arg Val Gln Lys Gly Gly Thr Glu Ser Gln Asp							
		245			250		255
ACC CTG TGT CAG AAC TGC CCC CGG GGA CCT TCT CTC CCA ATG GGA CCC							624
Thr Leu Cys Gln Asn Cys Pro Arg Gly Pro Ser Leu Pro Met Gly Pro							
		260			265		270
							275
TGG AGG AAT GTC AGC ACC AGA CCA AGT AAG TGAACCCGGG GGAGGCCAGC							674
Trp Arg Asn Val Ser Thr Arg Pro Ser Lys							
		280			285		
TCTGTGCCCT	GGGGAGGGGG	CTCCACGTTG	CTTCCCTGGG	AGATGACCGT	CTTCTCCAGC		734
AGAAAGGTTG	AAGGTCCCAC	CCTGAGCGGC	ACCCTGGTCA	CATGCCTGCG	TCCAGGAGAG		794
CTGCAGGGTG	AAGCCTGTGT	GCCCCAGATA	ACCCCTTCCA	TGGGCCCAGA	CAAAGCCTCA		854
TCAGATCTGA	GCTTCCTGGA	GGCTCAGGAT	GGGCCTTCCC	AGAAGCAGGC	CCAGAGGGAG		914
GCTGCCTCCA	GATCCCTGT	CCCCTGGGGC	TGTGGGTGTC	CCTGAATGTC	AGGGCCATGG		974



(2) INFORMATION FOR SEQ ID NO:8:

(A) LENGTH: 136 amino acids  
(B) TYPE: amino acid  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

Met	Leu	Gly	Thr	Ser	Gly	His	Leu	Val	Trp	Leu	Ser	Gln	Gly	Phe	Ser
1				5					10					15	
Leu	Ala	Gly	Arg	Pro	Gly	Ser	Ser	Pro	Trp	Pro	Val	Asp	Ala	Val	Leu
			20					25					30		
Ala	Cys	Gly	Trp	Cys	Pro	Gly	Leu	His	Val	Pro	Pro	Leu	Ser	Pro	Ser
		35					40					45			
Ser	Trp	Thr	Pro	Ala	Met	Gly	Leu	Arg	Ala	Ser	Arg	Asn	Cys	Ser	Arg
	50					55					60				
Thr	Glu	Asn	Ala	Val	Cys	Gly	Cys	Ser	Pro	Gly	His	Phe	Cys	Ile	Val
65					70					75					80
Gln	Asp	Gly	Asp	His	Cys	Ala	Ala	Cys	Arg	Ala	Tyr	Ala	Thr	Ser	Ser
				85					90					95	
Pro	Gly	Gln	Arg	Val	Gln	Lys	Gly	Gly	Thr	Glu	Ser	Gln	Asp	Thr	Leu
			100					105					110		
Cys	Gln	Asn	Cys	Pro	Arg	Gly	Pro	Ser	Leu	Pro	Met	Gly	Pro	Trp	Arg
		115					120					125			
Asn	Val	Ser	Thr	Arg	Pro	Ser	Lys								
130						135									

(2) INFORMATION FOR SEQ ID NO:9:

(A) LENGTH: 198 amino acids  
(B) TYPE: amino acid  
(C) STRANDEDNESS: not relevant  
(D) TOPOLOGY: not relevant

(11) MOLECULE TYPE: protein

(x1) SEQUENCE DESCRIPTION: SEQ ID NO:9:

Met	Ala	Pro	Val	Ala	Val	Trp	Ala	Ala	Leu	Ala	Val	Gly	Leu	Glu	Leu
1				5					10					15	
Trp	Ala	Ala	Ala	His	Ala	Leu	Pro	Ala	Gln	Val	Ala	Phe	Thr	Pro	Tyr
			20					25					30		
Ala	Pro	Glu	Pro	Gly	Ser	Thr	Cys	Arg	Leu	Arg	Glu	Tyr	Tyr	Asp	Gln
		35					40					45			

Thr Ala Gln Met Cys Cys Ser Lys Cys Ser Pro Gly Gln His Ala Lys  
50 55 60

Val Phe Cys Thr Lys Thr Ser Asp Thr Val Cys Asp Ser Cys Glu Asp  
65 70 75 80

Ser Thr Tyr Thr Gln Leu Trp Asn Trp Val Pro Glu Cys Leu Ser Cys  
85 90 95

Gly Ser Arg Cys Ser Ser Asp Gln Val Glu Thr Gln Ala Cys Thr Arg  
100 105 110

Glu Gln Asn Arg Ile Cys Thr Cys Arg Pro Gly Trp Tyr Cys Ala Leu  
115 120 125

Ser Lys Gln Glu Gly Cys Arg Leu Cys Ala Pro Leu Arg Lys Cys Arg  
130 135 140

Pro Gly Phe Gly Val Ala Arg Pro Gly Thr Glu Thr Ser Asp Val Val  
145 150 155 160

Cys Lys Pro Cys Ala Pro Gly Thr Phe Ser Asn Thr Thr Ser Ser Thr  
165 170 175

Asp Ile Cys Arg Pro His Gln Ile Cys Asn Val Val Ala Ile Pro Gly  
180 185 190

Asn Ala Ser Met Asp Ala  
195

(2) INFORMATION FOR SEQ ID NO:10:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 154 amino acids  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: not relevant  
 (D) TOPOLOGY: not relevant

(ii) MOLECULE TYPE: protein

(x1) SEQUENCE DESCRIPTION: SEQ ID NO:10:

Val Cys Pro Gln Gly Lys Tyr Ile His Pro Gln Asn Asn Ser Ile Cys  
1 5 10 15

Cys Thr Lys Cys His Lys Gly Thr Tyr Leu Tyr Asn Asp Cys Pro Gly  
20 25 30

Pro Gly Gln Asp Thr Asp Cys Arg Glu Cys Glu Ser Gly Ser Phe Thr  
35 40 45

Ala Ser Glu Asn His Leu Arg His Cys Leu Ser Cys Ser Lys Cys Arg  
50 55 60

Lys Glu Met Gly Gln Val Glu Ile Ser Ser Cys Thr Val Asp Arg Asp  
65 70 75 80

00340690-0699

(2) INFORMATION FOR SEO ID NO:11:

(11) MOLECULE TYPE: protein

Thr 1	Cys	Arg	Leu	Arg 5	Glu	Tyr	Tyr	Asp	Gln 10	Thr	Ala	Gln	Met	Cys 15	Cys
Ser	Lys	Cys	Ser 20	Pro	Gly	Gln	His	Ala 25	Lys	Val	Phe	Cys	Thr 30	Lys	Thr
Ser	Asp	Thr 35	Val	Cys	Asp	Ser	Cys 40	Glu	Asp	Ser	Thr	Tyr 45	Thr	Gln	Leu
Trp	Asn 50	Trp	Val	Pro	Glu	Cys 55	Leu	Ser	Cys	Gly	Ser 60	Arg	Cys	Ser	Ser
Asp 65	Gln	Val	Glu	Thr	Gln 70	Ala	Cys	Thr	Arg	Glu 75	Gln	Asn	Arg	Ile	Cys 80
Thr	Cys	Arg	Pro 85	Gly	Trp	Tyr	Cys	Ala	Leu 90	Ser	Lys	Gln	Glu	Gly 95	Cys
Arg	Leu	Cys	Ala 100	Pro	Leu	Arg	Lys	Cys 105	Arg	Pro	Gly	Phe	Gly 110	Val	Ala
Arg	Pro 115	Gly	Thr	Glu	Thr	Ser	Asp 120	Val	Val	Cys	Lys	Pro 125	Cys	Ala	Pro
Gly 130	Thr	Phe	Ser	Asn	Thr	Thr 135	Ser	Ser	Thr	Asp	Ile 140	Cys	Arg	Pro	His
Gln 145	Ile	Cys	Asn	Val	Val 150	Ala	Ile	Pro	Gly	Asn 155	Ala	Ser	Met	Asp	Ala 160

Val Cys Thr

(2) INFORMATION FOR SEQ ID NO:12:

- (i) SEQUENCE CHARACTERISTICS:  
    (A) LENGTH: 163 amino acids  
    (B) TYPE: amino acid  
    (C) STRANDEDNESS: not relevant  
    (D) TOPOLOGY: not relevant

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

Ala Cys Arg Glu Lys Gln Tyr Leu Ile Asn Ser Gln Cys Cys Ser Leu  
1                    5                    10                    15  
Cys Gln Pro Gly Gln Lys Leu Val Ser Asp Cys Thr Glu Pro Thr Glu  
                    20                    25                    30  
Thr Glu Cys Leu Pro Cys Gly Glu Ser Glu Phe Leu Asp Thr Trp Asn  
                    35                    40                    45  
Arg Glu Thr His Cys His Gln His Lys Tyr Cys Asp Pro Asn Leu Gly  
50                    55                    60  
Leu Arg Val Gln Gln Lys Gly Thr Ser Glu Thr Asp Thr Ile Cys Thr  
65                    70                    75                    80  
Cys Glu Glu Gly Trp His Cys Thr Ser Glu Ala Cys Glu Ser Cys Val  
                    85                    90                    95  
Leu His Arg Ser Cys Ser Pro Gly Phe Gly Val Lys Gln Ile Ala Thr  
                    100                    105                    110  
Gly Val Ser Asp Thr Ile Cys Glu Pro Cys Pro Val Gly Phe Phe Ser  
115                    120                    125  
Asn Val Ser Ser Ala Phe Glu Lys Cys His Pro Trp Thr Ser Cys Glu  
130                    135                    140  
Thr Lys Asp Leu Val Val Gln Gln Ala Gly Thr Asn Lys Thr Asp Val  
145                    150                    155                    160  
Val Cys Gly

(2) INFORMATION FOR SEQ ID NO:13:

- (i) SEQUENCE CHARACTERISTICS:  
    (A) LENGTH: 132 amino acids  
    (B) TYPE: amino acid  
    (C) STRANDEDNESS: not relevant  
    (D) TOPOLOGY: not relevant

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:

(2) INFORMATION FOR SEQ ID NO:14:

(11) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:

27

(2) INFORMATION FOR SEQ ID NO:15:

(1) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 30 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(11) MOLECULE TYPE: cDNA

(x1) SEQUENCE DESCRIPTION: SEQ ID NO:15:

CGCAAGCTTA TTGTGGGAGC TGCTGGTCCC

30

(2) INFORMATION FOR SEQ ID NO:16:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 25 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(11) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:16:

CGCGGATCCC GGAGCCCCCT GCTAC

25

(2) INFORMATION FOR SEQ ID NO:17:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 30 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(11) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:17:

CGCGGTACCA TTGTGGGAGC TGCTGGTCCC

30

(2) INFORMATION FOR SEQ ID NO:18:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 34 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(11) MOLECULE TYPE: cDNA

(x1) SEQUENCE DESCRIPTION: SEQ ID NO:18:

GCGCGGATCC ACCATGGAGC CTCCTGGAGA CTGG

34

05904660

(2) INFORMATION FOR SEQ ID NO:19:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 31 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:19:

GCGCGGTACC TCTACCCAG CAGGGGCGCC A

31

(2) INFORMATION FOR SEQ ID NO:20:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 34 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:20:

GCGCGGATCC ACCATGGAGC CTCCTGGAGA CTGG

34

(2) INFORMATION FOR SEQ ID NO:21:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 58 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:21:

GCGCTCTAGA TCAAGCGTAG TCTGGGACGT CGTATGGGTA GTGGTTTGGG CTCCTCCC

58

(2) INFORMATION FOR SEQ ID NO:22:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 34 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear



(xi) SEQUENCE DESCRIPTION: SEQ ID NO:22:

34

(i) SEQUENCE CHARACTERISTICS:

- SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 35 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(x1) SEQUENCE DESCRIPTION: SEQ ID NO:23:

35

(2) INFORMATION FOR SEQ ID NO:24:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 53 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(11) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:24:

53